

8.7: The Acid-Base Properties of Ions and Salts



Skills to Develop

- To understand hydrolysis
- To predict the effect a given ion will have on the pH of a solution

Depending on the acid–base properties of its component ions, a salt can dissolve in water to produce a neutral solution, a basic solution, or an acidic solution.



When a salt such as $NaCl$ dissolves in water, it produces $Na^+_{(aq)}$ and $Cl^-_{(aq)}$ ions. Using a Lewis approach, the Na^+ ion can be viewed as an acid because it is an electron pair acceptor, although its low charge and relatively large radius make it a **very** weak acid ($K_a < 1 \times 10^{-14}$). The Cl^- ion is the conjugate base of the strong acid HCl , so it has essentially no basic character ($K_b < 1 \times 10^{-14}$). Consequently, dissolving $NaCl$ in water has no effect on the pH of a solution, and the solution remains neutral.

Now let's compare this behavior to the behavior of aqueous solutions of potassium cyanide and sodium acetate. Again, the cations (K^+ and Na^+) have essentially no acidic character, but the anions (CN^- and $CH_3CO_2^-$) are weak bases that can react with water because they are the conjugate bases of the weak acids HCN and acetic acid, respectively and they have $K_a > 1 \times 10^{-14}$.



Neither reaction proceeds very far to the right as written because the formation of the weaker acid–base pair is favored. Both HCN and acetic acid are stronger acids than water, and hydroxide is a stronger base than either acetate or cyanide, so in both cases, the equilibrium lies to the left. Nonetheless, each of these reactions generates enough hydroxide ions to produce a basic solution. For example, the pH of a 0.1 M solution of sodium acetate or potassium cyanide at 25°C is 8.8 or 11.1, respectively. From Table 8.7.1 and Figure 8.7.1, we can see that CN^- is a stronger base ($pK_b = 4.79$) than acetate ($pK_b = 9.24$), which is consistent with KCN producing a more basic solution than sodium acetate at the same concentration.


In contrast, the conjugate acid of a weak base should be a weak acid (Equation 8.7.3). For example, ammonium chloride and pyridinium chloride are salts produced by reacting ammonia and pyridine, respectively, with HCl . As you already know, the chloride ion is such a weak base that it does not react with water. In contrast, the cations of the two salts are weak acids that react with water because they have a $K_b > 1 \times 10^{-14}$. The reactions are as follows:



Equation 8.7.3 indicates that H_3O^+ is a stronger acid than either NH_4^+ or $C_5H_5NH^+$, and conversely, ammonia and pyridine are both stronger bases than water. The equilibrium will therefore lie far to the left in both cases, favoring the weaker acid–base pair. The H_3O^+ concentration produced by the reactions is great enough, however, to decrease the pH of the solution significantly: the pH of a 0.10 M solution of ammonium chloride or pyridinium chloride at 25°C is 5.13 or 3.12, respectively. This is consistent with the information shown in Figure 16.2, indicating that the pyridinium ion is more acidic than the ammonium ion.

What happens with aqueous solutions of a salt such as ammonium acetate, where both the cation and the anion can react separately with water to produce an acid and a base, respectively? According to Figure 16.10, the ammonium ion will lower the pH , while according to Equation 8.7.4, the acetate ion will raise the pH . This particular case is unusual, in that the cation is as strong an acid as the anion is a base ($K_a \approx K_b$). Consequently, the two effects cancel, and the solution remains neutral. With salts in which the cation is a stronger acid than the anion is a base, the final solution has a $pH < 7.00$. Conversely, if the cation is a weaker acid than the anion is a base, the final solution has a $pH > 7.00$.

Solutions of simple salts of metal ions can also be acidic, even though a metal ion cannot donate a proton directly to water to produce H_3O^+ . Instead, a metal ion can act as a Lewis acid and interact with water, a Lewis base, by coordinating to a lone pair of electrons on the oxygen atom to form a hydrated metal ion (part (a) in Figure 8.7.1). A water molecule coordinated to a metal ion is more acidic than a free water molecule for two reasons. First, repulsive electrostatic interactions between the positively charged metal ion and the partially positively charged hydrogen atoms of the coordinated water molecule make it easier for the coordinated water to lose a proton.



Second, the positive charge on the Al^{3+} ion attracts electron density from the oxygen atoms of the water molecules, which decreases the electron density in the O–H bonds, as shown in part (b) in Figure 8.7.1. With less electron density between the O atoms and the H atoms, the O–H bonds are weaker than in a free H_2O molecule, making it easier to lose a H^+ ion.




Figure 8.7.1: Effect of a Metal Ion on the Acidity of Water (a) Reaction of the metal ion Al^{3+} with water to form the hydrated metal ion is an example of a Lewis acid–base reaction. (b) The positive charge on the aluminum ion attracts electron density from the oxygen atoms, which shifts electron density away from the O–H bonds. The decrease in electron density weakens the O–H bonds in the water molecules and makes it easier for them to lose a proton.

The magnitude of this effect depends on the following two factors (Figure 8.7.2):

1. The charge on the metal ion. A divalent ion (M^{2+}) has approximately twice as strong an effect on the electron density in a coordinated water molecule as a monovalent ion (M^+) of the same radius.
2. The radius of the metal ion. For metal ions with the same charge, the smaller the ion, the shorter the internuclear distance to the oxygen atom of the water molecule and the greater the effect of the metal on the electron density distribution in the water molecule.





Figure 8.7.2: The Effect of the Charge and Radius of a Metal Ion on the Acidity of a Coordinated Water Molecule. The contours show the electron density on the O atoms and the H atoms in both a free water molecule (left) and water molecules coordinated to Na^+ , Mg^{2+} , and Al^{3+} ions. These contour maps demonstrate that the smallest, most highly charged metal ion (Al^{3+}) causes the greatest decrease in electron density of the O–H bonds of the water molecule. Due to this effect, the acidity of hydrated metal ions increases as the charge on the metal ion increases and its radius decreases.

Thus aqueous solutions of small, highly charged metal ions, such as Al^{3+} and Fe^{3+} , are acidic:



The $[Al(H_2O)_6]^{3+}$ ion has a pK_a of 5.0, making it almost as strong an acid as acetic acid. Because of the two factors described previously, the most important parameter for predicting the effect of a metal ion on the acidity of coordinated water molecules is the charge-to-radius ratio of the metal ion.

Solutions of small, highly charged metal ions in water are acidic.

Reactions such as those discussed in this section, in which a salt reacts with water to give an acidic or basic solution, are often called hydrolysis reactions. Using a separate name for this type of reaction is unfortunate because it suggests that they are somehow different. In fact, hydrolysis reactions are just acid–base reactions in which the acid is a cation or the base is an anion; they obey the same principles and rules as all other acid–base reactions.

A hydrolysis reaction is an acid–base reaction.

Example 8.7.1

Predict whether aqueous solutions of these compounds are acidic, basic, or neutral.

- KNO_3
- $CrBr_3$
- Na_2SO_4

Given: compound

Asked for: acidity or basicity of aqueous solution

Strategy:

- Assess the acid–base properties of the cation and the anion. If the cation is a weak Lewis acid, it will not affect the pH of the solution. If the cation is the conjugate acid of a weak base or a relatively highly charged metal cation, however, it will react with water to produce an acidic solution.
- If the anion is the conjugate base of a strong acid, it will not affect the pH of the solution. If, however, the anion is the conjugate base of a weak acid, the solution will be basic.


Solution:

a

- The K^+ cation has a small positive charge (+1) and a relatively large radius (because it is in the fourth row of the periodic table), so its $K_a < 1 \times 10^{-14}$. It will not hydrolyze water or affect the pH .
- The NO_3^- anion is the conjugate base of a strong acid, so it has essentially no basic character ($K_b < 1 \times 10^{-14}$).
- Hence neither the cation nor the anion will react with water to produce H^+ or OH^- , and the solution will be neutral.

b.

- The Cr^{3+} ion is a relatively highly charged metal cation that should behave similarly to the Al^{3+} ion and form the $[Cr(H_2O)_6]^{3+}$ complex, which will behave as a weak acid:



($K_a > 1 \times 10^{-14}$)

- The Br^- anion is a very weak base (it is the conjugate base of the strong acid HBr), so it does not affect the pH of the solution ($K_b < 1 \times 10^{-14}$).
- Hence the solution will be acidic.

c.

- The Na^+ ion, like the K^+ , is a very weak acid ($K_a < 1 \times 10^{-14}$), so it should not affect the acidity of the solution.
- In contrast, SO_4^{2-} is the conjugate base of HSO_4^- , which is a weak acid. Hence the SO_4^{2-} ion, with a $K_b > 1 \times 10^{-14}$, will react with water as shown in Figure 16.6.
- The solution will be slightly basic.

Exercise 8.7.1

Predict whether aqueous solutions of the following are acidic, basic, or neutral.

- a. KI
- b. $Fe(ClO_4)_2$
- c. $NaHS$

Answer a

neutral

Answer b

acidic

Answer c

basic (due to the reaction of HS^- with water to form H_2S and OH^-)

Summary

A salt can dissolve in water to produce a neutral, a basic, or an acidic solution, depending on whether it contains the conjugate base of a weak acid as the anion (A^-), the conjugate acid of a weak base as the cation (BH^+), or both. Salts that contain small, highly charged metal ions produce acidic solutions in water. The reaction of a salt with water to produce an acidic or a basic solution is called a hydrolysis reaction.

Key Takeaways

- Acid–base reactions always contain two conjugate acid–base pairs.
- Each acid and each base has an associated ionization constant that corresponds to its acid or base strength.

Contributors

Modified by [Tom Neils](#) (Grand Rapids Community College)

8.7: The Acid-Base Properties of Ions and Salts is shared under a [CC BY-NC-SA 4.0](#) license and was authored, remixed, and/or curated by LibreTexts.