

5.11: R factor

The term **R factor** in crystallography is commonly taken to refer to the 'conventional' R factor

$R =$

 ParseError: invalid DekiScript ([click for details](#))

\backslash),

a measure of agreement between the amplitudes of the structure factors calculated from a crystallographic model and those from the original X-ray diffraction data. The R factor is calculated during each cycle of least-squares structure refinement to assess progress. The final R factor is one measure of model quality.

More generally, a variety of R factors may be determined to measure analogous residuals during least-squares optimization procedures. Where the refinement attempts to minimize the deviates of the squares of the structure factors (refinement against F^2), the R factor based on F^2 is used to monitor the progress of refinement:

$R(F^2) =$

 ParseError: invalid DekiScript ([click for details](#))

\backslash).

Likewise, refinement against I can be tracked using the Bragg R factor

$R_B =$

 ParseError: invalid DekiScript ([click for details](#))

\backslash).

Even for refinement against F^2 or I , the 'conventional' R factor may be calculated and quoted as a measure of model quality, in order to compare the resulting quality of models calculated at different times and with different refinement strategies.

The R factor is sometimes described as the **discrepancy index**.

This page titled [5.11: R factor](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Online Dictionary of Crystallography](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.