

1.2: VSEPR Theory and its Utility

Learning Objectives

In this lecture you will learn the following

- Definition of VSEPR.
- Hybridization concept.
- Prediction of shapes and geometries of molecules.
- Bent's rule and its application in understanding the structural parameters.

The Valence Shell Electron Pair Repulsion Theory (VSEPR)

VSEPR theory is an improved and extension of Lewis model but predicts the shapes of polyatomic molecules. This model was first suggested by Nevil Sidgwick and Herbert Powell in 1940 and later improved by Ronald Gillespie and Ronald Nyholm.

Prediction of molecular shapes and geometries was made easy by this model through the following simple steps.

1. Draw the Lewis structure.
2. Count the total number of bonds and lone pairs around the central atom. (Each single bond would involve one pair of electrons).
3. Arrange the bonding pairs and lone pairs in one of the standard geometries to minimize the electron-electron repulsion.
 - i. Lone pair electrons stay closer to the nucleus and also they spread out over a larger space than bond pairs and hence large angles between lone pairs.
 - ii. The repulsion follows the order $LP-LP > LP-BP > BP-BP$.
4. Multiple bonds should be considered as a single bonding region.

Steric numbers:

Another term called steric number is often used in VSEPR theory.

Steric number (SN) = No. of attached atom + No. of lone pairs. Since the lone pair—lone pair repulsions are maximum, the most stable geometry can be obtained by maximizing the distance between steric numbers on the central atom.

Molecular shapes are eventually determined by two parameters: Bond distance, separation between the nuclei of two bonded atoms in a straight line and the bond angle, the angle between any two bonds containing a common atom.

While mentioning the molecular shapes lone pairs may be ignored, however, while defining the geometry both the lone pairs and bond pairs should be considered.

For example: in water molecule the central oxygen atom is in tetrahedral environment with two lone pairs and two O—H bonds (or two bond pairs). The shape of the water molecule is therefore bent (two lone pairs are ignored).

Similarly, in ammonia, the nitrogen atom is in tetrahedral environment with three bonded pairs (three N—H bonds) and one lone pair. The shape of NH_3 molecule is pyramidal.

Predicting the molecular geometries

To begin with, draw the Lewis structure.

Count the number of bonding pairs and lone pairs around the central atom.

Arrange the bonding pairs and the lone pairs in one of the standard geometries thereby minimizing electron—electron repulsion.

Multiple bonds count as a single bonding region.

What is Bent's rule:

More electronegative substituents 'prefer' hybrid orbitals having less s-character, and more electropositive substituents 'prefer'

orbitals having more s-character.

The bond angles in CH_4 , CF_4 and CH_2F_2 can be explained using Bent's rule. While a carbon in CH_4 and CF_4 uses four identical sp^3 hybrids for bonding, in CH_2F_2 the hybrids used are not identical.

The C-F bonds are formed from sp^{3+x} hybrids, with slightly more p -character and less s -character than an sp^3 hybrid, and the hydrogen are bonded by sp^{3-x} hybrids, with slightly less p -character and slightly more s -character. Increasing the amount of p -character in the C-F bonds decreases the F-C-F bond angle, because for bonding by pure p -orbitals the bond angle would be decreased to 90° .

Molecular shapes determined by VSEPR theory


Molecule	Steric Number (Number Electron Pairs) (SN)	Geometry	Example
MA_2	2	Linear	BeCl_2
MA_3	3	Trigonal planar	BF_3
MA_4	4	Tetrahedral	SiF_4
MA_5	5	Trigonal bipyramidal	PF_5
MA_6	6	Octahedral	SF_6
MA_7	7	Pentagonal bipyramidal	IF_7

Molecule	SN	Number of lone pairs	Geometry	shape	Example
MA_2	2	0	Linear		CO_2
MA_3	3	0	Trigonal planar	Trigonal planar	SO_3
	3	1		angular	SO_2
MA_4	4	0	Tetrahedral	Tetrahedral	CH_4
		1		Trigonal pyramidal	NH_3
		2		Angular	H_2O
MA_5	5	0	Trigonal bipyramidal	Trigonal bipyramidal	AsF_5
		1		Seesaw	SF_4
		2		T-shaped	ClF_3
		3		linear	XeF_2
MA_6	6	0	Octahedral	Octahedral	SF_6
		1		Square pyramidal	BrF_5
		2		Square planar	XeF_4

Problems:

1. Draw the structure and depict the geometry around **Se** atoms in $[\text{Se}_3\text{O}_6\text{F}_3]^{3-}$, which is a symmetric ionic molecule with cyclic structure, using VSEPR model.

Solution




Apply VSEPR theory to the structure.

Se has six valence electrons. One Se—F and three Se—O (one terminal and two bridging) will add four more electrons to the valence shell of Se, so, $10 e = 5$ electron pairs out of which four are bonded pairs and one is a lone pair. A TBP environment is expected.

2. Predict and draw the structure of I_3^+ using VSEPR model.

Solution



Total number of electron at the central I atom is: $7 + 2 - 1$ (charge) = 8; 2BP + 2LP, should be tetrahedral and angular shape.

3. Which of H_2O and F_2O will have the larger X-O-X bond angle?

Solution:

F is more electronegative than H; therefore the space occupied by the O-H bonding pair in the O valence shell will be greater. Hence, H_2O will have the larger X-O-X bond angle.

4. Explain why the X-P-X bond angles for the series of POX_3 molecules decrease from X = Br (104.1°) to X = Cl (103.3°) to X = F (101.3°)

Solution:


Fluorine is the most electronegative halogen, so it will draw electron density in the P-F bond away from P atom; repulsion of the P-F bonding pairs will be less than the repulsion of P-Cl and P-Br bonding pairs, so the F-P-F bond angle will be the smallest.

5. Show that the following molecules and their corresponding shapes are correct, using VSEPR theory.

BCl_3 ,	trigonal planar
$[IF_5]^{2-}$,	pentagonal planar
$[NH_4]^+$,	tetrahedral
SF_6 ;	octahedral
XeF_4 ;	square planar
AsF_5 ;	trigonal bipyramidal
$Xe(O)F_4$,	square pyramid
IF_7 ,	pentagonal bipyramidal
$[H_3O]^+$	tetrahedral
H_2Se	tetrahedral

6. Work out the geometry of I_3^- ion using VSEPR model.

Solution



This page titled [1.2: VSEPR Theory and its Utility](#) is shared under a [CC BY-NC-SA 4.0](#) license and was authored, remixed, and/or curated by [M. S. Balakrishna & Prasenjit Ghosh](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.