


3.10: Biological Reduction


Addition to a carbonyl by a **semi-anionic** hydride, such as NaBH_4 , results in conversion of the carbonyl compound to an alcohol. The hydride from the BH_4^- anion acts as a nucleophile, adding H^- to the carbonyl carbon. A proton source can then protonate the oxygen of the resulting alkoxide ion, forming an alcohol.



Formally, that process is referred to as a reduction. Reduction generally means a reaction in which electrons are added to a compound; the compound that gains electrons is said to be reduced. Because hydride can be thought of as a proton plus two electrons, we can think of conversion of a ketone or an aldehyde to an alcohol as a two-electron reduction. An aldehyde plus two electrons and two protons becomes an alcohol.

Aldehydes, ketones and alcohols are very common features in biological molecules. Converting between these compounds is a frequent event in many biological pathways. However, semi-anionic compounds like sodium borohydride don't exist in the cell. Instead, a number of biological hydride donors play a similar role.

NADH is a common biological reducing agent. NADH is an acronym for nicotinamide adenine dinucleotide hydride. Instead of an anionic donor that provides a hydride to a carbonyl, NADH is actually a neutral donor. It supplies a hydride to the carbonyl under very specific circumstances. In doing so, it forms a cation, NAD^+ . However, NAD^+ is stabilized by the fact that its nicotinamide ring is aromatic; it was not aromatic in NADH.



This page titled [3.10: Biological Reduction](#) is shared under a [CC BY-NC 3.0](#) license and was authored, remixed, and/or curated by [Chris Schaller](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.