

## 6.13: The Rate of Collisions between Unlike Gas Molecules

We define the collision frequency,  $\tilde{\nu}_{12}$ , as the number of collision per unit time between a single molecule of type 1 and any of the molecules of type 2 present in the same container. We find  $\tilde{\nu}_{12} = N_2 \pi \sigma_{12}^2 \langle v_{12} \rangle$ . If there are  $N_1$  molecules of type 1 present in a unit volume of the gas, the total number of collisions between type 1 molecules and type 2 molecules is  $N_1$  times greater. For clarity, let us refer to the total number of such collisions, per unit volume and per unit time, as the **collision rate**,  $\rho_{12}$ . We have

$$\rho_{12} = N_1 \tilde{\nu}_{12} = N_1 N_2 \pi \sigma_{12}^2 \langle v_{12} \rangle = N_1 N_2 \sigma_{12}^2 \left( \frac{8\pi kT}{\mu} \right)^{1/2}$$

This page titled 6.13: The Rate of Collisions between Unlike Gas Molecules is shared under a [CC BY-SA 4.0](#) license and was authored, remixed, and/or curated by [Paul Ellgen](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

- **4.13: The Rate of Collisions between Unlike Gas Molecules** by [Paul Ellgen](#) is licensed [CC BY-SA 4.0](#). Original source: <https://www.amazon.com/Thermodynamics-Chemical-Equilibrium-Paul-Ellgen/dp/1492114278>.