


21.4: Enolates

For alkylation reactions of enolate anions to be useful, these intermediates must be generated in high concentration in the absence of other strong nucleophiles and bases. The aqueous base conditions used for the aldol condensation are not suitable because the enolate anions of simple carbonyl compounds are formed in very low concentration, and hydroxide or alkoxide bases induce competing S_N2 and $E2$ reactions of alkyl halides. It is necessary, therefore, to achieve complete conversion of aldehyde or ketone reactants to their enolate conjugate bases by treatment with a very strong base ($pK_a > 25$) in a non-hydroxylic solvent before any alkyl halides are added to the reaction system. Some bases that have been used for enolate anion formation are: NaH (sodium hydride, $pK_a > 45$), $NaNH_2$ (sodium amide, $pK_a = 34$), and $LiN[CH(CH_3)_2]_2$ (lithium diisopropylamide, LDA, $pK_a 36$). Ether solvents like tetrahydrofuran (THF) are commonly used for enolate anion formation. With the exception of sodium hydride and sodium amide, most of these bases are soluble in THF. Certain other strong bases, such as alkyl lithium and Grignard reagents, cannot be used to make enolate anions because they rapidly and irreversibly add to carbonyl groups. Nevertheless, these very strong bases are useful in making soluble amide bases. In the preparation of lithium diisopropylamide (LDA), for example, the only other product is the gaseous alkane butane.




LDA



Sodium Amide

Because of its solubility in THF, LDA is a widely used base for enolate anion formation. In this application, one equivalent of diisopropylamine is produced along with the lithium enolate, but this normally does not interfere with the enolate reactions and is easily removed from the products by washing with aqueous acid. Although the reaction of carbonyl compounds with sodium hydride is heterogeneous and slow, sodium enolates are formed with the loss of hydrogen, and no other organic compounds are produced.

The presence of these overlapping p orbitals gives α hydrogens (Hydrogens on carbons adjacent to carbonyls) special properties. In particular, α hydrogens are weakly acidic because the conjugate base, called an enolate, is stabilized through conjugation with the π orbitals of the carbonyl. The effect of the carbonyl is seen when comparing the pK_a for the α hydrogens of aldehydes ($\sim 16-18$), ketones ($\sim 19-21$), and esters ($\sim 23-25$) to the pK_a of an alkane (~ 50).




Of the two resonance structures of the enolate ion the one which places the negative charge on the oxygen is the most stable. This is because the negative charge will be better stabilized by the greater electronegativity of the oxygen.

Acidity of α -Hydrogens in Some Activated Compounds



Compound	RCH_2-NO_2	RCH_2-COR	$RCH_2-C\equiv N$	RCH_2-SO_2R

pK _a	9	20	25	25
-----------------	---	----	----	----

Examples




If the formed enolate is stabilized by more than one carbonyl it is possible to use a weaker base such as sodium ethoxide.




Because of the acidity of α hydrogens, carbonyls undergo keto-enol tautomerism. Tautomers are rapidly interconverted constitutional isomers, usually distinguished by a different bonding location for a labile hydrogen atom and a differently located double bond. The equilibrium between tautomers is not only rapid under normal conditions, but it often strongly favors one of the isomers (acetone, for example, is 99.999% keto tautomer). Even in such one-sided equilibria, evidence for the presence of the minor tautomer comes from the chemical behavior of the compound. Tautomeric equilibria are catalyzed by traces of acids or bases that are generally present in most chemical samples.

General Reaction of Enolates



The Ambident Character of Enolate Anions

Since the negative charge of an enolate anion is delocalized over the α -carbon and the oxygen, as shown earlier, electrophiles may bond to either atom. Reactants having two or more reactive sites are called **ambident**, so this term is properly applied to enolate anions. Modestly electrophilic reactants such as alkyl halides are not sufficiently reactive to combine with neutral enol tautomers, but the increased nucleophilicity of the enolate anion conjugate base permits such reactions to take place. Because alkylations are usually irreversible, their products should reflect the inherent (kinetic) reactivity of the different nucleophilic sites.



If an alkyl halide undergoes an S_N2 reaction at the carbon atom of an enolate anion the product is an alkylated aldehyde or ketone. On the other hand, if the S_N2 reaction occurs at oxygen the product is an ether derivative of the enol tautomer; such compounds are stable in the absence of acid and may be isolated and characterized. These alkylations (shown above) are irreversible under the conditions normally used for S_N2 reactions, so the product composition should provide a measure of the relative rates of substitution at carbon versus oxygen. It has been found that this competition is sensitive to a number of factors, including negative charge density, solvation, cation coordination and product stability.

Contributors

- William Reusch, Professor Emeritus ([Michigan State U.](#)), [Virtual Textbook of Organic Chemistry](#)
- Prof. Steven Farmer ([Sonoma State University](#))

21.4: Enolates is shared under a [CC BY-NC-SA 4.0](#) license and was authored, remixed, and/or curated by LibreTexts.