


## 9.1: Homolytic and Heterolytic Cleavage

For the reactions we learned so far, bond breaking occurs in the way that one part of the bond takes **both** electrons (the electron pair) of the bond away. For example of  $S_N1$  reaction, the leaving group Br leaves with the electron pair to form  $Br^-$  and carbocation intermediate.



example of heterolytic bond cleavage

This process is called **heterolytic bond cleavage**, the  $\sigma$  bond breaks heterolytically. As we have always been doing, an arrow with the **double-barbs** is used to show heterolytic cleavage, that is the transfer of electron pair specifically:



There is another type of bond breaking process, in which each part of the  $\sigma$  bond takes one electron away, as shown below:





### homolytic bond cleavage

This is called **homolytic cleavage**, or **homolysis**. The electron pair separate evenly to each part, and as a result both products contain a single electron. The species that contain one or more single electron is called radical (or free radical). Radicals are produced from homolytic cleavage. The arrow with sing-barb (like the shape of a fishhook) is used to show homolytic cleavage, that is single electron transfer specifically:



Homolysis occurs mainly for non-polar bonds, heat or light (delta is the symbol for heat;  $h\nu$  is used to show light) is needed to provide enough energy for initiating the process. For example:



Radical is another highly reactive reaction intermediate, because of the lack of octet. The substitution reaction we will learn in this chapter involves the radical intermediate.

This page titled [9.1: Homolytic and Heterolytic Cleavage](#) is shared under a [CC BY-NC-SA 4.0](#) license and was authored, remixed, and/or curated by [Xin Liu \(Kwantlen Polytechnic University\)](#).