

10.1: Acids and Bases Definitions

Learning Objectives

- Identify an Arrhenius acid and an Arrhenius base.
- Identify a Brønsted-Lowry acid and a Brønsted-Lowry base.
- Identify conjugate acid-base pairs in an acid-base reaction.


There are three major classifications of substances known as acids or bases. The theory developed by Svante Arrhenius in 1883, the Arrhenius definition, states that an acid produces hydrogen ions, H^+ , in solution and a base produces hydroxide ions, OH^- . Later, two more sophisticated and general theories were proposed. These theories are the Brønsted-Lowry and Lewis definitions of acids and bases. This section will cover the Arrhenius and Brønsted-Lowry theories; [Lewis theory](#) is discussed elsewhere.

The Arrhenius Theory of Acids and Bases

In 1884, the Swedish chemist Svante Arrhenius proposed two specific classifications of compounds, termed acids and bases. When dissolved in an aqueous solution, certain ions were released into the solution. An **Arrhenius acid** is a compound that *increases the concentration* of H^+ ions that are present when added to water. These H^+ ions form the **hydronium ion** (H_3O^+) when they combine with water molecules. This process is represented in a chemical equation by adding H_2O to the reactants side.



In this reaction, hydrochloric acid (HCl) dissociates completely into hydrogen (H^+) and chloride (Cl^-) ions when dissolved in water, thereby releasing H^+ ions into solution. Formation of the hydronium ion equation:



An **Arrhenius base** is a compound that *increases the concentration* of OH^- ions that are present when added to water. The dissociation is represented by the following equation:



In this reaction, sodium hydroxide ($NaOH$) dissociates into sodium (Na^+) and hydroxide (OH^-) ions when dissolved in water, thereby releasing OH^- ions into solution.

Arrhenius acids are substances which produce hydrogen ions in solution and Arrhenius bases are substances which produce hydroxide ions in solution.

Limitations to the Arrhenius Theory


The Arrhenius theory has many more limitations than the other two theories. The theory does not explain the weak base ammonia (NH_3), which in the presence of water, releases hydroxide ions into solution, but does not contain OH^- itself. The Arrhenius definition of acid and base is also limited to aqueous (i.e., water) solutions.

The Brønsted-Lowry Theory of Acids and Bases


In 1923, Danish chemist Johannes Brønsted and English chemist Thomas Lowry independently proposed new definitions for acids and bases, ones that focus on proton transfer. A **Brønsted-Lowry acid** is any species that can donate a proton (H^+) to another molecule. A **Brønsted-Lowry base** is any species that can accept a proton from another molecule. In short, a Brønsted-Lowry acid is a proton donor, while a Brønsted-Lowry base is a proton acceptor.

A Brønsted-Lowry acid is a proton donor, while a Brønsted-Lowry base is a proton acceptor.

Let us use the reaction of ammonia in water to demonstrate the Brønsted-Lowry definitions of an acid and a base. Ammonia and water molecules are reactants, while the ammonium ion and the hydroxide ion are products:




What has happened in this reaction is that the original water molecule has donated a hydrogen ion to the original ammonia molecule, which in turn has accepted the hydrogen ion. We can illustrate this as follows:




Because the water molecule *donates* a hydrogen ion to the ammonia, it is the Brønsted-Lowry acid, while the ammonia molecule—which *accepts* the hydrogen ion—is the Brønsted-Lowry base. Thus, ammonia acts as a base in both the Arrhenius sense and the Brønsted-Lowry sense.

Is an Arrhenius acid like hydrochloric acid still an acid in the Brønsted-Lowry sense? Yes, but it requires us to understand what really happens when HCl is dissolved in water. Recall that the hydrogen *atom* is a single proton surrounded by a single electron. To make the hydrogen ion, we remove the electron, leaving a bare proton. Do we really have bare protons floating around in aqueous solution? No, we do not. What really happens is that the H^+ ion attaches itself to H_2O to make H_3O^+ , which is called the *hydronium ion*. For most purposes, H^+ and H_3O^+ represent the same species, but writing H_3O^+ instead of H^+ shows that we understand that there are no bare protons floating around in solution. Rather, these protons are actually attached to solvent molecules.

With this in mind, how do we define HCl as an acid in the Brønsted-Lowry sense? Consider what happens when HCl is dissolved in H_2O :



We can depict this process using Lewis electron dot diagrams:



Now we see that a hydrogen ion is transferred from the HCl molecule to the H_2O molecule to make chloride ions and hydronium ions. As the hydrogen ion donor, HCl acts as a Brønsted-Lowry acid; the hydrogen ion acceptor, H_2O is a Brønsted-Lowry base. So HCl is an acid not just in the Arrhenius sense, but also in the Brønsted-Lowry sense. Moreover, by the Brønsted-Lowry definitions, H_2O is a base in the formation of aqueous HCl. So the Brønsted-Lowry definitions of an acid and a base classify the dissolving of HCl in water as a reaction between an acid and a base—although the Arrhenius definition would not have labeled H_2O a base in this circumstance.

Note: Acid and Base Definitions


- A Brønsted-Lowry acid is a proton (hydrogen ion) donor.
- A Brønsted-Lowry base is a proton (hydrogen ion) acceptor.
- **All** Arrhenius acids and bases are Brønsted-Lowry acids and bases as well. However, not all Brønsted-Lowry acids and bases are Arrhenius acids and bases.

Example 10.1.1

Aniline ($\text{C}_6\text{H}_5\text{NH}_2$) is slightly soluble in water. It has a nitrogen atom that can accept a hydrogen ion from a water molecule, just like the nitrogen atom in ammonia does. Write the chemical equation for this reaction and identify the Brønsted-Lowry acid and base.

Solution

$\text{C}_6\text{H}_5\text{NH}_2$ and H_2O are the reactants. When $\text{C}_6\text{H}_5\text{NH}_2$ accepts a proton from H_2O , it gains an extra H and a positive charge and leaves an OH^- ion behind. The reaction is as follows:



Because $\text{C}_6\text{H}_5\text{NH}_2$ accepts a proton, it is the Brønsted-Lowry base. The H_2O molecule, because it donates a proton, is the Brønsted-Lowry acid.

? Exercise 10.1.1

Identify the Brønsted-Lowry acid and the Brønsted-Lowry base in this chemical equation.



Answer

Brønsted-Lowry acid: H_2PO_4^- ; Brønsted-Lowry base: H_2O


? Exercise 10.1.2

Which of the following compounds is a Brønsted-Lowry base?


- HCl
- HPO_4^{2-}
- H_3PO_4
- NH_4^+
- CH_3NH_3^+

Answer

A Brønsted-Lowry Base is a proton acceptor, which means it will take in an H^+ . This eliminates HCl , H_3PO_4 , NH_4^+ and CH_3NH_3^+ because they are Brønsted-Lowry acids. They all give away protons. In the case of HPO_4^{2-} , consider the following equation:



Here, it is clear that HPO_4^{2-} is the acid since it donates a proton to water to make H_3O^+ and PO_4^{3-} . Now consider the following equation:




In this case, HPO_4^{2-} is the base since it accepts a proton from water to form H_2PO_4^- and OH^- . Thus, HPO_4^{2-} is an acid and base together, making it amphoteric.

Since HPO_4^{2-} is the only compound from the options that can act as a base, the answer is **(b) HPO_4^{2-}** .

Conjugate Acid-Base Pair

In reality, all acid-base reactions involve the transfer of protons between acids and bases. For example, consider the acid-base reaction that takes place when ammonia is dissolved in water. A water molecule (functioning as an acid) transfers a proton to an ammonia molecule (functioning as a base), yielding the conjugate base of water, OH^- , and the conjugate acid of ammonia, NH_4^+ :



In the reaction of ammonia with water to give ammonium ions and hydroxide ions, ammonia acts as a base by accepting a proton from a water molecule, which in this case means that water is acting as an acid. In the reverse reaction, an ammonium ion acts as an acid by donating a proton to a hydroxide ion, and the hydroxide ion acts as a base. The conjugate acid–base pairs for this reaction are NH_4^+/NH_3 and H_2O/OH^- .





Figure 10.1.1. The pairing of parent acids and bases with conjugate acids and bases.

ACID		BASE	
negligible	OH^-	O^{2-}	strong
	HS^-	S^{2-}	
weak	H_2O	OH^-	weak
	HPO_4^{2-}	PO_4^{3-}	
	HCO_3^-	CO_3^{2-}	
	NH_4^+	NH_3	
	HCN	CN^-	
	$H_2PO_4^-$	HPO_4^{2-}	
	HSO_3^-	SO_3^{2-}	
	H_2S	HS^-	
	H_2CO_3	HCO_3^-	
	$C_5H_5NH^+$	C_5H_5N	
	CH_3CO_2H	$CH_3CO_2^-$	
	HF	F^-	
	H_3PO_4	$H_2PO_4^-$	
	H_2SO_3	HSO_3^-	
	HSO_4^-	SO_4^{2-}	
strong	H_3O^+	H_2O	negligible
	HNO_3	NO_3^-	
	H_2SO_4	HSO_4^-	
	HCl	Cl^-	
	HBr	Br^-	

Figure 10.1.2: The Relative Strengths of Some Common Conjugate Acid-Base Pairs. The strongest acids are at the bottom left, and the strongest bases are at the top right. The conjugate base of a strong acid is a very weak base, and, conversely, the conjugate acid of a strong base is a very weak acid.

✓ Example 10.1.2

Identify the conjugate acid-base pairs in this equilibrium.




Solution

Similarly, in the reaction of acetic acid with water, acetic acid **donates** a proton to water, which acts as the base. In the reverse reaction, H_3O^+ is the acid that donates a proton to the acetate ion, which acts as the base.


Once again, we have two conjugate acid-base pairs:

- the parent acid and its conjugate base ($\text{CH}_3\text{CO}_2\text{H}/\text{CH}_3\text{CO}_2^-$) and
- the parent base and its conjugate acid ($\text{H}_3\text{O}^+/\text{H}_2\text{O}$).



✓ Example 10.1.3

Identify the conjugate acid-base pairs in this equilibrium.




Solution

One pair is H_2O and OH^- , where H_2O has one more H^+ and is the conjugate acid, while OH^- has one less H^+ and is the conjugate base.

The other pair consists of $(\text{CH}_3)_3\text{N}$ and $(\text{CH}_3)_3\text{NH}^+$, where $(\text{CH}_3)_3\text{NH}^+$ is the conjugate acid (it has an additional proton) and $(\text{CH}_3)_3\text{N}$ is the conjugate base.

? Exercise 10.1.3

Identify the conjugate acid-base pairs in this equilibrium.



Answer

H_2O (acid) and OH^- (base); NH_2^- (base) and NH_3 (acid)

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

10.1: Acids and Bases Definitions is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by LibreTexts.