

## 8.S: Solids, Liquids, and Gases (Summary)

*To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms in the following summary and ask yourself how they relate to the topics in the chapter.*

A **phase** is a certain form of matter that has the same physical properties throughout. Three phases are common: the solid, the liquid, and the gas phase. What determines the phase of a substance? Generally, the strength of the **intermolecular interactions** determines whether a substance is a solid, liquid, or gas under any particular conditions. **Covalent network bonding** is a very strong form of intermolecular interaction. Diamond is one example of a substance that has this intermolecular interaction. **Ionic interactions**, the forces of attraction due to oppositely charged ions, are also relatively strong. Covalent bonds are another type of interaction within molecules, but if the bonds are **polar covalent bonds**, then the unequal sharing of electrons can cause charge imbalances within molecules that cause interactions between molecules. These molecules are described as **polar**, and these interactions are called **dipole-dipole interactions**. A certain rather strong type of dipole-dipole interaction, involving a hydrogen atom, is called **hydrogen bonding**. On the other hand, equal sharing of electrons forms **nonpolar covalent bonds**, and the interactions between different molecules is less because the molecules are nonpolar. All substances have very weak **dispersion forces** (also called **London forces**) caused by the movement of electrons within the bonds themselves.

In the solid phase, intermolecular interactions are so strong that they hold the individual atoms or molecules in place. In many solids, the regular three-dimensional arrangement of particles makes a **crystal**. In other solids, the irregular arrangement of particles makes an **amorphous** solid. In liquids, the intermolecular interactions are strong enough to keep the particles of substance together but not in place. Thus, the particles are free to move over each other but still remain in contact.

In gases, the intermolecular interactions are weak enough that the individual particles are separated from each other in space. The **kinetic theory of gases** is a collection of statements that describe the fundamental behavior of all gases. Among other properties, gases exert a **pressure** on their container. Pressure is measured using units like **pascal**, **bar**, **atmosphere**, or **mmHg** (also called a **torr**).

There are several simple relationships between the variables used to describe a quantity of gas. These relationships are called **gas laws**. **Boyle's law** relates the pressure and volume of a gas, while **Charles's law** relates the volume and absolute temperature of a gas. The **combined gas law** relates the volume, pressure, and absolute temperature of a gas sample. All of these gas laws allow us to understand the changing conditions of a gas. The **ideal gas law** relates the pressure, volume, amount, and absolute temperature of a gas under any conditions. These four variables are related to the **ideal gas law constant**, which is the proportionality constant used to calculate the conditions of a gas. Because the conditions of a gas can change, a set of benchmark conditions called **standard temperature and pressure (STP)** is defined. Standard temperature is 0°C, and standard pressure is 1.00 atm.

This page titled [8.S: Solids, Liquids, and Gases \(Summary\)](#) is shared under a [CC BY-NC-SA 4.0](#) license and was authored, remixed, and/or curated by [Anonymous](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

- [8.S: Solids, Liquids, and Gases \(Summary\)](#) by Anonymous is licensed [CC BY-NC-SA 3.0](#). Original source: <https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological>.