

16.2: A Qualitative Description of Acid-Base Equilibria


Learning Objectives

- To understand the concept of conjugate acid–base pairs.
- To know the relationship between acid or base strength and the magnitude of K_a , K_b , pK_a , and pK_b .
- To understand the leveling effect.

We now turn our attention to acid–base reactions to see how the concepts of chemical equilibrium and equilibrium constants can deepen our understanding of this kind of chemical behavior. We begin with a qualitative description of acid–base equilibria in terms of the Brønsted–Lowry model and then proceed to a quantitative description in [Section 16.4](#).

Conjugate Acid–Base Pairs


In aqueous solutions, acids and bases can be defined in terms of the transfer of a proton from an acid to a base. Thus for every acidic species in an aqueous solution, there exists a species derived from the acid by the loss of a proton. These two species that differ by only a proton constitute a conjugate acid–base pair. For example, in the reaction of HCl with water ([Equation 16.1.1](#)), HCl, the *parent acid*, donates a proton to a water molecule, the *parent base*, thereby forming Cl^- . Thus HCl and Cl^- constitute a conjugate acid–base pair. By convention, we always write a conjugate acid–base pair as the acid followed by its conjugate base. In the reverse reaction, the Cl^- ion in solution acts as a base to accept a proton from H_3O^+ , forming H_2O and HCl. Thus H_3O^+ and H_2O constitute a second conjugate acid–base pair. In general, any acid–base reaction must contain *two* conjugate acid–base pairs, which in this case are HCl/Cl^- and $\text{H}_3\text{O}^+/\text{H}_2\text{O}$.



Note the Pattern

All acid–base reactions contain *two* conjugate acid–base pairs.

Similarly, in the reaction of acetic acid with water, acetic acid donates a proton to water, which acts as the base. In the reverse reaction, H_3O^+ is the acid that donates a proton to the acetate ion, which acts as the base. Once again, we have two conjugate acid–base pairs: the parent acid and its conjugate base ($\text{CH}_3\text{CO}_2\text{H}/\text{CH}_3\text{CO}_2^-$) and the parent base and its conjugate acid ($\text{H}_3\text{O}^+/\text{H}_2\text{O}$).



In the reaction of ammonia with water to give ammonium ions and hydroxide ions ([Equation 16.1.3](#)), ammonia acts as a base by accepting a proton from a water molecule, which in this case means that water is acting as an acid. In the reverse reaction, an ammonium ion acts as an acid by donating a proton to a hydroxide ion, and the hydroxide ion acts as a base. The conjugate acid–base pairs for this reaction are $\text{NH}_4^+/\text{NH}_3$ and $\text{H}_2\text{O}/\text{OH}^-$. Some common conjugate acid–base pairs are shown in [Figure 16.2.1](#).




Figure 16.2.1 The Relative Strengths of Some Common Conjugate Acid–Base Pairs The strongest acids are at the bottom left, and the strongest bases are at the top right. The conjugate base of a strong acid is a very weak base, and, conversely, the conjugate acid of a strong base is a very weak acid.

Acid–Base Equilibrium Constants: K_a , K_b , pK_a , and pK_b

The magnitude of the equilibrium constant for an ionization reaction can be used to determine the relative strengths of acids and bases. For example, the general equation for the ionization of a weak acid in water, where HA is the parent acid and A⁻ is its conjugate base, is as follows:



The equilibrium constant for this dissociation is as follows:

$$K = \frac{[H_3O^+][A^-]}{[H_2O][HA]} \quad (16.2.2)$$

As we noted earlier, the concentration of water is essentially constant for all reactions in aqueous solution, so [H₂O] in Equation 16.2.2 can be incorporated into a new quantity, the acid ionization constant (K_a). An equilibrium constant for the ionization (dissociation) of a weak acid (HA) with water, HA(aq) + H₂O(l) ⇌ H₃O(aq)⁺ + A⁻(aq), in which the concentration of water is treated as a constant: $K_a = [H_3O(aq)^+][A^-(aq)]/[HA]$, also called the *acid dissociation constant*:

$$K_a = K[H_2O] = \frac{[H_3O^+][A^-]}{[HA]} \quad (16.2.3)$$

Thus the numerical values of K and K_a differ by the concentration of water (55.3 M). Again, for simplicity, H₃O⁺ can be written as H⁺ in Equation 16.2.3. Keep in mind, though, that free H⁺ does not exist in aqueous solutions and that a proton is transferred to H₂O in all acid ionization reactions to form H₃O⁺. The larger the K_a , the stronger the acid and the higher the H⁺ concentration at equilibrium. Like all equilibrium constants, acid–base ionization constants are actually measured in terms of the *activities* of H⁺ or OH⁻, thus making them unitless. The values of K_a for a number of common acids are given in Table 16.2.1.

Table 16.2.1 Values of K_a , pK_a , K_b , and pK_b for Selected Acids (HA) and Their Conjugate Bases (A⁻)

Acid	HA	K_a	pK_a	A^-	K_b	pK_b
hydroiodic acid	HI	2×10^9	-9.3	I^-	5.5×10^{-24}	23.26
sulfuric acid (1)*	H_2SO_4	1×10^2	-2.0	HSO_4^-	1×10^{-16}	16.0
nitric acid	HNO_3	2.3×10^1	-1.37	NO_3^-	4.3×10^{-16}	15.37
hydronium ion	H_3O^+	1.0	0.00	H_2O	1.0×10^{-14}	14.00
sulfuric acid (2)*	HSO_4^-	1.0×10^{-2}	1.99	SO_4^{2-}	9.8×10^{-13}	12.01
hydrofluoric acid	HF	6.3×10^{-4}	3.20	F^-	1.6×10^{-11}	10.80
nitrous acid	HNO_2	5.6×10^{-4}	3.25	NO_2^-	1.8×10^{-11}	10.75
formic acid	HCO_2H	1.78×10^{-4}	3.750	HCO_2^-	5.6×10^{-11}	10.25
benzoic acid	$C_6H_5CO_2H$	6.3×10^{-5}	4.20	$C_6H_5CO_2^-$	1.6×10^{-10}	9.80
acetic acid	CH_3CO_2H	1.7×10^{-5}	4.76	$CH_3CO_2^-$	5.8×10^{-10}	9.24
pyridinium ion	$C_5H_5NH^+$	5.9×10^{-6}	5.23	C_5H_5N	1.7×10^{-9}	8.77
hypochlorous acid	HOCl	4.0×10^{-8}	7.40	OCl^-	2.5×10^{-7}	6.60
hydrocyanic acid	HCN	6.2×10^{-10}	9.21	CN^-	1.6×10^{-5}	4.79
ammonium ion	NH_4^+	5.6×10^{-10}	9.25	NH_3	1.8×10^{-5}	4.75
water	H_2O	1.0×10^{-14}	14.00	OH^-	1.00	0.00
acetylene	C_2H_2	1×10^{-26}	26.0	HC_2^-	1×10^{12}	-12.0
ammonia	NH_3	1×10^{-35}	35.0	NH_2^-	1×10^{21}	-21.0

*The number in parentheses indicates the ionization step referred to for a polyprotic acid.

Weak bases react with water to produce the hydroxide ion, as shown in the following general equation, where B is the parent base and BH^+ is its conjugate acid:



The equilibrium constant for this reaction is the base ionization constant (K_b). An equilibrium constant for the reaction of a weak base (B) with water, $B(aq) + H_2O(l) \rightleftharpoons BH^+(aq) + OH^-(aq)$, in which the concentration of water is treated as a constant: $K_b = [BH^+][OH^-]/[B]$, also called the *base dissociation constant*:

$$K_b = K[H_2O] = [BH^+][OH^-]/[B] \quad (16.2.5)$$

Once again, the concentration of water is constant, so it does not appear in the equilibrium constant expression; instead, it is included in the K_b . The larger the K_b , the stronger the base and the higher the OH^- concentration at equilibrium. The values of K_b for a number of common weak bases are given in [Table 16.2.2](#)

Table 16.2.2 Values of K_b , pK_b , K_a , and pK_a for Selected Weak Bases (B) and Their Conjugate Acids (BH^+)


Base	B	K_b	pK_b	BH^+	K_a	pK_a
hydroxide ion	OH^-	1.0	0.00*	H_2O	1.0×10^{-14}	14.00
phosphate ion	PO_4^{3-}	2.1×10^{-2}	1.68	HPO_4^{2-}	4.8×10^{-13}	12.32
dimethylamine	$(CH_3)_2NH$	5.4×10^{-4}	3.27	$(CH_3)_2NH_2^+$	1.9×10^{-11}	10.73

*As in [Table 16.2](#).

Base	B	K_b	pK_b	BH^+	K_a	pK_a
methylamine	CH_3NH_2	4.6×10^{-4}	3.34	$CH_3NH_3^+$	2.2×10^{-11}	10.66
trimethylamine	$(CH_3)_3N$	6.3×10^{-5}	4.20	$(CH_3)_3NH^+$	1.6×10^{-10}	9.80
ammonia	NH_3	1.8×10^{-5}	4.75	NH_4^+	5.6×10^{-10}	9.25
pyridine	C_5H_5N	1.7×10^{-9}	8.77	$C_5H_5NH^+$	5.9×10^{-6}	5.23
aniline	$C_6H_5NH_2$	7.4×10^{-10}	9.13	$C_6H_5NH_3^+$	1.3×10^{-5}	4.87
water	H_2O	1.0×10^{-14}	14.00	H_3O^+	1.0*	0.00

*As in Table 16.2.

There is a simple relationship between the magnitude of K_a for an acid and K_b for its conjugate base. Consider, for example, the ionization of hydrocyanic acid (HCN) in water to produce an acidic solution, and the reaction of CN^- with water to produce a basic solution:




The equilibrium constant expression for the ionization of HCN is as follows:

$$K_a = \frac{[H^+][CN^-]}{[HCN]} \quad (16.2.8)$$

The corresponding expression for the reaction of cyanide with water is as follows:

$$K_b = \frac{[OH^-][HCN]}{[CN^-]} \quad (16.2.9)$$

If we add Equation 16.2.8 and Equation 16.2.9, we obtain the following (recall from Chapter 15 that the equilibrium constant for the sum of two reactions is the product of the equilibrium constants for the individual reactions):



In this case, the sum of the reactions described by K_a and K_b is the equation for the autoionization of water, and the product of the two equilibrium constants is K_w :

$$K_a K_b = K_w \quad (16.2.10)$$

Thus if we know either K_a for an acid or K_b for its conjugate base, we can calculate the other equilibrium constant for any conjugate acid–base pair.

Just as with pH, pOH, and pK_w , we can use negative logarithms to avoid exponential notation in writing acid and base ionization constants, by defining pK_a as follows:

$$pK_a = -\log_{10} K_a \quad (16.2.11)$$

$$K_a = 10^{-pK_a} \quad (16.2.12)$$

and pK_b as

$$pK_b = -\log_{10} K_b \quad (16.2.13)$$

$$K_b = 10^{-pK_b} \quad (16.2.14)$$

Similarly, Equation 16.2.10, which expresses the relationship between K_a and K_b , can be written in logarithmic form as follows:

$$pK_a + pK_b = pK_w \quad (16.2.15)$$

At 25°C, this becomes

$$pK_a + pK_b = 14.00 \quad (16.2.16)$$


The values of pK_a and pK_b are given for several common acids and bases in Table 16.2.1 and Table 16.2.2, respectively, and a more extensive set of data is provided in Tables E1 and E2. Because of the use of negative logarithms, smaller values of pK_a correspond to larger acid ionization constants and hence stronger acids. For example, nitrous acid (HNO_2), with a pK_a of 3.25, is about a 1000 times stronger acid than hydrocyanic acid (HCN), with a pK_a of 9.21. Conversely, smaller values of pK_b correspond to larger base ionization constants and hence stronger bases.

The relative strengths of some common acids and their conjugate bases are shown graphically in Figure 16.2.1. The conjugate acid–base pairs are listed in order (from top to bottom) of increasing acid strength, which corresponds to decreasing values of pK_a . This order corresponds to decreasing strength of the conjugate base or increasing values of pK_b . At the bottom left of Figure 16.2 are the common strong acids; at the top right are the most common strong bases. Notice the inverse relationship between the strength of the parent acid and the strength of the conjugate base. Thus the conjugate base of a strong acid is a very weak base, and the conjugate base of a very weak acid is a strong base.

Note the Pattern


The conjugate base of a strong acid is a weak base and vice versa.

We can use the relative strengths of acids and bases to predict the direction of an acid–base reaction by following a single rule: *an acid–base equilibrium always favors the side with the weaker acid and base, as indicated by these arrows:*




In an acid–base reaction, the proton always reacts with the stronger base.


For example, hydrochloric acid is a strong acid that ionizes essentially completely in dilute aqueous solution to produce H_3O^+ and Cl^- ; only negligible amounts of HCl molecules remain undissociated. Hence the ionization equilibrium lies virtually all the way to the right, as represented by a single arrow:



In contrast, acetic acid is a weak acid, and water is a weak base. Consequently, aqueous solutions of acetic acid contain mostly acetic acid molecules in equilibrium with a small concentration of H_3O^+ and acetate ions, and the ionization equilibrium lies far to the left, as represented by these arrows:



Similarly, in the reaction of ammonia with water, the hydroxide ion is a strong base, and ammonia is a weak base, whereas the ammonium ion is a stronger acid than water. Hence this equilibrium also lies to the left:



Note the Pattern

All acid–base equilibria favor the side with the weaker acid and base. Thus the proton is bound to the stronger base.

Example 16.2.1

1. Calculate K_b and pK_b of the butyrate ion ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CO}_2^-$). The pK_a of butyric acid at 25°C is 4.83. Butyric acid is responsible for the foul smell of rancid butter.
2. Calculate K_a and pK_a of the dimethylammonium ion $[(\text{CH}_3)_2\text{NH}_2^+]$. The base ionization constant K_b of dimethylamine $[(\text{CH}_3)_2\text{NH}]$ is 5.4×10^{-4} at 25°C.

Given: pK_a and K_b

Asked for: corresponding K_b and pK_b , K_a and pK_a

Strategy:

The constants K_a and K_b are related as shown in [Equation 16.2.10](#). The pK_a and pK_b for an acid and its conjugate base are related as shown in [Equation 16.2.15](#) and [Equation 16.2.16](#). Use the relationships $pK = -\log K$ and $K = 10^{-pK}$ ([Equation 16.24](#) and [Equation 16.26](#)) to convert between K_a and pK_a or K_b and pK_b .

Solution:

1. We are given the pK_a for butyric acid and asked to calculate the K_b and the pK_b for its conjugate base, the butyrate ion. Because the pK_a value cited is for a temperature of 25°C, we can use [Equation 16.2.16](#): $pK_a + pK_b = pK_w = 14.00$. Substituting the pK_a and solving for the pK_b ,

$$4.83 + pK_b = 14.00$$

$$pK_b = 14.00 - 4.83 = 9.17$$

Because $pK_b = -\log K_b$, K_b is $10^{-9.17} = 6.8 \times 10^{-10}$.

2. In this case, we are given K_b for a base (dimethylamine) and asked to calculate K_a and pK_a for its conjugate acid, the dimethylammonium ion. Because the initial quantity given is K_b rather than pK_b , we can use [Equation 16.23](#): $K_a K_b = K_w$. Substituting the values of K_b and K_w at 25°C and solving for K_a ,

$$K_a(5.4 \times 10^{-4}) = 1.01 \times 10^{-14}$$

$$K_a = 1.9 \times 10^{-11}$$

Because $pK_a = -\log K_a$, we have $pK_a = -\log(1.9 \times 10^{-11}) = 10.72$. We could also have converted K_b to pK_b to obtain the same answer:

$$pK_b = -\log(5.4 \times 10^{-4}) = 3.27$$

$$pK_a + pK_b = 14.00$$

$$pK_a = 10.73$$

$$K_a = 10^{-pK_a} = 10^{-10.73} = 1.9 \times 10^{-11}$$

If we are given any one of these four quantities for an acid or a base (K_a , pK_a , K_b , or pK_b), we can calculate the other three.

Exercise

Lactic acid [$\text{CH}_3\text{CH}(\text{OH})\text{CO}_2\text{H}$] is responsible for the pungent taste and smell of sour milk; it is also thought to produce soreness in fatigued muscles. Its pK_a is 3.86 at 25°C. Calculate K_a for lactic acid and pK_b and K_b for the lactate ion.

Answer: $K_a = 1.4 \times 10^{-4}$ for lactic acid; $pK_b = 10.14$ and $K_b = 7.2 \times 10^{-11}$ for the lactate ion

Solutions of Strong Acids and Bases: The Leveling Effect

You will notice in [Table 16.2.1](#) that acids like H_2SO_4 and HNO_3 lie *above* the hydronium ion, meaning that they have pK_a values less than zero and are stronger acids than the H_3O^+ ion. Recall from [Chapter 8](#) that the acidic proton in virtually all oxoacids is bonded to one of the oxygen atoms of the oxoanion. Thus nitric acid should properly be written as HONO_2 . Unfortunately, however, the formulas of oxoacids are almost always written with hydrogen on the left and oxygen on the right, giving HNO_3 instead. In fact, all six of the common strong acids that we first encountered in [Chapter 8](#) have pK_a values less than zero, which means that they have a greater tendency to lose a proton than does the H_3O^+ ion. Conversely, the conjugate bases of these strong acids are weaker bases than water. Consequently, the proton-transfer equilibria for these strong acids lie far to the right, and adding any of the common strong acids to water results in an essentially stoichiometric reaction of the acid with water to form a solution of the H_3O^+ ion and the conjugate base of the acid.

Although K_a for HI is about 10^8 greater than K_a for HNO_3 , the reaction of either HI or HNO_3 with water gives an essentially stoichiometric solution of H_3O^+ and I^- or NO_3^- . In fact, a 0.1 M aqueous solution of any strong acid actually contains 0.1 M H_3O^+ , *regardless of the identity of the strong acid*. This phenomenon is called the leveling effect. The phenomenon that makes H_3O^+ the strongest acid that can exist in water. Any species that is a stronger acid than H_3O^+ is leveled to the strength of H_3O^+ in aqueous solution.: any species that is a stronger acid than the conjugate acid of water (H_3O^+) is leveled to the strength of H_3O^+ in aqueous



solution because H_3O^+ is the strongest acid that can exist in equilibrium with water. Consequently, it is impossible to distinguish between the strengths of acids such as HI and HNO_3 in aqueous solution, and an alternative approach must be used to determine their relative acid strengths.

One method is to use a solvent such as anhydrous acetic acid. Because acetic acid is a stronger acid than water, it must also be a weaker base, with a lesser tendency to accept a proton than H_2O . Measurements of the conductivity of 0.1 M solutions of both HI and HNO_3 in acetic acid show that HI is completely dissociated, but HNO_3 is only partially dissociated and behaves like a weak acid *in this solvent*. This result clearly tells us that HI is a stronger acid than HNO_3 . The relative order of acid strengths and approximate K_a and $\text{p}K_a$ values for the strong acids at the top of [Table 16.2.1](#) were determined using measurements like this and different nonaqueous solvents.

Note the Pattern

In aqueous solutions, $[\text{H}_3\text{O}^+]$ is the strongest acid and OH^- is the strongest base that can exist in equilibrium with H_2O .


The leveling effect applies to solutions of strong bases as well: In aqueous solution, any base stronger than OH^- is leveled to the strength of OH^- because OH^- is the strongest base that can exist in equilibrium with water. Salts such as K_2O , NaOCH_3 (sodium methoxide), and NaNH_2 (sodamide, or sodium amide), whose anions are the conjugate bases of species that would lie below water in [Table 16.3](#), are all strong bases that react essentially completely (and often violently) with water, accepting a proton to give a solution of OH^- and the corresponding cation:



Other examples that you may encounter are potassium hydride (KH) and organometallic compounds such as methyl lithium (CH_3Li).

Polyprotic Acids and Bases

As you learned in [Chapter 8](#), *polyprotic acids* such as H_2SO_4 , H_3PO_4 , and H_2CO_3 contain more than one ionizable proton, and the protons are lost in a stepwise manner. The fully protonated species is always the strongest acid because it is easier to remove a proton from a neutral molecule than from a negatively charged ion. Thus acid strength decreases with the loss of subsequent protons, and, correspondingly, the $\text{p}K_a$ increases. Consider H_2SO_4 , for example:





The equilibrium in the first reaction lies far to the right, consistent with H_2SO_4 being a strong acid. In contrast, in the second reaction, appreciable quantities of both HSO_4^- and SO_4^{2-} are present at equilibrium.

Note the Pattern

For a polyprotic acid, acid strength decreases and the $\text{p}K_a$ increases with the sequential loss of each proton.

The hydrogen sulfate ion (HSO_4^-) is both the conjugate base of H_2SO_4 and the conjugate acid of SO_4^{2-} . Just like water, HSO_4^- can therefore act as *either* an acid or a base, depending on whether the other reactant is a stronger acid or a stronger base. Conversely, the sulfate ion (SO_4^{2-}) is a *polyprotic base* that is capable of accepting two protons in a stepwise manner:



Like any other conjugate acid–base pair, the strengths of the conjugate acids and bases are related by $\text{p}K_a + \text{p}K_b = \text{p}K_w$. Consider, for example, the $\text{HSO}_4^-/\text{SO}_4^{2-}$ conjugate acid–base pair. From [Table 16.2.1](#), we see that the $\text{p}K_a$ of HSO_4^- is 1.99. Hence the $\text{p}K_b$ of SO_4^{2-} is $14.00 - 1.99 = 12.01$. Thus sulfate is a rather weak base, whereas OH^- is a strong base, so the equilibrium shown above lies to the left. The HSO_4^- ion is also a very weak base [$\text{p}K_a$ of $\text{H}_2\text{SO}_4 = 2.0$, $\text{p}K_b$ of $\text{HSO}_4^- = 14 - (-2.0) = 16$], which is consistent with what we expect for the conjugate base of a strong acid. Thus the equilibrium also lies almost completely to the left. Once again, equilibrium favors the formation of the weaker acid–base pair.

Example 16.2.2

Predict whether the equilibrium for each reaction lies to the left or the right as written.

- $NH_4^+_{(aq)} + PO_4^{3-}_{(aq)} \rightleftharpoons NH_{3(aq)} + HPO_4^{2-}_{(aq)}$
- $CH_3CH_2CO_2H_{(aq)} + CN^-_{(aq)} \rightleftharpoons CH_3CH_2CO_2^-_{(aq)} + HCN_{(aq)}$

Given: balanced chemical equation


Asked for: equilibrium position

Strategy:


Identify the conjugate acid–base pairs in each reaction. Then refer to [Table 16.2.1](#) [Table 16.2.2](#), and [Figure 16.2.1](#) to determine which is the stronger acid and base. Equilibrium always favors the formation of the weaker acid–base pair.

Solution:

1. The conjugate acid–base pairs are NH_4^+/NH_3 and HPO_4^{2-}/PO_4^{3-} . According to [Table 16.2](#) and [Table 16.3](#), NH_4^+ is a stronger acid ($pK_a = 9.25$) than HPO_4^{2-} ($pK_a = 12.32$), and PO_4^{3-} is a stronger base ($pK_b = 1.68$) than NH_3 ($pK_b = 4.75$). The equilibrium will therefore lie to the right, favoring the formation of the weaker acid–base pair:



2. The conjugate acid–base pairs are $CH_3CH_2CO_2H/CH_3CH_2CO_2^-$ and HCN/CN^- . According to [Table 16.2.1](#), HCN is a weak acid ($pK_a = 9.21$) and CN^- is a moderately weak base ($pK_b = 4.79$). Propionic acid ($CH_3CH_2CO_2H$) is not listed in [Table 16.2.1](#), however. In a situation like this, the best approach is to look for a similar compound whose acid–base properties *are* listed. For example, propionic acid and acetic acid are identical except for the groups attached to the carbon atom of the carboxylic acid ($-CH_2CH_3$ versus $-CH_3$), so we might expect the two compounds to have similar acid–base properties. In particular, we would expect the pK_a of propionic acid to be similar in magnitude to the pK_a of acetic acid. (In fact, the pK_a of propionic acid is 4.87, compared to 4.76 for acetic acid, which makes propionic acid a slightly weaker acid than acetic acid.) Thus propionic acid should be a significantly stronger acid than HCN . Because the stronger acid forms the weaker conjugate base, we predict that cyanide will be a stronger base than propionate. The equilibrium will therefore lie to the right, favoring the formation of the weaker acid–base pair:



Exercise

Predict whether the equilibrium for each reaction lies to the left or the right as written.


1. $H_2O_{(l)} + HS^-_{(aq)} \rightleftharpoons OH^-_{(aq)} + H_2S_{(aq)}$
2. $HCO_2^-_{(aq)} + HSO_4^-_{(aq)} \rightleftharpoons HCO_2H_{(aq)} + SO_4^{2-}_{(aq)}$

Answer

1. left
2. left

Acid–Base Properties of Solutions of Salts



We can also use the relative strengths of conjugate acid–base pairs to understand the acid–base properties of solutions of salts. In [Chapter 8](#), you learned that a *neutralization reaction* can be defined as the reaction of an acid and a base to produce a salt and water. That is, another cation, such as Na^+ , replaces the proton on the acid. An example is the reaction of CH_3CO_2H , a weak acid, with $NaOH$, a strong base:



Depending on the acid–base properties of its component ions, however, a salt can dissolve in water to produce a neutral solution, a basic solution, or an acidic solution.



When a salt such as NaCl dissolves in water, it produces $\text{Na}^+(\text{aq})$ and $\text{Cl}^-(\text{aq})$ ions. Using a Lewis approach, the Na^+ ion can be viewed as an acid because it is an electron pair acceptor, although its low charge and relatively large radius make it a very weak acid. The Cl^- ion is the conjugate base of the strong acid HCl, so it has essentially no basic character. Consequently, dissolving NaCl in water has no effect on the pH of a solution, and the solution remains neutral.

Now let's compare this behavior to the behavior of aqueous solutions of potassium cyanide and sodium acetate. Again, the cations (K^+ and Na^+) have essentially no acidic character, but the anions (CN^- and CH_3CO_2^-) are weak bases that can react with water because they are the conjugate bases of the weak acids HCN and acetic acid, respectively.



Neither reaction proceeds very far to the right as written because the formation of the weaker acid–base pair is favored. Both HCN and acetic acid are stronger acids than water, and hydroxide is a stronger base than either acetate or cyanide, so in both cases, the equilibrium lies to the left. Nonetheless, each of these reactions generates enough hydroxide ions to produce a basic solution. For example, the pH of a 0.1 M solution of sodium acetate or potassium cyanide at 25°C is 8.8 or 11.1, respectively. From [Table 16.2.2](#) and [Figure 16.2.1](#), we can see that CN^- is a stronger base ($\text{pK}_\text{b} = 4.79$) than acetate ($\text{pK}_\text{b} = 9.24$), which is consistent with KCN producing a more basic solution than sodium acetate at the same concentration.


In contrast, the conjugate acid of a weak base should be a weak acid ([Figure 16.2.1](#)). For example, ammonium chloride and pyridinium chloride are salts produced by reacting ammonia and pyridine, respectively, with HCl. As you already know, the chloride ion is such a weak base that it does not react with water. In contrast, the cations of the two salts are weak acids that react with water as follows:



[Figure 16.2.1](#) shows that H_3O^+ is a stronger acid than either NH_4^+ or $\text{C}_5\text{H}_5\text{NH}^+$, and conversely, ammonia and pyridine are both stronger bases than water. The equilibrium will therefore lie far to the left in both cases, favoring the weaker acid–base pair. The H_3O^+ concentration produced by the reactions is great enough, however, to decrease the pH of the solution significantly: the pH of a 0.10 M solution of ammonium chloride or pyridinium chloride at 25°C is 5.13 or 3.12, respectively. This is consistent with the information shown in [Figure 16.2.1](#), indicating that the pyridinium ion is more acidic than the ammonium ion.

What happens with aqueous solutions of a salt such as ammonium acetate, where both the cation *and* the anion can react separately with water to produce an acid and a base, respectively? According to Eq 16.2.25, the ammonium ion will lower the pH, while according to Eq 16.2.24, the acetate ion will raise the pH. This particular case is unusual, in that the cation is as strong an acid as the anion is a base ($\text{pK}_\text{a} \approx \text{pK}_\text{b}$). Consequently, the two effects cancel, and the solution remains neutral. With salts in which the cation is a stronger acid than the anion is a base, the final solution has a $\text{pH} < 7.00$. Conversely, if the cation is a weaker acid than the anion is a base, the final solution has a $\text{pH} > 7.00$.

Solutions of simple salts of metal ions can also be acidic, even though a metal ion cannot donate a proton directly to water to produce H_3O^+ . Instead, a metal ion can act as a Lewis acid and interact with water, a Lewis base, by coordinating to a lone pair of electrons on the oxygen atom to form a hydrated metal ion (part (a) in [Figure 16.2.2](#)), as discussed in [Chapter 8](#). A water molecule coordinated to a metal ion is more acidic than a free water molecule for two reasons. First, repulsive electrostatic interactions between the positively charged metal ion and the partially positively charged hydrogen atoms of the coordinated water molecule make it easier for the coordinated water to lose a proton.



Second, the positive charge on the Al^{3+} ion attracts electron density from the oxygen atoms of the water molecules, which decreases the electron density in the O–H bonds, as shown in part (b) in Figure 16.2.2. With less electron density between the O atoms and the H atoms, the O–H bonds are weaker than in a free H_2O molecule, making it easier to lose a H^+ ion.




Figure 16.2.2 Effect of a Metal Ion on the Acidity of Water (a) Reaction of the metal ion Al^{3+} with water to form the hydrated metal ion is an example of a Lewis acid–base reaction. (b) The positive charge on the aluminum ion attracts electron density from the oxygen atoms, which shifts electron density away from the O–H bonds. The decrease in electron density weakens the O–H bonds in the water molecules and makes it easier for them to lose a proton.

The magnitude of this effect depends on the following two factors (Figure 16.2.3):

1. **The charge on the metal ion.** A divalent ion (M^{2+}) has approximately twice as strong an effect on the electron density in a coordinated water molecule as a monovalent ion (M^+) of the same radius.
2. **The radius of the metal ion.** For metal ions with the same charge, the smaller the ion, the shorter the internuclear distance to the oxygen atom of the water molecule and the greater the effect of the metal on the electron density distribution in the water molecule.





Figure 16.2.3 The Effect of the Charge and Radius of a Metal Ion on the Acidity of a Coordinated Water Molecule The contours show the electron density on the O atoms and the H atoms in both a free water molecule (left) and water molecules coordinated to Na^+ , Mg^{2+} , and Al^{3+} ions. These contour maps demonstrate that the smallest, most highly charged metal ion (Al^{3+}) causes the greatest decrease in electron density of the O–H bonds of the water molecule. Due to this effect, the acidity of hydrated metal ions increases as the charge on the metal ion increases and its radius decreases.

Thus aqueous solutions of small, highly charged metal ions, such as Al^{3+} and Fe^{3+} , are acidic:



The $[\text{Al}(\text{H}_2\text{O})_6]^{3+}$ ion has a pK_a of 5.0, making it almost as strong an acid as acetic acid. Because of the two factors described previously, the most important parameter for predicting the effect of a metal ion on the acidity of coordinated water molecules is the *charge-to-radius ratio* of the metal ion. A number of pairs of metal ions that lie on a diagonal line in the periodic table, such as Li^+ and Mg^{2+} or Ca^{2+} and Y^{3+} , have different sizes and charges but similar charge-to-radius ratios. As a result, these pairs of metal ions have similar effects on the acidity of coordinated water molecules, and they often exhibit other significant similarities in chemistry as well.

Note the Pattern

Solutions of small, highly charged metal ions in water are acidic.

Reactions such as those discussed in this section, in which a salt reacts with water to give an acidic or basic solution, are often called hydrolysis reactions. A chemical reaction in which a salt reacts with water to yield an acidic or a basic solution. Using a

separate name for this type of reaction is unfortunate because it suggests that they are somehow different. In fact, hydrolysis reactions are just acid–base reactions in which the acid is a cation or the base is an anion; they obey the same principles and rules as all other acid–base reactions.

Note the Pattern

A hydrolysis reaction is an acid–base reaction.

Example 16.2.3

Predict whether aqueous solutions of these compounds are acidic, basic, or neutral.

1. KNO_3
2. $\text{CrBr}_3 \cdot 6\text{H}_2\text{O}$
3. Na_2SO_4

Given: compound

Asked for: acidity or basicity of aqueous solution

Strategy:

A Assess the acid–base properties of the cation and the anion. If the cation is a weak Lewis acid, it will not affect the pH of the solution. If the cation is the conjugate acid of a weak base or a relatively highly charged metal cation, however, it will react with water to produce an acidic solution.


B If the anion is the conjugate base of a strong acid, it will not affect the pH of the solution. If, however, the anion is the conjugate base of a weak acid, the solution will be basic.

Solution:

1. **A** The K^+ cation has a small positive charge (+1) and a relatively large radius (because it is in the fourth row of the periodic table), so it is a very weak Lewis acid.

B The NO_3^- anion is the conjugate base of a strong acid, so it has essentially no basic character (Table 16.1). Hence neither the cation nor the anion will react with water to produce H^+ or OH^- , and the solution will be neutral.

2. **A** The Cr^{3+} ion is a relatively highly charged metal cation that should behave similarly to the Al^{3+} ion and form the $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$ complex, which will behave as a weak acid:



B The Br^- anion is a very weak base (it is the conjugate base of the strong acid HBr), so it does not affect the pH of the solution. Hence the solution will be acidic.

3. **A** The Na^+ ion, like the K^+ , is a very weak acid, so it should not affect the acidity of the solution.

B In contrast, SO_4^{2-} is the conjugate base of HSO_4^- , which is a weak acid. Hence the SO_4^{2-} ion will react with water as shown in Figure 16.6 to give a slightly basic solution.

Exercise

Predict whether aqueous solutions of the following are acidic, basic, or neutral.

1. KI
2. $\text{Mg}(\text{ClO}_4)_2$
3. NaHS

Answer

1. neutral
2. acidic
3. basic (due to the reaction of HS^- with water to form H_2S and OH^-)

Summary

Two species that differ by only a proton constitute a **conjugate acid–base pair**. The magnitude of the equilibrium constant for an ionization reaction can be used to determine the relative strengths of acids and bases. For an aqueous solution of a weak acid, the dissociation constant is called the **acid ionization constant (K_a)**. Similarly, the equilibrium constant for the reaction of a weak base with water is the **base ionization constant (K_b)**. For any conjugate acid–base pair, $K_a K_b = K_w$. Smaller values of pK_a correspond to larger acid ionization constants and hence stronger acids. Conversely, smaller values of pK_b correspond to larger base ionization constants and hence stronger bases. At 25°C, $pK_a + pK_b = 14.00$. Acid–base reactions always proceed in the direction that produces the weaker acid–base pair. No acid stronger than H_3O^+ and no base stronger than OH^- can exist in aqueous solution, leading to the phenomenon known as the **leveling effect**. Polyprotic acids (and bases) lose (and gain) protons in a stepwise manner, with the fully protonated species being the strongest acid and the fully deprotonated species the strongest base. A salt can dissolve in water to produce a neutral, a basic, or an acidic solution, depending on whether it contains the conjugate base of a weak acid as the anion (A^-), the conjugate acid of a weak base as the cation (BH^+), or both. Salts that contain small, highly charged metal ions produce acidic solutions in water. The reaction of a salt with water to produce an acidic or a basic solution is called a **hydrolysis reaction**.

Key Takeaways

- Acid–base reactions always contain two conjugate acid–base pairs.
- Each acid and each base has an associated ionization constant that corresponds to its acid or base strength.

Key Equations

Acid ionization constant

Equation 16.2.13: $K_a = K[H_2O] = \frac{[H_3O^+][A^-]}{[HA]}$

Base ionization constant

Equation 16.2.15: $K_b = K[H_2O] = [BH^+][OH^-][B]$

Relationship between K_a and K_b of a conjugate acid–base pair

Equation 16.2.20: $K_a K_b = K_w$

Definition of pK_a

Equation 16.2.21: $pK_a = -\log_{10} K_a$

Equation 16.2.22: $K_a = 10^{-pK_a}$

Definition of pK_b

Equation 16.2.23: $pK_b = -\log_{10} K_b$

Equation 16.2.25: $K_b = 10^{-pK_b}$

Relationship between pK_a and pK_b of a conjugate acid–base pair

Equation 16.2.25: $pK_a + pK_b = pK_w$

Equation 16.29: $pK_a + pK_b = 14.00$ at 25 °C


Conceptual Problems

1. Identify the conjugate acid–base pairs in each equilibrium.

1. $HSO_4^-(aq) + H_2O(l) \rightleftharpoons SO_4^{2-}(aq) + H_3O^+(aq)$
2. $C_3H_7NO_2(aq) + H_3O^+(aq) \rightleftharpoons C_3H_8NO_2^+(aq) + H_2O(l)$
3. $CH_3CO_2H(aq) + NH_3(aq) \rightleftharpoons CH_3CO_2^-(aq) + NH_4^+(aq)$
4. $SbF_5(aq) + 2HF(aq) \rightleftharpoons H_2F^+(aq) + SbF_6^-(aq)$


2. Identify the conjugate acid–base pairs in each equilibrium.

1. $HF(aq) + H_2O(l) \rightleftharpoons H_3O^+(aq) + F^-(aq)$
2. $CH_3CH_2NH_2(aq) + H_2O(l) \rightleftharpoons CH_3CH_2NH_3^+(aq) + OH^-(aq)$




3. Salts such as NaH contain the hydride ion (H^-). When sodium hydride is added to water, it produces hydrogen gas in a highly vigorous reaction. Write a balanced chemical equation for this reaction and identify the conjugate acid–base pairs.


4. Write the expression for K_a for each reaction.



5. Write an expression for the ionization constant K_b for each reaction.



6. Predict whether each equilibrium lies primarily to the left or to the right.



7. Species that are strong bases in water, such as CH_3^- , NH_2^- , and S^{2-} , are leveled to the strength of OH^- , the conjugate base of H_2O . Because their relative base strengths are indistinguishable in water, suggest a method for identifying which is the strongest base. How would you distinguish between the strength of the acids HIO_3 , H_2SO_4 , and $HClO_4$?

8. Is it accurate to say that a 2.0 M solution of H_2SO_4 , which contains two acidic protons per molecule, is 4.0 M in H^+ ? Explain your answer.

9. The alkalinity of soil is defined by the following equation: $\text{alkalinity} = [HCO_3^-] + 2[CO_3^{2-}] + [OH^-] - [H^+]$. The source of both HCO_3^- and CO_3^{2-} is H_2CO_3 . Explain why the basicity of soil is defined in this way.

10. Why are aqueous solutions of salts such as $CaCl_2$ neutral? Why is an aqueous solution of $NaNH_2$ basic?

11. Predict whether aqueous solutions of the following are acidic, basic, or neutral.

1. Li_3N
2. NaH
3. KBr
4. $C_2H_5NH_3^+Cl^-$

12. When each compound is added to water, would you expect the pH of the solution to increase, decrease, or remain the same?


1. $LiCH_3$
2. $MgCl_2$
3. K_2O
4. $(CH_3)_2NH_2^+Br^-$




13. Which complex ion would you expect to be more acidic— $Pb(H_2O)_4^{2+}$ or $Sn(H_2O)_4^{2+}$? Why?

14. Would you expect $Sn(H_2O)_4^{2+}$ or $Sn(H_2O)_6^{4+}$ to be more acidic? Why?

15. Is it possible to arrange the hydrides LiH , RbH , KH , CsH , and NaH in order of increasing base strength in aqueous solution? Why or why not?

Answer





- 2.
- 3.
- 4.
- 5.
- 6.
- 7.
- 8.
- 9.
- 10.
- 11.
- 12.
- 13.
- 14.
- 15.

Numerical Problems

1. Arrange these acids in order of increasing strength.

- acid A: $pK_a = 1.52$
- acid B: $pK_a = 6.93$
- acid C: $pK_a = 3.86$

Given solutions with the same initial concentration of each acid, which would have the highest percent ionization?

2. Arrange these bases in order of increasing strength:

- base A: $pK_b = 13.10$
- base B: $pK_b = 8.74$
- base C: $pK_b = 11.87$

Given solutions with the same initial concentration of each base, which would have the highest percent ionization?

3. Calculate the K_a and the pK_a of the conjugate acid of a base with each pK_b value.

1. 3.80
2. 7.90
3. 13.70
4. 1.40
5. -2.50

4. Benzoic acid is a food preservative with a pK_a of 4.20. Determine the K_b and the pK_b for the benzoate ion.

5. Determine K_a and pK_a of boric acid $[B(OH)_3]$, solutions of which are occasionally used as an eyewash; the pK_b of its conjugate base is 4.80.

Answers

1. acid B < acid C < acid A (strongest)

2.

3. 1. $K_a = 6.3 \times 10^{-11}$; $pK_a = 10.20$
2. $K_a = 7.9 \times 10^{-7}$; $pK_a = 6.10$

3. $K_a = 0.50$; $pK_a = 0.30$

4. $K_a = 2.5 \times 10^{-13}$; $pK_a = 12.60$

5. $K_a = 3.2 \times 10^{-17}$; $pK_a = 16.50$

4.

5. $K_a = 6.3 \times 10^{-10}$ $pK_a = 9.20$

Contributors

- Anonymous

Modified by Joshua B. Halpern

This page titled [16.2: A Qualitative Description of Acid-Base Equilibria](#) is shared under a [CC BY-NC-SA 3.0](#) license and was authored, remixed, and/or curated by [Anonymous](#).