

11.4: Enthalpy Change is a Measure of the Heat Evolved or Absorbed

When physical or chemical changes occur, they are generally accompanied by a transfer of energy. The **law of conservation of energy** states that in any physical or chemical process, energy is neither created nor destroyed. In other words, the entire energy in the universe is conserved. In order to better understand the energy changes taking place during a reaction, we need to define two parts of the universe, called the system and the surroundings. The **system** is the specific portion of matter in a given space that is being studied during an experiment or an observation. The **surroundings** are everything in the universe that is not part of the system. In practical terms for a laboratory chemist, the system is the particular chemicals being reacted, while the surroundings is the immediate vicinity within the room. During most processes, energy is exchanged between the system and the surroundings. If the system loses a certain amount of energy, that same amount of energy is gained by the surroundings. If the system gains a certain amount of energy, that energy is supplied by the surroundings.

A chemical reaction or physical change is **endothermic** if heat is absorbed by the system from the surroundings. In the course of an endothermic process, the system gains heat from the surroundings and so the temperature of the surroundings decreases. The quantity of heat for a process is represented by the letter q . The sign of q for an endothermic process is positive because the system is gaining heat. A chemical reaction or physical change is **exothermic** if heat is released by the system into the surroundings. Because the surroundings are gaining heat from the system, the temperature of the surroundings increases. The sign of q for an exothermic process is negative because the system is losing heat.




Figure 11.4.1: (A) Endothermic reaction. (B) Exothermic reaction.

Enthalpy


Heat changes in chemical reactions are often measured in the laboratory under conditions in which the reacting system is open to the atmosphere. In that case, the system is at a constant pressure. **Enthalpy** (H) is the heat content of a system at constant pressure. Chemists routinely measure changes in enthalpy of chemical systems as reactants are converted into products. The heat that is absorbed or released by a reaction at constant pressure is the same as the enthalpy change, and is given the symbol ΔH . Unless otherwise specified, all reactions in this material are assumed to take place at constant pressure.

The change in enthalpy of a reaction is a measure of the differences in enthalpy of the reactants and products. The enthalpy of a system is determined by the energies needed to break chemical bonds and the energies needed to form chemical bonds. Energy needs to be put into the system in order to break chemical bonds, as they do not come apart spontaneously in most cases. Bond formation to produce products will involve release of energy. The change in enthalpy shows the trade-offs made in these two processes. Does it take more energy to break bonds than that needed to form bonds? If so, the reaction is endothermic and the enthalpy change is positive. If more energy is produced in bond formation than that needed for bond breaking, the reaction is exothermic and the enthalpy is negative.

Several factors influence the enthalpy of a system. Enthalpy is an extensive property, determined in part by the amount of material we work with. The state of reactants and products (solid, liquid, or gas) influences the enthalpy value for a system. The direction of the reaction affects the enthalpy value. A reaction that takes place in the opposite direction has the same numerical enthalpy value, but the opposite sign.

Thermochemical Equation

When methane gas is combusted, heat is released, making the reaction exothermic. Specifically, the combustion of 1 mol of methane releases 890.4 kilojoules of heat energy. This information can be shown as part of the balanced equation:



The equation tells us that 1 mol of methane combines with 2 mol of oxygen to produce 1 mol of carbon dioxide and 2 mol of water. In the process, 890.4 kJ is released and so it is written as a product of the reaction. A **thermochemical equation** is a chemical equation that includes the enthalpy change of the reaction. The process in the above thermochemical equation can be shown visually in Figure 11.4.2






Figure 11.4.2: (A) As reactants are converted to products in an exothermic reaction, enthalpy is released into the surroundings. The enthalpy change of the reaction is negative. (B) As reactants are converted to products in an endothermic reaction, enthalpy is absorbed from the surroundings. The enthalpy change of the reaction is positive.

In the combustion of methane example, the enthalpy change is negative because heat is being released by the system. Therefore, the overall enthalpy of the system decreases. The **heat of reaction** is the enthalpy change for a chemical reaction. In the case above, the heat of reaction is -890.4 kJ . The thermochemical reaction can also be written in this way:




Heats of reaction are typically measured in kilojoules. It is important to include the physical states of the reactants and products in a thermochemical equation as the value of the ΔH depends on those states.


Endothermic reactions absorb energy from the surroundings as the reaction occurs. When 1 mol of calcium carbonate decomposes into 1 mol of calcium oxide and 1 mol of carbon dioxide, 177.8 kJ of heat is absorbed. The process is shown visually in Figure 11.4.2B. The thermochemical reaction is shown below.



Because the heat is absorbed by the system, the 177.8 kJ is written as a reactant. The heat of reaction is positive for an endothermic reaction.



The way in which a reaction is written influences the value of the enthalpy change for the reaction. Many reactions are reversible, meaning that the product(s) of the reaction are capable of combining and reforming the reactant(s). If a reaction is written in the reverse direction, the sign of the ΔH changes. For example, we can write an equation for the reaction of calcium oxide with carbon dioxide to form calcium carbonate.



The reaction is exothermic and thus the sign of the enthalpy change is negative.




Stoichiometric Calculations and Enthalpy Changes

Chemistry problems that involve enthalpy changes can be solved by techniques similar to stoichiometry problems. Refer again to the combustion reaction of methane. Since the reaction of 1 mol of methane released 890.4 kJ, the reaction of 2 mol of methane would release $2 \times 890.4 \text{ kJ} = 1781 \text{ kJ}$. The reaction of 0.5 mol of methane would release $\frac{890.4 \text{ kJ}}{2} = 445.2 \text{ kJ}$. As with other stoichiometry problems, the moles of a reactant or product can be linked to mass or volume.

✓ Example 11.4.1

Sulfur dioxide gas reacts with oxygen to form sulfur trioxide in an exothermic reaction, according to the following thermochemical equation.



Calculate the enthalpy change that occurs when 58.0 g of sulfur dioxide is reacted with excess oxygen.

Solution

Step 1: List the known quantities and plan the problem.

- Mass $\text{SO}_2 = 58.0 \text{ g}$
- Molar mass $\text{SO}_2 = 64.07 \text{ g/mol}$
- $\Delta H = -198 \text{ kJ}$ for the reaction of 2 mol SO_2

Unknown

- $\Delta H = ? \text{ kJ}$

The calculation requires two steps. The mass of SO_2 is converted to moles. Then the moles of SO_2 is multiplied by the conversion factor of $\left(\frac{-198 \text{ kJ}}{2 \text{ mol SO}_2} \right)$.

Step 2: Solve.

$$\Delta H = 58.0 \text{ g SO}_2 \times \frac{1 \text{ mol SO}_2}{64.07 \text{ g SO}_2} \times \frac{-198 \text{ kJ}}{2 \text{ mol SO}_2} = 89.6 \text{ kJ}$$

Step 3: Think about your result.

The mass of sulfur dioxide is slightly less than 1 mol. Since 198 kJ is released for every 2 mol of SO_2 that reacts, the heat released when about 1 mol reacts is one half of 198. The 89.6 kJ is slightly less than half of 198. The sign of ΔH is negative because the reaction is exothermic.

This page titled [11.4: Enthalpy Change is a Measure of the Heat Evolved or Absorbed](#) is shared under a [mixed](#) license and was authored, remixed, and/or curated by [Anonymous](#).

- [8.8: Enthalpy Change is a Measure of the Heat Evolved or Absorbed](#) by Henry Agnew, Marisa Alviar-Agnew is licensed [CK-12](#). Original source: <https://www.ck12.org/c/chemistry/>.