

## 8.1: Solids, Liquids, and Gases- A Molecular Comparison

### Learning Objectives

- To be familiar with the kinetic molecular description of solids, liquids, and gases.

The *physical* properties of a substance depends upon its physical state. Water vapor, liquid water and ice all have the same *chemical* properties, but their *physical* properties are considerably different. In general *covalent bonds* determine: molecular shape, bond energies, *chemical* properties, while *intermolecular forces* (non-covalent bonds) influence the *physical* properties of liquids and solids. The kinetic molecular theory of gases gives a reasonably accurate description of the behavior of gases. A similar model can be applied to liquids, but it must take into account the nonzero volumes of particles and the presence of strong intermolecular attractive forces.



Figure 8.1.1: The three common states of matter. From the left, they are solid, liquid, and gas, represented by an ice sculpture, a drop of water, and the air around clouds, respectively. Images used with permission from Wikipedia.

The *state* of a substance depends on the balance between the *kinetic energy* of the individual particles (molecules or atoms) and the *intermolecular forces*. The kinetic energy keeps the molecules apart and moving around, and is a function of the temperature of the substance. The intermolecular forces are attractive forces that try to draw the particles together (Figure 8.1.2). As discussed previously, gasses are very sensitive to temperatures and pressure. However, these also affect liquids and solids too. Heating and cooling can change the *kinetic energy* of the particles in a substance, and so, we can change the physical state of a substance by heating or cooling it. Increasing the pressure on a substance forces the molecules closer together, which *increases* the strength of intermolecular forces.




Figure 8.1.2: Molecular level picture of gases, liquids and solids.

Below is an overview of the general properties of the three different phases of matter.

### Properties of Gases

- A collection of widely separated molecules
- The kinetic energy of the molecules is greater than any attractive forces between the molecules
- The lack of any significant attractive force between molecules allows a gas to expand to fill its container
- If attractive forces become large enough, then the gases exhibit [non-ideal behavior](#)

### Properties of Liquids

- The intermolecular attractive forces are strong enough to hold molecules close together
- Liquids are more dense and less compressible than gasses
- Liquids have a definite volume, independent of the size and shape of their container

- The attractive forces are **not** strong enough, however, to keep neighboring molecules in a fixed position and molecules are free to move past or slide over one another

┆ *Thus, liquids can be poured and assume the shape of their containers.*

### Properties of Solids

- The intermolecular forces between neighboring molecules are strong enough to keep them locked in position
- Solids (like liquids) are not very compressible due to the lack of space between molecules
- If the molecules in a solid adopt a highly ordered packing arrangement, the structures are said to be **crystalline**

┆ *Due to the strong intermolecular forces between neighboring molecules, solids are rigid.*

### Contributors and Attributions

- [Mike Blaber](#) ([Florida State University](#))

---

This page titled [8.1: Solids, Liquids, and Gases- A Molecular Comparison](#) is shared under a [mixed](#) license and was authored, remixed, and/or curated by [Anonymous](#).

- [11.1: A Molecular Comparison of Gases, Liquids, and Solids](#) is licensed [CC BY-NC-SA 3.0](#).