


Mechanism of Vision

The molecule cis-retinal can absorb light at a specific wavelength. When visible light hits the cis-retinal, the cis-retinal undergoes an **isomerization**, or change in molecular arrangement, to all-trans-retinal. The new form of trans-retinal does not fit as well into the protein, and so a series of geometry changes in the protein begins. The resulting complex is referred to as bathorhodopsin (there are other intermediates in this process, but we'll ignore them for now).



The reaction above shows Lysine side-chain from the opsin react with 11-cis-retinal when stimulated. By removing the oxygen atom from the retinal and two hydrogen atom from the free amino group of the lysine, the linkage show on the picture above is formed, and it is called Schiff base.

Signal Transduction Pathway

As the protein changes its geometry, it initiates a cascade of biochemical reactions that results in changes in charge so that a large potential difference builds up across the plasma membrane. This potential difference is passed along to an adjoining nerve cell as an electrical impulse. The nerve cell carries this impulse to the brain, where the visual information is interpreted.

The light image is mapped on the surface of the retina by activating a series of light-sensitive cells known as rods and cones or photoreceptors. The rods and cones convert the light into electrical impulses which are transmitted to the brain via nerve fibers. The brain then determines, which nerve fibers carried the electrical impulse activate by light at certain photoreceptors, and then creates an image.

The retina is lined with many millions of photoreceptor cells that consist of two types: 7 million cones provide color information and sharpness of images, and 120 million rods are extremely sensitive detectors of white light to provide night vision. The tops of the rods and cones contain a region filled with membrane-bound discs, which contain the molecule cis-retinal bound to a protein called opsin. The resulting complex is called rhodopsin or "visual purple".



In human eyes, rod and cones react to light stimulation, and a series of chemical reactions happen in cells. These cells receive light, and pass on signals to other receiver cells. This chain of process is class signal transduction pathway. Signal transduction pathway is a mechanism that describe the ways cells react and response to stimulation.

References

1. Biochemistry, L. Stryer (W.H. Freeman and Co, San Francisco, 1975).
2. *The Cambridge Guide to the Material World*, Rodney Cotterill (Cambridge University Press, Cambridge, 1985)

Contributors

- {{template.ContribOphardt()}}

15.3: Vision is shared under a [CC BY-NC-SA 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/) license and was authored, remixed, and/or curated by LibreTexts.

- [Vision and Light](#) is licensed [CC BY-NC-SA 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/).