

9.16: ADDITION OF CARBENES TO ALKENES - CYCLOPROPANE SYNTHESIS

Learning Objective

- predict the products of carbene additions to alkenes


INTRODUCTION

Carbenes were once only thought of as short lived intermediates. The reactions of this section only deal with these short lived carbenes which are mostly prepared in situ, in conjunction with the main reaction. However, there do exist so called persistent carbenes. These persistent carbenes are stabilized by a variety of methods often including aromatic rings or transition metals. In general a carbene is neutral and has 6 valence electrons, 2 of which are non bonding. These electrons can either occupy the same sp^2 hybridized orbital to form a singlet carbene (with paired electrons), or two different sp^2 orbitals to form a triplet carbene (with unpaired electrons). The chemistry of triplet and singlet carbenes is quite different but can be oversimplified to the statement: singlet carbenes usually retain stereochemistry while triplet carbenes do not. The carbenes discussed in this section are singlet and thus retain stereochemistry.

The reactivity of a singlet carbene is concerted and similar to that of electrophilic or nucleophilic addition (although, triplet carbenes react like biradicals, explaining why stereochemistry is not retained). The highly reactive nature of carbenes leads to very fast reactions in which the rate determining step is generally carbene formation.


PREPARATION OF METHYLENE

The preparation of methylene starts with the yellow gas diazomethane, CH_2N_2 . Diazomethane can be exposed to light, heat or copper to facilitate the loss of nitrogen gas and the formation of the simplest carbene methylene. The process is driven by the formation of the nitrogen gas which is a very stable molecule.




CARBENE REACTION WITH ALKENES

A carbene such as methylene will react with an alkene which will break the double bond and result with a cyclopropane. The reaction will usually leave stereochemistry of the double bond unchanged. As stated before, carbenes are generally formed along with the main reaction; hence the starting material is diazomethane not methylene.



In the above case *cis*-2-butene is converted to *cis*-1,2-dimethylcyclopropane. Likewise, below the *trans* configuration is maintained.




ADDITIONAL TYPES OF CARBENES AND CARBENOIDS

In addition to the general carbene with formula R_2C there exist a number of other compounds that behave in much the same way as carbenes in the synthesis of cyclopropane. **Halogenated carbenes** are formed from halomethanes. An example is dichlorocarbene, Cl_2C . These halogenated carbenes will form cyclopropanes in the same manner as methylene but with the interesting presence of two halogen atoms in place of the hydrogen atoms.


Carbenoids are substances that form cyclopropanes like carbenes but are not technically carbenes. One common example is the stereospecific Simmon-Smith reaction which utilizes the carbenoid ICH_2ZnI . The carbenoid is formed in situ via the mixing of a Zn-Cu couple with CH_2I_2 . Since this reacts the same as a carbene, the same methods can be applied to determine the product. An example of this is given as problem 5.

Exercise 9.16.1


1. Knowing that cycloalkenes react much the same as regular alkenes what would be the expected structure of the product of cyclohexene and diazomethane facilitated by copper metal?
2. What would be the result of a Simmons-Smith reaction that used *trans*-3-pentene as a reagent?
3. What starting material could be used to form *cis*-1,2-diethylcyclopropane?
4. What would the following reaction yield?



5. Draw the product of this reaction. What type of reaction is this?




6. Predict the following products. Will they be the same product?




Answer


1. The product will be a bicyclic ring, Bicyclo[4.1.0]heptane.




2. The stereochemistry will be retained making a cyclopropane with *trans* methyl and ethyl groups. *Trans*-1-ethyl-2-methylcyclopropane
3. The *cis* configuration will be maintained from reagent to product so we would want to start with *cis*-3-hexene. A Simmons Smith reagent, or methylene could be used as the carbene or carbenoid.
4. The halogenated carbene will react the same as methylene yielding, *cis*-1,1-dichloro-2,3dimethylcyclopropane.



5. This is a Simmons-Smith reaction which uses the carbenoid formed by the CH_2I_2 and Zn-Cu. The reaction results in the same product as if methylene was used and retains stereospecificity. Iodine metal and the Zn-Cu are not part of the product. The product is *trans*-1,2-ethyl-methylcyclopropane.



6. No they will not be the same product, they will be isomers of each other.



REFERENCES

1. Vollhardt, K. Peter C. and Schore, Neil E. Organic Chemistry: Structure and Function. New York: Bleyer, Brennan, 2007.
2. Abdel-Wahab, Aboel-Magd A. Ahmed, Saleh A. and Dürr, Heinz. "Carbene Formation by Extrusion of Nitrogen" in CRC Handbook of Organic Photochemistry and Photobiology. CRC Press, 2004.

CONTRIBUTORS AND ATTRIBUTIONS

- Dr. Dietmar Kennepohl FCIC (Professor of Chemistry, [Athabasca University](#))
- Prof. Steven Farmer ([Sonoma State University](#))
- Paul Tisher

9.16: Addition of Carbenes to Alkenes - Cyclopropane Synthesis is shared under a [not declared](#) license and was authored, remixed, and/or curated by LibreTexts.