

11.6: IUPAC Nomenclature

Learning Objectives

- To name alkanes by the IUPAC system and write formulas for alkanes given IUPAC names

As noted in previously, the number of isomers increases rapidly as the number of carbon atoms increases. There are 3 pentanes, 5 hexanes, 9 heptanes, and 18 octanes. It would be difficult to assign unique individual names that we could remember. A systematic way of naming hydrocarbons and other organic compounds has been devised by the International Union of Pure and Applied Chemistry (IUPAC). These rules, used worldwide, are known as the IUPAC System of Nomenclature. (Some of the names we used earlier, such as isobutane, isopentane, and neopentane, do not follow these rules and are called *common names*.) A stem name (Table 11.6.1) indicates the number of carbon atoms in the longest continuous chain (LCC). Atoms or groups attached to this carbon chain, called *substituents*, are then named, with their positions indicated by numbers. For now, we will consider only those substituents called alkyl groups.

Table 11.6.1: Stems That Indicate the Number of Carbon Atoms in Organic Molecules

Stem	Number
meth-	1
eth-	2
prop-	3
but-	4
pent-	5
hex-	6
hept-	7
oct-	8
non-	9
dec-	10

An alkyl group is a group of atoms that results when one hydrogen atom is removed from an alkane. The group is named by replacing the *-ane* suffix of the parent hydrocarbon with *-yl*. For example, the -CH_3 group derived from methane (CH_4) results from subtracting one hydrogen atom and is called a *methyl group*. The alkyl groups we will use most frequently are listed in Table 11.6.2 Alkyl groups are not independent molecules; they are parts of molecules that we consider as a unit to name compounds systematically.

Table 11.6.2: Common Alkyl Groups

Parent Alkane		Alkyl Group		Condensed Structural Formula
methane	$\begin{array}{c} \text{H} \\ \\ \text{H}-\text{C}-\text{H} \\ \\ \text{H} \end{array}$	methyl	$\begin{array}{c} \text{H} \\ \\ \text{H}-\text{C}- \\ \\ \text{H} \end{array}$	CH ₃ —


*There are four butyl groups, two derived from butane and two from isobutane. We will introduce the other three where appropriate.

Parent Alkane		Alkyl Group		Condensed Structural Formula
ethane		ethyl		CH_3CH_2-
propane		propyl		$\text{CH}_3\text{CH}_2\text{CH}_2-$
		isopropyl		$(\text{CH}_3)_2\text{CH}-$
butane		butyl*		$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2-$

*There are four butyl groups, two derived from butane and two from isobutane. We will introduce the other three where appropriate.

Simplified IUPAC rules for naming alkanes are as follows (demonstrated in Example 11.6.1).

- Name alkanes according to the LCC (longest continuous chain) of carbon atoms in the molecule (rather than the total number of carbon atoms).** This LCC, considered the parent chain, determines the base name, to which we add the suffix *-ane* to indicate that the molecule is an alkane.
- If the hydrocarbon is branched, number the carbon atoms of the LCC.** Numbers are assigned in the direction that gives the lowest numbers to the carbon atoms with attached substituents. Hyphens are used to separate numbers from the names of substituents; commas separate numbers from each other. (The LCC need not be written in a straight line; for example, the LCC in the following has five carbon atoms.)




- Place the names of the substituent groups in alphabetical order before the name of the parent compound.** If the same alkyl group appears more than once, the numbers of all the carbon atoms to which it is attached are expressed. If the same group appears more than once on the same carbon atom, the number of that carbon atom is repeated as many times as the group appears. Moreover, the number of identical groups is indicated by the Greek prefixes *di-*, *tri-*, *tetra-*, and so on. These prefixes are *not* considered in determining the alphabetical order of the substituents. For example, ethyl is listed before dimethyl; the *di-* is simply ignored. The last alkyl group named is prefixed to the name of the parent alkane to form one word.



When these rules are followed, every unique compound receives its own exclusive name. The rules enable us to not only name a compound from a given structure but also draw a structure from a given name. The best way to learn how to use the IUPAC system is to put it to work, not just memorize the rules. It's easier than it looks.

✓ Example 11.6.1

Name each compound.

1.








Solution

1. The LCC has five carbon atoms, and so the parent compound is pentane (rule 1). There is a methyl group (rule 2) attached to the second carbon atom of the pentane chain. The name is therefore 2-methylpentane.
2. The LCC has six carbon atoms, so the parent compound is hexane (rule 1). Methyl groups (rule 2) are attached to the second and fifth carbon atoms. The name is 2,5-dimethylhexane.
3. The LCC has eight carbon atoms, so the parent compound is octane (rule 1). There are methyl and ethyl groups (rule 2), both attached to the fourth carbon atom (counting from the *right* gives this carbon atom a lower number; rule 3). The correct name is thus 4-ethyl-4-methyloctane.

? Exercise 11.6.1

Name each compound.

1.



✓ Example 11.6.2


Draw the structure for each compound.

- a. 2,3-dimethylbutane
- b. 4-ethyl-2-methylheptane


Solution

In drawing structures, always start with the parent chain.


- a. The parent chain is butane, indicating four carbon atoms in the LCC.




Then add the groups at their proper positions. You can number the parent chain from either direction as long as you are consistent; just don't change directions before the structure is done. The name indicates two methyl (CH_3) groups, one on the second carbon atom and one on the third.




Finally, fill in all the hydrogen atoms, keeping in mind that each carbon atom must have four bonds.



- The parent chain is heptane in this case, indicating seven carbon atoms in the LCC. $-C-C-C-C-C-C-C-$
Adding the groups at their proper positions gives



Filling in all the hydrogen atoms gives the following condensed structural formulas:



Note that the bonds (dashes) can be shown or not; sometimes they are needed for spacing.

? Exercise 11.6.2

Draw the structure for each compound.

- 4-ethyloctane
- 3-ethyl-2-methylpentane
- 3,3,5-trimethylheptane

Key Takeaway

- Alkanes have both common names and systematic names, specified by IUPAC.

This page titled [11.6: IUPAC Nomenclature](#) is shared under a [CC BY-NC-SA 4.0](#) license and was authored, remixed, and/or curated by [Eden Francis](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

- [12.5: IUPAC Nomenclature](#) by Anonymous is licensed [CC BY-NC-SA 3.0](#). Original source: <https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological>.