

2.2: Internal Energy

The internal energy of a system is identified with the random, disordered motion of molecules; the total (internal) energy in a system includes potential and kinetic energy. This is contrast to external energy which is a function of the sample with respect to the outside environment (e.g. kinetic energy if the sample is moving or potential energy if the sample is at a height from the ground etc). The symbol for Internal Energy Change is ΔU .

Energy on a smaller scale

- Internal energy includes energy on a microscopic scale
- It is the sum of all the microscopic energies such as:
 1. translational kinetic energy
 2. vibrational and rotational kinetic energy
 3. potential energy from intermolecular forces

Example

One gram of water at zero °Celsius compared with one gram of copper at zero °Celsius do NOT have the same internal energy because even though their kinetic energies are equal, water has a much higher potential energy causing its internal energy to be much greater than the copper's internal energy.

Internal Energy Change Equations

The [first law of thermodynamics](#) states:

$$dU = dq + dw \quad (2.2.1)$$

where dq is heat and dw is work.


An [isolated system](#) cannot exchange heat or work with its surroundings making the change in internal energy equal to zero:

$$dU_{\text{isolated system}} = 0 \quad (2.2.2)$$

Therefore, in an isolated system:

$$dq = -dw \quad (2.2.3)$$

Energy is Conserved



$$dU_{\text{isolated system}} = dU_{\text{system}} + dU_{\text{surroundings}} \quad (2.2.4)$$

$$dU_{\text{system}} = -dU_{\text{surroundings}} \quad (2.2.5)$$

The signs of internal energy

- Energy *entering* the system is **POSITIVE (+)**, meaning heat is *absorbed*, $q > 0$. Work is thus done *on* the system, $w > 0$
- Energy *leaving* the system is **NEGATIVE (-)**, meaning heat is *given off* by the system, $q < 0$ and work is done *by* the system, $w < 0$

Quick Notes

- A system *contains* ONLY Internal Energy
- A system does NOT *contain* energy in the form of heat or work
- Heat and work only exist during a change in the system; they are path functions
- Internal energy is a state function

Outside Links

- Levine, Ira N. "Thermodynamic internal energy of an ideal gas of rigid rotors." J. Chem. Educ. **1985**: 62, 53.

Contributors

- Lorraine Alborzfar (UCD)

2.2: Internal Energy is shared under a [not declared](#) license and was authored, remixed, and/or curated by LibreTexts.