

12.S: STRUCTURE DETERMINATION - MASS SPECTROMETRY AND INFRARED SPECTROSCOPY (SUMMARY)

CONCEPTS & VOCABULARY

12.1 Introduction

- Spectroscopy describes several techniques used by chemists to understand chemical structures and bonds.

12.2 Mass Spectrometry of Small Molecules - Magnetic Sector Instruments

- Mass spectrometers consist of an **ion source**, **mass analyzer** and detector.
- There are several common **ion sources** including **electron ionization** and **chemical ionization**.
- Upon ionization, a molecular ion is formed (the molecule after losing a single electron) which will break into smaller pieces (fragments).
- Fragments that are charged will appear in the mass spectrum and are helpful in identifying the parent molecule.
- The most abundant ion in a mass spectrum is called the **base peak**.
- The ion with the same mass as the parent molecule is called the **molecular ion**.
- Isotopes of carbon and hydrogen lead to common M+1 peaks.
- The x-axis of a mass spectrum is **m/z** - the mass to charge ratio, which in practice equals the mass of the ion.

12.3 Interpreting Mass Spectra

- Uncharged particles do not appear in mass spectra.
- The y-axis of a mass spectrum is the relative abundance, with the base peak set at 100 as the most abundant ion.
- Abundance of ions is related to their stability.

12.4 Mass Spectrometry of Some Common Functional Groups

12.5 Mass Spectrometry in Biological - Time-of-flight (TOF) Instruments

12.6 Spectroscopy and the Electromagnetic Spectrum

- Electromagnetic radiation is composed of waves where shorter wavelengths correspond to higher energy radiation.
- Electromagnetic radiation can also be thought of as a stream of particles called **photons**.
- The electromagnetic spectrum is made up of many types of radiation including infrared, ultraviolet, and visible lights as well as x-rays, gamma rays, microwaves, and radio waves.
- Molecular spectroscopy works by exposing a chemical sample to electromagnetic radiation. It will only absorb radiation with energy that corresponds to some excited state, while all other energies will pass through unabsorbed.

12.7 Infrared Spectroscopy

- When infrared radiation is absorbed, molecules will move to a higher vibrational energy state.
- Examples of molecular vibrations include bending and stretching of bonds. These vibrations can be symmetric or asymmetric.
- In general, more polar bonds have stronger IR absorption.
- IR spectra typically use wavenumbers (cm^{-1}) as units for the x-axis.
- The y-axis for IR spectra is usually % transmittance, with 100% at the top of the spectrum and absorbances looking like valleys (or downward peaks).

12.8 Interpreting Infrared Spectra

- Functional groups have standard regions within the IR spectrum where they absorb.
- The general regions include hydrogen bonding (O-H and N-H), carbon-hydrogen bonds, triple bonds, carbonyls, alkenes, and fingerprint region.

12.9 Infrared Spectra of Some Common Functional Groups

SKILLS TO MASTER

- Skill 12.1 Determine specific atoms from mass spectra based on molecular ion and M+2 peaks (N, Cl, Br).
- Skill 12.2 Interpret mass spectra fragments - recognizing common fragments.
- Skill 12.3 Interpret infrared spectra to determine functional groups that are present or absent.

MEMORIZATION TASKS (MT)

- MT 12.1 Memorize common mass spectra fragments.
 - MT 12.2 Memorize common functional group regions in infrared spectroscopy.
-

This page titled [12.S: Structure Determination - Mass Spectrometry and Infrared Spectroscopy \(Summary\)](#) is shared under a [CC BY-SA 4.0](#) license and was authored, remixed, and/or curated by [Layne Morsch](#).

- [12.S: Structure Determination - Mass Spectrometry and Infrared Spectroscopy \(Summary\)](#) by Layne Morsch is licensed [CC BY-SA 4.0](#).