

14.5: The pH of Solutions of Weak Bases

The pH of a solution of a weak base can be calculated in a way which is very similar to that used for a weak acid. Instead of an acid constant K_a , a base constant K_b must be used. If a weak base B accepts protons from water according to the equation










then the base constant is defined by the expression

$$K_b = \frac{[BH^+][OH^-]}{[B]} \quad (14.5.2)$$

A list of K_b values for selected bases arranged in order of strength is given in the table below. This table is part of our larger collection of acid-base resources.

Table 14.5.1: The Base Constants for Some Bases at 25°C. Taken from Hogfelt, E. Perrin, D. D. Stability Constants of Metal Ion Complexes, 1st ed. Oxford; New Pergamon, 1979-1982. International Union of Pure and Applied Chemistry, Commission on Equilibrium. ISBN: 0080209580

Base	Formula and Ionization Equation	K_b	Molecular Shape
Ammonia	$NH_3 + H_2O \rightleftharpoons NH_4^+ + OH^-$	1.77×10^{-5}	
Aniline	$C_6H_5NH_2 + H_2O \rightleftharpoons C_6H_5NH_3^+ + OH^-$	3.9×10^{-10}	
Carbonate ion	$CO_3^{2-} + H_2O \rightleftharpoons HCO_3^- + OH^-$	2.1×10^{-4}	
Hydrazine	$N_2H_4 + H_2O \rightleftharpoons N_2H_5^+ + OH^-$ $N_2H_5^+ + H_2O \rightleftharpoons N_2H_6^{2+} + OH^-$	$K_1 = 1.2 \times 10^{-6}$ $K_2 = 1.3 \times 10^{-15}$	
Hydride ion	$H^- + H_2O \rightarrow H_2 + OH^-$	1.0	
Phosphate ion	$PO_4^{3-} + H_2O \rightleftharpoons HPO_4^{2-} + OH^-$	5.9×10^{-3}	

Base	Formula and Ionization Equation	K_b	Molecular Shape
Pyridine	$C_5H_5N + H_2O \rightleftharpoons C_5H_5NH^+ + OH^-$	1.6×10^{-9}	

To find the pH we follow the same general procedure as in the case of a weak acid. If the stoichiometric concentration of the base is indicated by c_b , the result is entirely analogous to [equation 4 in the section on the pH of weak acids](#); namely,

$$K_b = \frac{[OH^-]^2}{c_b - [OH^-]} \quad (14.5.3)$$

Under most circumstances we can make the approximation

$$c_b - [OH^-] \approx c_b$$

in which case Equation 14.5.3 reduces to the approximation

$$[OH^-] \approx \sqrt{K_b c_b} \quad (14.5.4)$$

which is identical to the expression obtained in the acid case (approximation shown in [equation 6 in the section on the pH of weak acids](#)) except that OH^- replaces H_3O^+ and b replaces a . Once we have found the hydroxide-ion concentration from this approximation, we can then easily find the pOH, and from it the pH.

✓ Example 14.5.1: pH using K_b

Using the value for K_b listed in the table, find the pH of 0.100 M NH_3 .

Solution

It is not a bad idea to guess an approximate pH before embarking on the calculation. Since we have a dilute solution of a weak base, we expect the solution to be only mildly basic. A pH of 13 or 14 would be too basic, while a pH of 8 or 9 is too close to neutral. A pH of 10 or 11 seems reasonable. Using Equation 14.5.4 we have

$$\begin{aligned} [OH^-] &= \sqrt{K_b c_b} \\ &= \sqrt{1.8 \times 10^{-5} \text{ mol L}^{-1} \times 0.100 \text{ mol L}^{-1}} \\ &= \sqrt{1.8 \times 10^{-6} \text{ mol}^2 \text{ L}^{-2}} \\ &= 1.34 \times 10^{-3} \text{ mol L}^{-1} \end{aligned}$$

Checking the accuracy of the approximation, we find

$$\frac{[OH^-]}{c_b} = \frac{1.34 \times 10^{-3}}{0.1} \approx 1 \text{ percent}$$

The approximation is valid, and we thus proceed to find the pOH.

$$pOH = -\log \frac{[OH^-]}{\text{mol L}^{-1}} = -\log(1.34 \times 10^{-3}) = 2.87$$

From which

$$pH = 14.00 - pOH = 14.00 - 2.87 = 11.13$$

This calculated value checks well with our initial guess.

Occasionally we will find that the approximation

$$c_b - [OH^-] \approx c_b$$

is not valid, in which case we must use a series of successive approximations similar to that outlined above for acids. The appropriate formula can be derived from Equation 14.5.3 and reads

$$[OH^-] \approx \sqrt{K_b(c_b - [OH^-])}$$

This page titled 14.5: The pH of Solutions of Weak Bases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Ed Vitz, John W. Moore, Justin Shorb, Xavier Prat-Resina, Tim Wendorff, & Adam Hahn.