

15.7.1: Lecture Demonstrations

Calorimeter Constant

To a styrofoam cup calorimeter containing 250 mL of water at 24.0 °C is added a known amount of heat as 250 mL of water from a second styrofoam cup at 32.0 °C. The final temperature, measured by a computer-interfaced thermistor, is 27.7 °C. The computer-generated plot of T vs. time should be projected^[1]. What is the calorimeter constant?

$$\Delta T_{\text{calorimeter}} = 27.7\text{ }^{\circ}\text{C} - 24.0\text{ }^{\circ}\text{C} = 3.7\text{ }^{\circ}\text{C}$$

$$\Delta T_{\text{cold}} = 27.5\text{ }^{\circ}\text{C} - 24.0\text{ }^{\circ}\text{C} = 3.7\text{ }^{\circ}\text{C}$$

$$\Delta T_{\text{hot}} = 27.7 - 32.0 = -4.3\text{ }^{\circ}\text{C}$$

$$q_{\text{hot}} = m \times \text{S.H.} \times \Delta T = 250\text{ g} \times 4.18\text{ J/g}^{\circ}\text{C} \times -4.3^{\circ}\text{C} = -4494\text{ J}$$

$$q_{\text{cold}} = m \times \text{S.H.} \times \Delta T = 250 \times 4.18 \times 3.7\text{ }^{\circ}\text{C} = 3867\text{ J}$$

$$q_{\text{calorimeter}} + q_{\text{hot}} + q_{\text{cold}} = 0$$

$$q_{\text{calorimeter}} - 4494 + 3867 = 0$$

$$q_{\text{calorimeter}} = 627\text{ J.}$$

$$C = Q_{\text{calorimeter}}/T = 627\text{ J} / 3.7\text{ }^{\circ}\text{C} = 169\text{ J/}^{\circ}\text{C}$$

The Ammonium Nitrate "Cold Pack"^[2]

A styrofoam cup calorimeter contains 250 mL of water at 25.0°C. Solid NH₄NO₃ (5 g) is added, and the temperature falls to 23.8°C. The computer-generated plot of T vs. time should be projected^[3]. The calorimeter is found to absorb 169 J to change its temperature 1 °C, so it is said to have a calorimeter constant of 169 J/°C. What is the enthalpy change for the dissolution reaction?

$$q_{\text{water}} = m \times \text{S.H.} \times \Delta T = 250\text{ g} \times 4.18\text{ J/g}^{\circ}\text{C} \times -1.2^{\circ}\text{C} \\ = -1254\text{ J}$$

$$q_{\text{calorim}} = C \times \Delta T = 169 \times -1.2^{\circ}\text{C} = -203\text{ J}$$

$$q_{\text{tot}} = -1254 + -203 = -1457\text{ J}$$

$$q_{\text{rxn}} = +1457\text{ J.}$$

$$\Delta H_{\text{rxn}} = q\text{ (kJ)} / n\text{ (mol)}$$

$$n = m / M = 5\text{ g NH}_4\text{NO}_3 / 80\text{ g/mol} = 0.057\text{ mol}$$

$$\Delta H_{\text{rxn}} = q/n = 1.457\text{ kJ} / 0.057\text{ mol} = +25.6\text{ kJ/mol}$$

The reaction is spontaneous even though it is endothermic, because of the large positive entropy change resulting from water association with the separate ions in solution.

References

1. We used Vernier LoggerPro(R) software
2. J. Chem. Educ., 2004, 81 (1), p 64A
3. We used Vernier LoggerPro(R) software

Contributors

- Ed Vitz (Kutztown University), [John W. Moore](#) (UW-Madison), [Justin Shorb](#) (Hope College), [Xavier Prat-Resina](#) (University of Minnesota Rochester), [Tim Wendorff](#), and [Adam Hahn](#).

This page titled [15.7.1: Lecture Demonstrations](#) is shared under a [CC BY-NC-SA 4.0](#) license and was authored, remixed, and/or curated by [Ed Vitz](#), [John W. Moore](#), [Justin Shorb](#), [Xavier Prat-Resina](#), [Tim Wendorff](#), & [Adam Hahn](#).