

TABLE OF CONTENTS

Licensing

1: Thermodynamics

- 1.1: Thermodynamic Variables and Equations of State
- 1.2: The First Law of Thermodynamics
- 1.3: Extended Explanations- The First Law of Thermodynamics
 - 1.3.1: Overview of Classical Thermodynamics
 - 1.3.2: Pressure-Volume Work
 - 1.3.3: Work and Heat are not State Functions
 - 1.3.4: Energy is a State Function
 - 1.3.5: An Adiabatic Process is a Process in which No Energy as Heat is Transferred
 - 1.3.6: The Temperature of a Gas Decreases in a Reversible Adiabatic Expansion
 - 1.3.7: Work and Heat Have a Simple Molecular Interpretation
 - 1.3.8: Pressure-Volume Work
 - 1.3.9: Heat Capacity is a Path Function
 - 1.3.10: Relative Enthalpies Can Be Determined from Heat Capacity Data and Heats of Transition
 - 1.3.11: Enthalpy Changes for Chemical Equations are Additive
 - 1.3.12: Heats of Reactions Can Be Calculated from Tabulated Heats of Formation
 - 1.3.13: The Temperature Dependence of ΔH
 - 1.3.14: Enthalpy is a State Function
 - 1.3.E: The First Law of Thermodynamics (Exercises)
- 1.4: Thermochemistry
- 1.5: The Second Law of Thermodynamics
- 1.6: The Boltzmann Distribution and the Statistical Definition of Entropy
- 1.7: The Gibbs and Helmholtz Energy
- 1.8: Extended Explanations - Helmholtz and Gibbs Energies
 - 1.8.1: Helmholtz Energy
 - 1.8.2: Gibbs Energy Determines the Direction of Spontaneity at Constant Pressure and Temperature
 - 1.8.3: The Maxwell Relations
 - 1.8.4: The Enthalpy of an Ideal Gas is Independent of Pressure
 - 1.8.5: Thermodynamic Functions have Natural Variables
 - 1.8.6: The Standard State for a Gas is an Ideal Gas at 1 Bar
 - 1.8.7: The Gibbs-Helmholtz Equation
 - 1.8.8: Fugacity Measures Nonideality of a Gas
 - 1.8.E: Helmholtz and Gibbs Energies (Exercises)
- 1.9: Phase Equilibria and Mixtures
- 1.10: Extended Explanations - Phase Equilibria
 - 1.10.1: Phase Diagrams
 - 1.10.2: A Phase Diagram Summarizes the Solid-Liquid-Gas Behavior of a Substance
 - 1.10.3: Gibbs Energies and Phase Diagrams
 - 1.10.4: The Chemical Potentials of a Pure Substance in Two Phases in Equilibrium
 - 1.10.5: The Clausius-Clapeyron Equation
 - 1.10.6: Chemical Potential Can be Evaluated From a Partition Function
 - 1.10.E: Phase Equilibria (Exercises)
- 1.11: Colligative Properties
- 1.12: Osmotic Pressure can Determine Molecular Masses

2: Extended Explanations - Solutions I - Volatile Solutes

- 2.1: A Mixture is a Combination of Two or More Substances
- 2.2: The Gibbs-Duhem Equation Relates Chemical Potential and Composition at Equilibrium
- 2.3: Chemical Potential of Each Component Has the Same Value in Each Phase in Which the Component Appears
- 2.4: Ideal Solutions obey Raoult's Law
- 2.5: Most Solutions are Not Ideal
- 2.6: Vapor Pressures of Volatile Binary Solutions
- 2.7: Activities of Nonideal Solutions
- 2.8: Activities are Calculated with Respect to Standard States
- 2.9: Gibbs Energy of Mixing of Binary Solutions in Terms of the Activity Coefficient
- 2.E: Solutions I- Liquid-Liquid Solutions (Exercises)

3: Extended Explanations - Solutions II - Nonvolatile Solutes

- 3.1: Standard State of Nonvolatile Solutions
- 3.2: The Activities of Nonvolatile Solutes
- 3.3: Electrolytes Solutions are Nonideal at Low Concentrations
- 3.4: The Debye-Hückel Theory
- 3.5: Extending Debye-Hückel Theory to Higher Concentrations
- 3.6: Homework Problems

[Index](#)

[Glossary](#)

[Detailed Licensing](#)

[Detailed Licensing](#)