

1.4: Phase Space

We construct a Cartesian space in which each of the $6N$ coordinates and momenta is assigned to one of $6N$ mutually orthogonal axes. Phase space is, therefore, a $6N$ dimensional space. A point in this space is specified by giving a particular set of values for the $6N$ coordinates and momenta. Denote such a point by

$$x = (p_1, \dots, p_N, r_1, \dots, r_N)$$

x is a $6N$ dimensional vector. Thus, the time evolution or trajectory of a system as specified by Hamilton's equations of motion, can be expressed by giving the phase space vector, x as a function of time.




Figure 1.4.1: Evolution of an ensemble of classical systems in phase space (top). The systems are a massive particle in a one-dimensional potential well (red curve, lower figure). The initially compact ensemble becomes swirled up over time.

The law of conservation of energy, expressed as a condition on the phase space vector:

$$H(x(t)) = \text{const} = E$$

defines a $6N - 1$ dimensional hypersurface in phase space on which the trajectory must remain.

This page titled 1.4: Phase Space is shared under a [CC BY-NC-SA 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/) license and was authored, remixed, and/or curated by [Mark Tuckerman](https://chem.libretexts.org/@go/page/5100).