


## 1.30: From Coordinate Space to Momentum Space and Back

The 2s state of the one-dimensional hydrogen atom is used to illustrate transformations back and forth between the coordinate and momentum representations.


$$\Psi_2(x) := \frac{1}{\sqrt{8}} \cdot x \cdot (2 - x) \cdot \exp\left(-\frac{x}{2}\right)$$



The 2s state is Fourier transformed into momentum space (using atomic units) and the magnitude of the momentum wave function is displayed.

$$\langle p | \Psi_2 \rangle = \int_0^\infty \langle p | x \rangle \langle x | \Psi_2 \rangle dx \quad \text{where} \quad \langle p | x \rangle = \frac{1}{\sqrt{2\pi}} \exp\left(\frac{-ipx}{\hbar}\right)$$

$$\Psi_2(p) := \frac{1}{\sqrt{2 \cdot \pi}} \int_0^\infty \exp(-i \cdot p \cdot x) \cdot \Psi_2(x) dx \quad \text{simplify} \rightarrow \frac{2}{\pi^{\frac{1}{2}}} \cdot \frac{2 \cdot i \cdot p - 1}{(2 \cdot i \cdot p + 1)^3}$$




The return to coordinate space is carried out in the numeric mode, integrating over the range of momentum values shown above ( $\pm 10$  is effectively  $\pm\infty$ ).

$$\langle x | \Psi_2 \rangle = \int_{-\infty}^{\infty} \langle x | p \rangle \langle p | \Psi_2 \rangle dp \quad \text{where} \quad \langle x | p \rangle = \frac{1}{\sqrt{2\pi}} \exp\left(\frac{ipx}{\hbar}\right)$$

$$\Psi_2(x) := \int_{-10}^{10} \frac{1}{\sqrt{2 \cdot \pi}} \cdot \exp(i \cdot p \cdot x) \cdot \Psi_2(p) dp$$

The graphical display below shows that we have successfully returned to coordinate space.



This page titled [1.30: From Coordinate Space to Momentum Space and Back](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Frank Rioux](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.