

6.6: BCl₃

C_{3v} Symmetry - BCl₃

The infrared spectrum of BCl₃ shows vibrational bands at 995, 480, and 244 cm⁻¹, while Raman bands appear at 995, 471, and 244 cm⁻¹. Is the geometry of the molecule trigonal pyramid (C_{3v}) or trigonal planar (D_{3h})? Is your answer to this question consistent with chemical bonding principles (VSEPR)? Assign symmetry labels to the vibrational bands and identify the stretches and bends.

$$\begin{aligned}
 C_{3v} = & \begin{pmatrix} C_3 & \sigma_v \\ 1 & 1 & 1 \\ 1 & 1 & -1 \\ 2 & -1 & 0 \end{pmatrix} \begin{matrix} 1 : z, x^2 + y^2, z^2 \\ A_2 : R_z \\ E : (x, y)(R_x, R_y), (x^2 + y^2, xy)(xz, yz) \end{matrix} & C_{3v} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} & \Gamma_{uma} = \begin{pmatrix} 4 \\ 1 \\ 2 \end{pmatrix} & \Gamma_{bonds} = \begin{pmatrix} 3 \\ 0 \\ 1 \end{pmatrix} \\
 A_1 = (C_{3v}^T)^{<1>} & A_2 = (C_{3v}^T)^{<2>} & E = (C_{3v}^T)^{<3>} & h = \sum C_{3v} \\
 \Gamma_{tot} = \overline{[\Gamma_{uma}(A_1 + E)]} & \Gamma_{tot}^T = (12 \ 0 \ 2) & \Gamma_{vib} = \Gamma_{tot} - A_1 - A_2 - 2E \\
 i = 1..3 & \text{Vib}_i = \frac{\sum [C_{3v}(C_{3v}^T)^{<i>} \Gamma_{vib}]}{h} & \text{Vib} = \begin{pmatrix} 2 \\ 0 \\ 2 \end{pmatrix} \begin{matrix} 1 : z, x^2 + y^2, z^2 \\ A_2 : R_z \\ E : (x, y)(R_x, R_y), (x^2 + y^2, xy)(xz, yz) \end{matrix} \\
 \Gamma_{stretch} = \Gamma_{bonds} & \text{Stretch}_i = \frac{\sum [C_{3v}(C_{3v}^T)^{<i>} \Gamma_{stretch}]}{h} & \text{Stretch} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \begin{matrix} 1 : z, x^2 + y^2, z^2 \\ A_2 : R_z \\ E : (x, y)(R_x, R_y), (x^2 + y^2, xy)(xz, yz) \end{matrix} \\
 \Gamma_{bend} = \Gamma_{vib} - \Gamma_{stretch} & \text{Bend}_i = \frac{\sum [C_{3v}(C_{3v}^T)^{<i>} \Gamma_{bend}]}{h} & \text{Bend} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \begin{matrix} 1 : z, x^2 + y^2, z^2 \\ A_2 : R_z \\ E : (x, y)(R_x, R_y), (x^2 + y^2, xy)(xz, yz) \end{matrix}
 \end{aligned}$$

This analysis predicts that there should be 4 IR and 4 Raman active vibrations. It also predicts that the IR and Raman vibrations should be coincident. This is not in agreement with the experimental data, so BCl₃ does not have trigonal pyramidal geometry.

This analysis also predicts that the stretches should have A₁ and E symmetry, but since it has already been concluded that the molecule does not have C_{3v} symmetry this result will not be discussed further.

D_{3h} Symmetry - BCl₃

$$\begin{aligned}
 C_{3v} = & \begin{pmatrix} C_3 & C_2 & \sigma_h & S_3 & \sigma_v \\ 1 & 1 & 11 & 1 & 1 \\ 1 & 1 & -1 & 1 & 1 \\ 2 & -1 & 0 & 2 & -1 \\ 1 & 1 & 1 & -1 & -1 \\ 1 & 1 & -1 & -1 & 1 \\ 2 & -1 & 0 & -2 & 1 \end{pmatrix} \begin{matrix} 1' : x^2 + y^2, z^2 \\ A_2' : R_z \\ E' : (x, y), (x^2 + y^2, xy) \\ A_1'' : \\ A_2'' : z \\ E'' : (R_x, R_y), (xz, yz) \end{matrix} & D_{3h} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 1 \\ 2 \\ 3 \end{bmatrix} & \Gamma_{uma} = \begin{bmatrix} 4 \\ 1 \\ 2 \\ 4 \\ 1 \\ 2 \end{bmatrix} & \Gamma_{bonds} = \begin{bmatrix} 3 \\ 0 \\ 1 \\ 3 \\ 0 \\ 1 \end{bmatrix} \\
 A_1 = (C_{D_{3h}}^T)^{<1>} & A_2 = (C_{D_{3h}}^T)^{<2>} & E = (C_{D_{3h}}^T)^{<3>} & A_{11} = (C_{D_{3h}}^T)^{<4>} \\
 A_{21} = (C_{D_{3h}}^T)^{<5>} & E_1 = (C_{D_{3h}}^T)^{<6>} & h = \sum D_{3h} & \Gamma_{tot} = \overline{[\Gamma_{uma}(A_{21} + E)]} \\
 \Gamma_{tot}^T = (12 \ 0 \ -2 \ 4 \ -2 \ 2) & \Gamma_{vib} = \Gamma_{tot} - A_2 - E - A_{21} - E_1 & i = 1..6 \\
 \Gamma_{stretch} = \Gamma_{bonds} & \Gamma_{bend} = \Gamma_{vib} - \Gamma_{stretch}
 \end{aligned}$$

$$\begin{aligned}
 \text{Vib}_i &= \frac{\sum_h [D3h(C_{D3h}^T)^{<i>}\Gamma_{vib}]}{h} & \text{Stretch}_i &= \frac{\sum_h [D3h(C_{D3h}^T)^{<i>}\Gamma_{stretch}]}{h} & \text{Bend}_i &= \frac{\sum_h [D3h(C_{D3h}^T)^{<i>}\Gamma_{bend}]}{h} \\
 \text{Vib} &= \begin{bmatrix} 1 \\ 0 \\ 2 \\ 0 \\ 1 \\ 0 \end{bmatrix} \begin{matrix} 1' : x^2 + y^2, z^2 \\ A'_2 : Rz \\ E' : (x, y), (x^2 + y^2, xy) \\ A1'' : \\ A2'' : z \\ E'' : (Rx, Ry), (xz, yz) \end{matrix} & \text{Stretch} &= \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \begin{matrix} 1' : x^2 + y^2, z^2 \\ A'_2 : Rz \\ E' : (x, y), (x^2 + y^2, xy) \\ A1'' : \\ A2'' : z \\ E'' : (Rx, Ry), (xz, yz) \end{matrix} \\
 \text{Bend} &= \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 1 \\ 0 \end{bmatrix} \begin{matrix} 1' : x^2 + y^2, z^2 \\ A'_2 : Rz \\ E' : (x, y), (x^2 + y^2, xy) \\ A1'' : \\ A2'' : z \\ E'' : (Rx, Ry), (xz, yz) \end{matrix}
 \end{aligned}$$

This analysis shows that there are 3 IR active modes (2E', A2'') and 3 Raman active modes (A1', E'). It also show that there are two coincidences (2E'). Thus, D_{3h} symmetry for BCl₃ is consistent with the spectroscopic data.

Thus the specific assignments are: A1' = 471; A2'' = 480; E' = 995; E' = 244. From the analysis above it can be seen that the stretches occur at 995 and 471, and the bends are at 480 and 244 cm⁻¹.

VSEPR theory would predict a trigonal planar arrangement of chlorine atoms around the electron deficient boron, so the symmetry analysis and the bonding theory are in agreement with each other and the spectroscopic measurements.

This page titled [6.6: BCl₃](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Frank Rioux](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.