


5.18: Another Look at the Double-Slit Experiment

Illumination of a double-slit screen with a coherent particle beam leads to a Schrödinger "cat state" that can be represented by a linear superposition (unnormalized) of two Gaussian wavepackets. The probability distribution function in coordinate space, $|\Psi(x)|^2$, at the slit-screen for this "cat state" is shown below.

$$x = -10, -9.99 \dots 10 \quad \Psi(x) = \exp[-(x-5)^2] + \exp[-(x+5)^2] \quad P(x) = (|\Psi(x)|)^2$$



The slits localize the particle in the x -direction which leads to a spread in the x -component of the momentum required by the uncertainty principle, $\Delta x \Delta p_x > \hbar/4$. The momentum wave function is obtained by a Fourier transform of the coordinate-space wave function.

$$\Phi(p_x) = \frac{1}{\sqrt{2\pi}} \left[\int_{-\infty}^{\infty} \exp(-ip_x x) [\exp[-(x-5)^2] + \exp[-(x+5)^2]] dx \right]$$


Evaluation of the integral yields,

$$\Phi(p_x) = \frac{1}{\sqrt{2}} \left[\exp\left[\frac{-1}{4}p_x(p_x + 20i)\right] + \exp\left[\frac{-1}{4}p_x(p_x - 20i)\right] \right]$$

The momentum probability function in the x -direction is $|\Phi(p_x)|^2$ and simplifies to the expression given below when evaluated.

$$P(p_x) = 2 \exp\left(\frac{-1}{2}p_x^2\right) \cos(5p_x)^2$$

This momentum probability function is displayed below.



Because the arrival at position x on the detection screen is proportional to p_x it is also proportional to $|\Phi(p_x)|^2$. In other words, the particle distribution at the detection screen is determined by the momentum distribution at the slit screen. This means the position measurement at the detection screen is effectively a measurement of the p_x . Therefore, the particle distribution at the detector screen will have the same shape as shown in the figure above.

In summary, the double-slit experiment clearly reveals the three essential steps in a quantum mechanical experiment:

1. State preparation (interaction of the incident beam with the slit-screen)
2. Measurement of an observable (arrival of scattered beam at the detection screen)
3. Calculation of expected results of the measurement step

*The preparation of this tutorial was stimulated by reading "Quantum interference with slits" by Thomas Marcella which appeared in *European Journal of Physics* **23**, 615-621 (2002). This paper offers a lucid and novel quantum mechanical analysis of a very important experiment.

Additional references:

R. P. Feynman, R. B. Leighton, and M. Sands, *The Feynman Lectures on Physics*, Volume 3; Addison-Wesley; Reading, 1965, Chapters 1 and 3.

R. P. Feynman, *The Character of Physical Law*; MIT Press: Cambridge, 1967, Chapter 6.

A. Tonomura, J. Endo, T. Matsuda, T. Kawasaki, and H. Exawa, "Demonstration of single-electron buildup of an interference pattern" *Am. J. Phys.* **57**, 117-120 (1989).

D. Leibfried, T. Pfau, and C. Monroe, "Shadows and Mirrors: Reconstructing Quantum States of Atom Motion" *Phys. Today* **51(4)**, 22-28 (1998).

The double-slit experiment with single electrons was recently selected (informally) as physics most beautiful experiment. The following web reference traces the history of double-slit interference experiments from the time of Thomas Young to the present, presenting numerous literature references in the process: <http://physicsweb.org/article/world/15/9/1>.

This page titled [5.18: Another Look at the Double-Slit Experiment](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [Frank Rioux](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.