

18.2: The Single-Determinant Wavefunction

The simplest trial function of the form given above is the single Slater determinant function:

$$|\Psi\rangle = |\phi_1\phi_2\phi_3\dots\phi_N\rangle.$$

For such a function, the CI part of the energy minimization is absent (the classic papers in which the SCF equations for closed- and open-shell systems are treated are C. C. J. Roothaan, Rev. Mod. Phys. 23 , 69 (1951); 32 , 179 (1960)) and the density matrices simplify greatly because only one spin-orbital occupancy is operative. In this case, the orbital optimization conditions reduce to:

$$\hat{F}\phi_i = \sum_j \epsilon_{i,j}\phi_j,$$

where the so-called Fock operator \hat{F} is given by

$$\hat{F}\phi_i = h\phi_i + \sum_{j(\text{occupied})} [\hat{J}_j - \hat{K}_j] \phi_i.$$

The coulomb (\hat{J}_j) and exchange (\hat{K}_j) operators are defined by the relations:

$$\hat{J}_j\phi_i = \int \phi_j^*(r')\phi_j(r') \frac{1}{|r-r'|} d\tau' \phi_i(r)$$

and

$$\hat{K}_j\phi_i = \int \phi_j^*(r')\phi_i(r') \frac{1}{|r-r'|} d\tau' \phi_j(r).$$

Again, the integration implies integration over the spin variables associated with the ϕ_j (and, for the exchange operator, ϕ_i), as a result of which the exchange integral vanishes unless the spin function of ϕ_j is the same as that of ϕ_i ; the coulomb integral is non-vanishing no matter what the spin functions of ϕ_j and ϕ_i .

The sum over coulomb and exchange interactions in the Fock operator runs only over those spin-orbitals that are occupied in the trial Ψ . Because a unitary transformation among the orbitals that appear in $|\Psi\rangle$ leaves the determinant unchanged (this is a property of determinants- $\det(UA) = \det(U)\det(A) = 1\det(A)$, if U is a unitary matrix), it is possible to choose such a unitary transformation to make the $\epsilon_{i,j}$ matrix diagonal. Upon so doing, one is left with the so-called **canonical Hartree-Fock equations**:

$$\hat{F}\phi_i = \epsilon_i\phi_j,$$

where ϵ_i is the diagonal value of the $\epsilon_{i,j}$ matrix after the unitary transformation has been applied; that is, ϵ_i is an eigenvalue of the $\epsilon_{i,j}$ matrix. These equations are of the eigenvalue-eigenfunction form with the Fock operator playing the role of an effective one-electron Hamiltonian and the ϕ_i playing the role of the one-electron eigenfunctions.

It should be noted that the Hartree-Fock equations $\hat{F}\phi_i = \epsilon_i\phi_j$ possess solutions for the spin-orbitals which appear in Ψ (the so-called **occupied** spin-orbitals) as well as for orbitals which are not occupied in Ψ (the so-called **virtual** spin-orbitals). In fact, the F operator is hermitian, so it possesses a complete set of orthonormal eigenfunctions; only those which appear in Ψ appear in the coulomb and exchange potentials of the Fock operator. The physical meaning of the occupied and virtual orbitals will be clarified later in this Chapter (Section VII.A).

This page titled [18.2: The Single-Determinant Wavefunction](#) is shared under a [CC BY-NC-SA 4.0](#) license and was authored, remixed, and/or curated by [Jack Simons](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.