

5.6: Entropy and Disorder

A common interpretation of entropy is that it is somehow a measure of chaos or randomness. There is some utility in that concept. Given that entropy is a measure of the dispersal of energy in a system, the more chaotic a system is, the greater the dispersal of energy will be, and thus the greater the entropy will be. Ludwig Boltzmann (1844 – 1906) (O'Connor & Robertson, 1998) understood this concept well, and used it to derive a statistical approach to calculating entropy. Boltzmann proposed a method for calculating the entropy of a system based on the number of energetically equivalent ways a system can be constructed.

Boltzmann proposed an expression, which in its modern form is:

$$S = k_b \ln(W) \quad (5.6.1)$$

This rather famous equation is etched on Boltzmann's grave marker in commemoration of his profound contributions to the science of thermodynamics (Figure 5.6.1).



Figure 5.6.1: Ludwig Boltzmann (1844 - 1906)

✓ Example 5.6.1:

Calculate the entropy of a carbon monoxide crystal, containing 1.00 mol of CO, and assuming that the molecules are randomly oriented in one of two equivalent orientations.

Solution

Using the Boltzmann formula (Equation 5.6.1):

$$S = nK \ln(W)$$

And using $W = 2$, the calculation is straightforward.

$$\begin{aligned} S &= \left(1.00 \text{ mol} \cot \frac{6.022 \times 10^{23}}{1 \text{ mol}} \right) (1.38 \times 10^{-23} \text{ J/K}) \ln 2 \\ &= 5.76 \text{ J/K} \end{aligned}$$

This page titled 5.6: Entropy and Disorder is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Patrick Fleming.