


## 9.8: STEREOCHEMISTRY OF REACTIONS - HYDRATION OF CHIRAL ALKENES

### Learning Objective

- discern the stereochemical differences between the EAR of chiral and achiral alkenes

### STEREOCHEMISTRY - THE SUBTLE DETAILS

In the previous section, the addition of water to the achiral alkene produced a racemic mixture of two enantiomeric alcohols. They are produced in equal amounts so the mixture is optically inactive. What would occur if we carried out a similar reaction on a chiral alkene? Consider (S)-3-methyl-1-pentene reacting with water (acid catalyzed). Proton addition produces a carbocation intermediate that is chiral (\* denotes stereogenic centre). That intermediate does not have a plane of symmetry and therefore attack by water is not equal from the top and bottom. This ultimately produces R and S products in a non 50:50 ratio as shown in the mechanism below.




### Exercise

- Predict the products of the following reaction showing stereochemistry.



### Answer

- The products are diastereomers of one another.



### CONTRIBUTORS AND ATTRIBUTIONS

- Dr. Dietmar Kennepohl FCIC (Professor of Chemistry, [Athabasca University](#))
- Prof. Steven Farmer ([Sonoma State University](#))

9.8: Stereochemistry of Reactions - Hydration of Chiral Alkenes is shared under a [CC BY-NC-SA 4.0](#) license and was authored, remixed, and/or curated by LibreTexts.