


## 12.14: MORE NMR EXAMPLES

### ADDITIONAL NMR EXAMPLES


For each molecule, predict the number of signals in the  $^1\text{H}$ -NMR and the  $^{13}\text{C}$ -NMR spectra (do not count split peaks - eg. a quartet counts as only one signal). Assume that diastereotopic groups are non-equivalent.



**P5.2:** For each of the 20 common amino acids, predict the number of signals in the proton-decoupled  $^{13}\text{C}$ -NMR spectrum.

**P5.3:** Calculate the chemical shift value (expressed in Hz, to one decimal place) of each sub-peak on the  $^1\text{H}$ -NMR doublet signal below. Do this for:

- a spectrum obtained on a 300 MHz instrument
- a spectrum obtained on a 100 MHz instrument




**P5.4:** Consider a quartet signal in an  $^1\text{H}$ -NMR spectrum obtained on a 300 MHz instrument. The chemical shift is recorded as 1.7562 ppm, and the coupling constant is  $J = 7.6$  Hz. What is the chemical shift, expressed to the nearest 0.1 Hz, of the furthest downfield sub-peak in the quartet? What is the resonance frequency (again expressed in Hz) of this sub-peak?)

**P5.5:** One easily recognizable splitting pattern for the aromatic proton signals from disubstituted benzene structures is a pair of doublets. Does this pattern indicate *ortho*, *meta*, or *para* substitution?

**P5.6 :**Match spectra below to their corresponding structures A-F.

Structures:



#### Spectrum 1

$\delta$	splitting integration	
4.13	q	2
2.45	t	2
1.94	quintet	1
1.27	t	3

#### Spectrum 2

$\delta$	splitting integration	
3.68	s	3
2.99	t	2
1.95	quintet	1

#### Spectrum 3

$\delta$	splitting integration	
4.14	q	1
2.62	s	1
1.26	t	1.5

#### Spectrum 4

$\delta$	splitting integration	
4.14	q	4
3.22	s	1
1.27	t	6
1.13	s	9

#### Spectrum 5


$\delta$	splitting integration	
4.18	q	1
1.92	q	1
1.23	t	1.5
0.81	t	1.5

#### Spectrum 6


$\delta$	splitting integration	
3.69	s	1.5
2.63	s	1

**P5.7:** Match spectra 7-12 below to their corresponding structures G-L .


Structures:




**G**




**H**




**I**



**J**



**K**



**L**

Spectrum 7:

$\delta$	splitting integration	
9.96	d	1
5.88	d	1
2.17	s	3
1.98	s	3

Spectrum 8:

$\delta$	splitting integration	
9.36	s	1
6.55	q	1
2.26	q	2
1.99	d	3
0.96	t	3

Spectrum 9:

$\delta$	splitting integration	
9.57	s	1
6.30	s	1
6.00	s	1
1.84	s	3

Spectrum 10:

$\delta$	splitting integration	
9.83	t	1
2.27	d	2
1.07	s	9

Spectrum 11:






$\delta$	splitting integration	
9.75	t	1
2.30	dd	2
2.21	m	1
0.98	d	6

Spectrum 12:

$\delta$	splitting integration	
8.08	s	1
4.13	t	2
1.70	m	2
0.96	t	3

**P5.8:** Match the  $^1\text{H}$ -NMR spectra 13-18 below to their corresponding structures M-R .

Structures:



Spectrum 13:

$\delta$	splitting integration	
8.15	d	1
6.33	d	1

Spectrum 14: 1-723C (structure O)

$\delta$	splitting integration	
6.05	s	1
2.24	s	3

Spectrum 15:

$\delta$	splitting integration	
8.57	s (b)	1
7.89	d	1
6.30	d	1
2.28	s	3

Spectrum 16:

$\delta$	splitting integration	
9.05	s (b)	1
8.03	s	1
6.34	s	1
5.68	s (b)	1
4.31	s	2

Spectrum 17:


$\delta$	splitting	integration
7.76	d	1
7.57	s (b)	1
6.44	d	1
2.78	q	2
1.25	t	3

#### Spectrum 18:

$\delta$	splitting	integration
4.03	s	1
2.51	t	1
2.02	t	1

**P5.9:** Match the  $^1\text{H}$ -NMR spectra 19-24 below to their corresponding structures S-X.

Structures:



#### Spectrum 19:

$\delta$	splitting	integration
9.94	s	1
7.77	d	2
7.31	d	2
2.43	s	3

#### Spectrum 20:

$\delta$	splitting	integration
10.14	s	2
8.38	s	1
8.17	d	2
7.75	t	1

#### Spectrum 21:

$\delta$	splitting	integration
9.98	s	1
7.81	d	2
7.50	d	2

#### Spectrum 22:

$\delta$	splitting integration	
7.15-7.29	m	2.5
2.86	t	1
2.73	t	1
2.12	s	1.5

Spectrum 23:







$\delta$	splitting integration	
7.10	d	1
6.86	d	1
3.78	s	1.5
3.61	s	1
2.12	s	1.5

Spectrum 24:

$\delta$	splitting integration	
7.23-7.30	m	1
3.53	s	1

**P5.10:** Match the  $^1\text{H}$ -NMR spectra 25-30 below to their corresponding structures AA-FF.

Structures:



Spectrum 25:

$\delta$	splitting integration	
9.96	s	1
7.79	d	2
7.33	d	2
2.72	q	2
1.24	t	3

Spectrum 26:

$\delta$	splitting integration	
9.73	s	1
7.71	d	2
6.68	d	2
3.06	s	6

Spectrum 27:

$\delta$	splitting integration	
7.20-7.35	m	10
5.12	s	1
2.22	s	3

Spectrum 28:

$\delta$	splitting integration	
8.08	s	1
7.29	d	2
6.87	d	2
5.11	s	2
3.78	s	3

Spectrum 29:


$\delta$	splitting integration	
7.18	d	1
6.65	m	1.5
3.2	q	2
1.13	t	3

Spectrum 30:


$\delta$	splitting integration	
8.32	s	1
4.19	t	2
2.83	t	2
2.40	s	3

**P5.11:** Match the  $^1\text{H}$ -NMR spectra 31-36 below to their corresponding structures GG-LL


Structures:




**GG**




**HH**




**II**



**JJ**



**KK**



**LL**

Spectrum 31:

$\delta$	splitting integration	
6.98	d	1
6.64	d	1
6.54	s	1
4.95	s	1
2.23	s	3
2.17	s	3

Spectrum 32:

$\delta$	splitting integration	
7.08	d	1
6.72	d	1
6.53	s	1
4.81	s	1
3.15	7-tet	1
2.24	s	3
1.22	d	6

Spectrum 33:

$\delta$	splitting integration	
7.08	d	2
6.71	d	2
6.54	s	1
3.69	s	3
3.54	s	2

Spectrum 34:

$\delta$	splitting integration	
9.63	s	1
7.45	d	2
6.77	d	2
3.95	q	2
2.05	s	3
1.33	t	3

Spectrum 35:

$\delta$	splitting integration	
9.49	s	1
7.20	d	2
6.49	d	2
4.82	s	2
1.963	s	3

Spectrum 36:



$\delta$	splitting	integration
9.58	s(b)	1
9.31	s	1
7.36	d	1
6.67	s	1
6.55	d	1
2.21	s	3
2.11	s	3

**P5.12:** Use the NMR data given to deduce structures.

a ) Molecular formula:  $C_5H_8O$

$^1H$ -NMR:

$\delta$	splitting	integration
9.56	s	1
	d	
6.25	(J~1 Hz)	1
	d	
5.99	(J~1 Hz)	1
	d	
2.27	q	2
1.18	t	3

$^{13}C$ -NMR

$\delta$	DEPT
194.60	CH
151.77	C
132.99	CH <sub>2</sub>
20.91	CH <sub>2</sub>
11.92	CH <sub>3</sub>

b) Molecular formula:  $C_7H_{14}O_2$

$^1H$ -NMR:

$\delta$	splitting	integration
3.85	d	2
2.32	q	2
1.93	m	1
1.14	t	3
0.94	d	6

$^{13}C$ -NMR

$\delta$	DEPT
174.47	C
70.41	CH <sub>2</sub>
27.77	CH
27.64	CH <sub>2</sub>
19.09	CH <sub>3</sub>
9.21	CH <sub>3</sub>

c) Molecular formula:  $C_5H_{12}O$

$^1H$ -NMR:

$\delta$	splitting integration	
3.38	s	2H
2.17	s	1H
0.91	s	9H

$^{13}C$ -NMR

$\delta$	DEPT
73.35	$CH_2$
32.61	C
26.04	$CH_3$

d) Molecular formula:  $C_{10}H_{12}O$

$^1H$ -NMR:

$\delta$	splitting integration	
7.18-7.35	m	2.5
3.66	s	1
2.44	q	1
1.01	t	1.5


$^{13}C$ -NMR

$\delta$	DEPT
208.79	C
134.43	C
129.31	CH
128.61	CH
126.86	CH
49.77	$CH_2$
35.16	$CH_2$
7.75	$CH_3$

P5.13:


$^{13}C$ -NMR data is given for the molecules shown below. Complete the peak assignment column of each NMR data table.

a)




$\delta$	DEPT	carbon #
161.12	CH	
65.54	$CH_2$	
21.98	$CH_2$	
10.31	$CH_3$	

b)




$\delta$	DEPT carbon #
194.72	C
149.10	C
146.33	CH
16.93	CH <sub>2</sub>
14.47	CH <sub>3</sub>
12.93	CH <sub>3</sub>

c)




$\delta$	DEPT carbon #
171.76	C
60.87	CH <sub>2</sub>
58.36	C
24.66	CH <sub>2</sub>
14.14	CH <sub>3</sub>
8.35	CH <sub>3</sub>

d)



$\delta$	DEPT carbon #
173.45	C
155.01	C
130.34	CH
125.34	C
115.56	CH
52.27	CH <sub>3</sub>
40.27	CH <sub>2</sub>


e)




$\delta$	DEPT carbon #
147.79	C
129.18	CH
115.36	CH
111.89	CH
44.29	CH <sub>2</sub>
12.57	CH <sub>3</sub>

**P5.14:** You obtain the following data for an unknown sample. Deduce its structure.

<sup>1</sup>H-NMR:




<sup>13</sup>C-NMR:




Mass Spectrometry:

$\delta$	splitting	integration
4.3	septet	0.0735
3.4	triplet	0.661
1.9	sextet	0.665
1.7	doublet	0.441
1.0	triplet	1.00

**C5.2:** Researchers wanted to investigate a reaction which can be catalyzed by the enzyme alcohol dehydrogenase in yeast. They treated 4'-acetylpyridine (1) with living yeast, and isolated the alcohol product(s) (some combination of 2A and 2B).



- a) Will the products 2A and 2B have identical or different  $^1\text{H-NMR}$  spectra? Explain.
- b) Suggest a  $^1\text{H-NMR}$  experiment that could be used to determine what percent of starting material (1) got turned into product (2A and 2B).
- c) With purified 2A/2B, the researchers carried out the subsequent reaction shown below to make 3A and 3B, known as 'Mosher's esters'. Do 3A and 3B have identical or different  $^1\text{H-NMR}$  spectra? Explain.



d) Explain, very specifically, how the researchers could use  $^1\text{H-NMR}$  to determine the relative amounts of 2A and 2B formed in the reaction catalyzed by yeast enzyme.

12.14: More NMR Examples is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.