

3.3: Vector Addition and Subtraction- Analytical Methods

Learning Objectives

By the end of this section, you will be able to:

- Understand the rules of vector addition and subtraction using analytical methods.
- Apply analytical methods to determine vertical and horizontal component vectors.
- Apply analytical methods to determine the magnitude and direction of a resultant vector.

Analytical methods of vector addition and subtraction employ geometry and simple trigonometry rather than the ruler and protractor of graphical methods. Part of the graphical technique is retained, because vectors are still represented by arrows for easy visualization. However, analytical methods are more concise, accurate, and precise than graphical methods, which are limited by the accuracy with which a drawing can be made. Analytical methods are limited only by the accuracy and precision with which physical quantities are known.

Resolving a Vector into Perpendicular Components

Analytical techniques and right triangles go hand-in-hand in physics because (among other things) motions along perpendicular directions are independent. We very often need to separate a vector into perpendicular components. For example, given a vector like A in Figure 3.3.1, we may wish to find which two perpendicular vectors, A_x and A_y , add to produce it.




Figure 3.3.1: The vector A , with its tail at the origin of an x , y -coordinate system, is shown together with its x - and y -components, A_x and A_y . These vectors form a right triangle. The analytical relationships among these vectors are summarized below.

A_x and A_y are defined to be the components of A along the x - and y -axes. The three vectors A , A_x , and A_y form a right triangle:

$$A_x + A_y = A. \quad (3.3.1)$$

Note that this relationship between vector components and the resultant vector holds only for vector quantities (which include both magnitude and direction). The relationship does not apply for the magnitudes alone. For example, if $A_x = 3m$ east, $A_y = 4m$ north, and $A = 5m$ north-east, then it is true that the vectors $A_x + A_y = A$. However, it is not true that the sum of the magnitudes of the vectors is also equal. That is,

$$3m + 4m \neq 5m \quad (3.3.2)$$

Thus,

$$A_x + A_y \neq A \quad (3.3.3)$$

If the vector A is known, then its magnitude A (its length) and its angle θ (its direction) are known. To find A_x and A_y , its x - and y -components, we use the following relationships for a right triangle.

$$A_x = A \cos \theta \quad (3.3.4)$$

and

$$A_y = A \sin \theta. \quad (3.3.5)$$




Figure 3.3.2: The magnitudes of the vector components A_x and A_y can be related to the resultant vector A and the angle θ with trigonometric identities. Here we see that $A_x = A \cos \theta$ and $A_y = A \sin \theta$.

Suppose, for example, that A is the vector representing the total displacement of the person walking in a city considered in Kinematics in Two Dimensions: An Introduction and Vector Addition and Subtraction: Graphical Methods.




Figure 3.3.3: We can use the relationships $A_x = A \cos \theta$ and $A_y = A \sin \theta$ to determine the magnitude of the horizontal and vertical component vectors in this example.

Then $A = 10.3$ blocks and $\theta = 29.1^\circ$, so that

$$A_x = A \cos \theta = (10.3 \text{ blocks})(\cos 29.1^\circ) = 9.0 \text{ blocks} \quad (3.3.6)$$

$$A_y = A \sin \theta = (10.3 \text{ blocks})(\sin 29.1^\circ) = 5.0 \text{ blocks} \quad (3.3.7)$$

Calculating a Resultant Vector

If the perpendicular components A_x and A_y of a vector A are known, then A can also be found analytically. To find the magnitude A and direction θ of a vector from its perpendicular components A_x and A_y , we use the following relationships:

$$A = \sqrt{A_x^2 + A_y^2} \quad (3.3.8)$$

$$\theta = \tan^{-1}(A_y/A_x) \quad (3.3.9)$$




Figure 3.3.4: The magnitude and direction of the resultant vector can be determined once the horizontal and vertical components A_x and A_y have been determined.

Note that the equation $A = \sqrt{A_x^2 + A_y^2}$ is just the Pythagorean theorem relating the legs of a right triangle to the length of the hypotenuse. For example, if A_x and A_y are 9 and 5 blocks, respectively, then $A = \sqrt{9^2 + 5^2} = 10.3$ blocks, again consistent with the example of the person walking in a city. Finally, the direction is $\theta = \tan^{-1}(5/9) = 29.1^\circ$, as before.

DETERMINING VECTORS AND VECTOR COMPONENTS WITH ANALYTICAL METHODS

Equations $A_x = A \cos \theta$ and $A_y = A \sin \theta$ are used to find the perpendicular components of a vector—that is, to go from A and θ to A_x and A_y . Equations $A = \sqrt{A_x^2 + A_y^2}$ and $\theta = \tan^{-1}(A_y / A_x)$ are used to find a vector from its perpendicular components—that is, to go from A_x and A_y to A and θ . Both processes are crucial to analytical methods of vector addition and subtraction.

Adding Vectors Using Analytical Methods

To see how to add vectors using perpendicular components, consider Figure 3.3.5, in which the vectors A and B are added to produce the resultant R .




Figure 3.3.5: Vectors A and B are two legs of a walk, and R is the resultant or total displacement. You can use analytical methods to determine the magnitude and direction of R .

If A and B represent two legs of a walk (two displacements), then R is the total displacement. The person taking the walk ends up at the tip of R . There are many ways to arrive at the same point. In particular, the person could have walked first in the x -direction and then in the y -direction. Those paths are the x - and y -components of the resultant, R_x and R_y . If we know R_x and R_y , we can find R and θ using the equations $A = \sqrt{A_x^2 + A_y^2}$ and $\theta = \tan^{-1}(A_y / A_x)$. When you use the analytical method of vector addition, you can determine the components or the magnitude and direction of a vector.

Step 1. Identify the x - and y -axes that will be used in the problem. Then, find the components of each vector to be added along the chosen perpendicular axes. Use the equations $A_x = A \cos \theta$ and $A_y = A \sin \theta$ to find the components. In Figure, these components are A_x , A_y , B_x , and B_y . The angles that vectors A and B make with the x -axis are θ_A and θ_B , respectively.




Figure 3.3.6: To add vectors A and B , first determine the horizontal and vertical components of each vector. These are the dotted vectors A_x , A_y , B_x , and B_y shown in the image.

Step 2. Find the components of the resultant along each axis by adding the components of the individual vectors along that axis. That is, as shown in Figure 3.3.6, the x -component of the resultant is $R_x = A_x + B_x$ and the y -component is $R_y = A_y + B_y$.

$$R_x = A_x + B_x \quad (3.3.10)$$

and

$$R_y = A_y + B_y. \quad (3.3.11)$$




Figure 3.3.7: The magnitude of the vectors A_x and B_x add to give the magnitude R_x of the resultant vector in the horizontal direction. Similarly, the magnitudes of the vectors A_y and B_y add to give the magnitude R_y of the resultant vector in the vertical direction.

Components along the same axis, say the x -axis, are vectors along the same line and, thus, can be added to one another like ordinary numbers. The same is true for components along the y -axis. (For example, a 9-block eastward walk could be taken in two legs, the first 3 blocks east and the second 6 blocks east, for a total of 9, because they are along the same direction.) So resolving vectors into components along common axes makes it easier to add them. Now that the components of R are known, its magnitude and direction can be found.

Step 3. To get the magnitude R of the resultant, use the Pythagorean theorem:

$$R = \sqrt{R_x^2 + R_y^2} \quad (3.3.12)$$

Step 4. To get the direction of the resultant:

$$\theta = \tan^{-1}(R_y/R_x) \quad (3.3.13)$$

The following example illustrates this technique for adding vectors using perpendicular components.

Example 3.3.1: Adding Vectors Using Analytical Methods

Add the vector A to the vector B shown in Figure, using perpendicular components along the x - and y -axes. The x - and y -axes are along the east–west and north–south directions, respectively. Vector A represents the first leg of a walk in which a person walks 53.0m in a direction 20.0° north of east. Vector B represents the second leg, a displacement of 34.0m in a direction 63.0° north of east.




Figure 3.3.8: Vector A has magnitude 53.0m and direction 20.0° north of the x -axis. Vector B has magnitude 34.0m and direction 63.0° north of the x -axis. You can use analytical methods to determine the magnitude and direction of R .

Strategy

The components of A and B along the x - and y -axes represent walking due east and due north to get to the same ending point. Once found, they are combined to produce the resultant.

Solution

Following the method outlined above, we first find the components of A and B along the x - and y -axes. Note that $A = 53.0\text{m}$, $\theta_A = 20.0^\circ$, $B = 34.0\text{m}$, and $\theta_B = 63.0^\circ$. We find the x -components by using $A_x = A\cos\theta$, which gives

$$A_x = A\cos\theta_A = (53.0\text{m})(\cos 20.0^\circ)(53.0\text{m})(0.940) = 49.8\text{m} \quad (3.3.14)$$

and

$$B_x = B\cos\theta_B = (34.0\text{m})(\cos 63.0^\circ)(34.0\text{m})(0.454) = 15.4\text{m}. \quad (3.3.15)$$

Similarly, the y -components are found using $A_y = A\sin\theta_A$:

$$A_y = A\sin\theta_A = (53.0\text{m})(\sin 20.0^\circ)(53.0\text{m})(0.342) = 18.1\text{m} \quad (3.3.16)$$

and

$$B_y = B \sin \theta_B = (34.0 \text{ m})(\sin 63.0^\circ)(34.0 \text{ m})(0.891) = 30.3 \text{ m}. \quad (3.3.17)$$

The x- and y-components of the resultant are thus

$$R_x = A_x + B_x = 49.8 \text{ m} + 15.4 \text{ m} = 65.2 \text{ m} \quad (3.3.18)$$

and

$$R_y = A_y + B_y = 18.1 \text{ m} + 30.3 \text{ m} = 48.4 \text{ m}. \quad (3.3.19)$$

Now we can find the magnitude of the resultant by using the Pythagorean theorem:

$$R = \sqrt{R_x^2 + R_y^2} = \sqrt{(65.2)^2 + (48.4)^2} \text{ m} \quad (3.3.20)$$

so that

$$R = 81.2 \text{ m}. \quad (3.3.21)$$

Finally, we find the direction of the resultant:

$$\theta = \tan^{-1}(R_y/R_x) = +\tan^{-1}(48.4/65.2). \quad (3.3.22)$$

Thus,

$$\theta = \tan^{-1}(0.742) = 36.6^\circ. \quad (3.3.23)$$




Figure 3.3.9: Using analytical methods, we see that the magnitude of R is 81.2 m and its direction is 36.6° north of east.

Discussion

This example illustrates the addition of vectors using perpendicular components. Vector subtraction using perpendicular components is very similar—it is just the addition of a negative vector.

Subtraction of vectors is accomplished by the addition of a negative vector. That is, $A - B \equiv A + (-B)$. Thus, *the method for the subtraction of vectors using perpendicular components is identical to that for addition*. The components of $-B$ are the negatives of the components of B . The x- and y-components of the resultant $A - B = R$ are thus

$$R_x = A_x + (-B_x) \quad (3.3.24)$$

and

$$R_y = A_y + (-B_y) \quad (3.3.25)$$

and the rest of the method outlined above is identical to that for addition. (See Figure 3.3.10)

Analyzing vectors using perpendicular components is very useful in many areas of physics, because perpendicular quantities are often independent of one another. The next module, Projectile Motion, is one of many in which using perpendicular components helps make the picture clear and simplifies the physics.





Figure 3.3.10. The components of $-B$ are the negatives of the components of B . The method of subtraction is the same as that for addition.

PHET EXPLORATIONS: VECTOR ADDITION

Learn how to add vectors. Drag vectors onto a graph, change their length and angle, and sum them together. The magnitude, angle, and components of each vector can be displayed in several formats.

Vector Addition




Explore 1D



Explore 2D



Lab



Equations



Summary

- The analytical method of vector addition and subtraction involves using the Pythagorean theorem and trigonometric identities to determine the magnitude and direction of a resultant vector.
- The steps to add vectors A and B using the analytical method are as follows:

Step 1: Determine the coordinate system for the vectors. Then, determine the horizontal and vertical components of each vector using the equations

$$A_x = A \cos \theta$$

$$B_x = B \cos \theta$$

and

$$A_y = A \sin \theta$$
$$B_y = B \sin \theta.$$

Step 2: Add the horizontal and vertical components of each vector to determine the components R_x and R_y of the resultant vector, \mathbf{R} :

$$R_x = A_x + B_x$$

and

$$R_y = A_y + B_y .$$

Step 3: Use the Pythagorean theorem to determine the magnitude, R , of the resultant vector \mathbf{R} :

$$R = \sqrt{R_x^2 + R_y^2} .$$

Step 4: Use a trigonometric identity to determine the direction, θ , of \mathbf{R} :

$$\theta = \tan^{-1}(R_y/R_x) .$$

Glossary

analytical method

the method of determining the magnitude and direction of a resultant vector using the Pythagorean theorem and trigonometric identities

This page titled [3.3: Vector Addition and Subtraction- Analytical Methods](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [OpenStax](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.