

CHAPTER OVERVIEW

11: Time-Independent Perturbation Theory

Consider the following very commonly occurring problem. The Hamiltonian of a quantum mechanical system is written

$$H = H_0 + H_1. \quad (11.1)$$

Here, H_0 is a simple Hamiltonian whose eigenvalues and eigenstates are known exactly. H_1 introduces some interesting additional physics into the problem, but is sufficiently complicated that when we add it to H_0 we can no longer find the exact energy eigenvalues and eigenstates. However, H_1 can, in some sense (which we shall specify more precisely later on), be regarded as small compared to H_0 . Can we find approximate eigenvalues and eigenstates of the modified Hamiltonian, $H_0 + H_1$, by performing some sort of perturbation expansion about the eigenvalues and eigenstates of the original Hamiltonian, H_0 ? Let us investigate.

Incidentally, in this chapter, we shall only discuss so-called *time-independent perturbation theory*, in which the modification to the Hamiltonian, H_1 , has no explicit dependence on time. It is also assumed that the unperturbed Hamiltonian, H_0 , is time independent.

- [11.1: Exercises](#)
- [11.2: Improved Notation](#)
- [11.3: Two-State System](#)
- [11.4: Non-Degenerate Perturbation Theory](#)
- [11.5: Quadratic Stark Effect](#)
- [11.6: Degenerate Perturbation Theory](#)
- [11.7: Linear Stark Effect](#)
- [11.8: Fine Structure of Hydrogen](#)
- [11.9: Zeeman Effect](#)
- [11.10: Hyperfine Structure](#)

This page titled [11: Time-Independent Perturbation Theory](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).