

University of Texas at Austin  
Introductory Quantum Mechanics

Richard Fitzpatrick

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (<https://LibreTexts.org>) and like the hundreds of other texts available within this powerful platform, it is freely available for reading, printing and "consuming." Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects.

Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



The LibreTexts mission is to unite students, faculty and scholars in a cooperative effort to develop an easy-to-use online platform for the construction, customization, and dissemination of OER content to reduce the burdens of unreasonable textbook costs to our students and society. The LibreTexts project is a multi-institutional collaborative venture to develop the next generation of open-access texts to improve postsecondary education at all levels of higher learning by developing an Open Access Resource environment. The project currently consists of 14 independently operating and interconnected libraries that are constantly being optimized by students, faculty, and outside experts to supplant conventional paper-based books. These free textbook alternatives are organized within a central environment that is both vertically (from advance to basic level) and horizontally (across different fields) integrated.

The LibreTexts libraries are Powered by [NICE CXOne](#) and are supported by the Department of Education Open Textbook Pilot Project, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptations contact [info@LibreTexts.org](mailto:info@LibreTexts.org). More information on our activities can be found via Facebook (<https://facebook.com/Libretexts>), Twitter (<https://twitter.com/libretexts>), or our blog (<http://Blog.Libretexts.org>).

This text was compiled on 04/01/2025

# TABLE OF CONTENTS

## Licensing

## 1: Probability Theory

- 1.1: What is Probability?
- 1.2: Combining Probabilities
- 1.3: Mean, Variance, and Standard Deviation
- 1.4: Continuous Probability Distributions
- 1.5: Exercises

## 2: Wave-Particle Duality

- 2.1: Wavefunctions
- 2.2: Plane-Waves
- 2.3: Representation of Waves via Complex Functions
- 2.4: Classical Light-Waves
- 2.5: Photoelectric Effect
- 2.6: Quantum Theory of Light
- 2.7: Classical Interferences of Light Waves
- 2.8: Quantum Interference of Light
- 2.9: Particles
- 2.10: Wave-Packets
- 2.11: Evolution of Wave-Packets
- 2.12: Schrodinger's Equation and Wavefunction Collapse
- 2.13: Exercises

## 3: Fundamentals of Quantum Mechanics

- 3.1: Schrodinger's Equation
- 3.2: Normalization of the Wavefunction
- 3.3: Expectation Values (Averages) and Variances
- 3.4: Ehrenfest's Theorem
- 3.5: Operators
- 3.6: Momentum Representation
- 3.7: Heisenberg's Uncertainty Principle
- 3.8: Eigenstates and Eigenvalues
- 3.9: Measurement
- 3.10: Stationary States
- 3.11: Exercises

## 4: One-Dimensional Potentials

- 4.1: Infinite Potential Well
- 4.2: Square Potential Barrier
- 4.3: WKB Approximation
- 4.4: Cold Emission
- 4.5: Alpha Decay
- 4.6: Square Potential Well
- 4.7: Simple Harmonic Oscillator
- 4.E: One-Dimensional Potentials (Exercises)

## 5: Multi-Particle Systems

- 5.1: Fundamental Concepts of Multi-Particle Systems
- 5.2: Non-interacting Particles
- 5.3: Two-Particle Systems
- 5.4: Identical Particles
- 5.E: Multi-Particle Systems (Exercises)

## 6: Three-Dimensional Quantum Mechanics

- 6.1: Fundamental Concepts
- 6.2: Particle in Box
- 6.3: Degenerate Electron Gases
- 6.4: White Dwarf Stars
- 6.5: Exercises

## 7: Orbital Angular Momentum

- 7.1: Angular Momentum Operators
- 7.2: Representation of Angular Momentum
- 7.3: Eigenstates of Angular Momentum
- 7.4: Eigenvalues of  $L_z$
- 7.5: Eigenvalues of  $L^2$
- 7.6: Spherical Harmonics
- 7.E: Orbital Angular Momentum (Exercises)

## 8: Central Potentials

- 8.1: Derivation of Radial Equation
- 8.2: Infinite Spherical Potential Well
- 8.3: Hydrogen Atom
- 8.4: Rydberg Formula
- 8.E: Central Potentials (Exercises)

## 9: Spin Angular Momentum

- 9.1: Spin Operators
- 9.2: Spin Space
- 9.3: Eigenstates of  $S_z$  and  $S^2$
- 9.4: Pauli Representation
- 9.5: Spin Precession
- 9.E: Spin Angular Momentum (Exercises)

## 10: Addition of Angular Momentum

- 10.1: General Principles of Angular Momentum
- 10.2: Angular Momentum in Hydrogen Atom
- 10.3: Two Spin One-Half Particles
- 10.E: Addition of Angular Momentum (Exercises)

## 11: Time-Independent Perturbation Theory

- 11.1: Exercises
- 11.2: Improved Notation
- 11.3: Two-State System



- 11.4: Non-Degenerate Perturbation Theory
- 11.5: Quadratic Stark Effect
- 11.6: Degenerate Perturbation Theory
- 11.7: Linear Stark Effect
- 11.8: Fine Structure of Hydrogen
- 11.9: Zeeman Effect
- 11.10: Hyperfine Structure

## 12: Time-Dependent Perturbation Theory

- 12.1: Preliminary Analysis
- 12.2: Two-State System
- 12.3: Spin Magnetic Resonance
- 12.4: Perturbation Expansion
- 12.5: Harmonic Perturbation
- 12.6: Electromagnetic Radiation
- 12.7: Electric Dipole Approximation
- 12.8: Spontaneous Emission
- 12.9: Radiation from Harmonic Oscillator
- 12.10: Selection Rules (Hydrogen Atoms)
- 12.11: 2P-1S Transitions in Hydrogen
- 12.12: Intensity Rules
- 12.13: Forbidden Transitions
- 12.E: Time-Dependent Perturbation Theory (Exercises)

## 13: Variational Methods

- 13.1: Variational Principle
- 13.2: Helium Atom
- 13.3: Hydrogen Molecule Ion

## 14: Scattering Theory

- 14.1: Fundamentals of Scattering Theory
- 14.2: Born Approximation
- 14.3: Partial Waves
- 14.4: Optical Theorem
- 14.5: Determination of Phase-Shifts
- 14.6: Hard-Sphere Scattering
- 14.7: Low-Energy Scattering
- 14.8: Resonances

## 15: Bibliography

- 15.1: Section 1-
- 15.2: Section 2-
- 15.3: Section 3-
- 15.4: Section 4-
- 15.5: Section 5-
- 15.6: Section 6-

## Index

[Index](#)

[Glossary](#)

[Detailed Licensing](#)

## Licensing

---

*A detailed breakdown of this resource's licensing can be found in [Back Matter/Detailed Licensing](#).*

## CHAPTER OVERVIEW

### 1: Probability Theory

This chapter is devoted to a brief, and fairly low-level, introduction to a branch of mathematics known as *probability theory*.

[1.1: What is Probability?](#)

[1.2: Combining Probabilities](#)

[1.3: Mean, Variance, and Standard Deviation](#)

[1.4: Continuous Probability Distributions](#)

[1.5: Exercises](#)

---

This page titled [1: Probability Theory](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 1.1: What is Probability?

---

What is the scientific definition of probability? Consider an observation made on a general system,  $S$ . This can result in any one of a number of different possible outcomes. Suppose that we wish to find the probability of some general outcome,  $X$ . In order to ascribe a probability, we have to consider the system as a member of a large set,  $\Sigma$ , of similar, and similarly prepared, systems. Mathematicians call such a group an *ensemble* (which is just the French for “group”). So, consider an ensemble,  $\Sigma$ , of similar systems,  $S$ . The probability of the outcome  $X$  is defined as the ratio of the number of systems in the ensemble that exhibit this outcome to the total number of systems, in the limit that the latter number tends to infinity. We can write this symbolically as

$$P(X) = \lim_{\Omega(\Sigma) \rightarrow \infty} \frac{\Omega(X)}{\Omega(\Sigma)}, \quad (1.1.1)$$

where  $\Omega(\Sigma)$  is the total number of systems in the ensemble, and  $\Omega(X)$  the number of systems exhibiting the outcome  $X$ . We can see that the probability  $P(X)$  must be a real number lying between 0 and 1. The probability is zero if no systems exhibit the outcome  $X$ , even when the number of systems goes to infinity. This is just another way of saying that there is no chance of the outcome  $X$ . The probability is unity if all systems exhibit the outcome  $X$  in the limit that the number of systems goes to infinity. This is another way of saying that the outcome  $X$  is bound to occur.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [1.1: What is Probability?](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 1.2: Combining Probabilities

Consider two distinct possible outcomes,  $X$  and  $Y$ , of an observation made on the system  $S$ , with probabilities of occurrence  $P(X)$  and  $P(Y)$ , respectively. Let us determine the probability of obtaining either the outcome  $X$  or the outcome  $Y$ , which we shall denote  $P(X \mid Y)$ . From the basic definition of probability,

$$P(X \mid Y) = \lim_{\Omega(\Sigma) \rightarrow \infty} \frac{\Omega(X \mid Y)}{\Omega(\Sigma)}, \quad (1.2.1)$$

where  $\Omega(X \mid Y)$  is the number of systems in the ensemble that exhibit either the outcome  $X$  or the outcome  $Y$ . Now,

$$\Omega(X \mid Y) = \Omega(X) + \Omega(Y) \quad (1.2.2)$$

if the outcomes  $X$  and  $Y$  are mutually exclusive (which must be the case if they are two distinct outcomes). Thus,

$$P(X \mid Y) = P(X) + P(Y), \quad (1.2.3)$$

which means that the probability of obtaining either the outcome  $X$  or the outcome  $Y$  is the sum of the individual probabilities of  $X$  and  $Y$ . For instance, with a six-sided die the probability of throwing any particular number (one to six) is  $1/6$ , because all of the possible outcomes are considered to be equally likely. It follows, from what has just been said, that the probability of throwing either a one or a two is simply  $1/6 + 1/6$ , which equals  $1/3$ .

Let us denote all of the  $M$ , say, possible outcomes of an observation made on the system  $S$  by  $X_i$ , where  $i$  runs from 1 to  $M$ . Let us determine the probability of obtaining any one of these outcomes. This quantity is unity, from the basic definition of probability, because each of the systems in the ensemble must exhibit one of the possible outcomes. But, this quantity is also equal to the sum of the probabilities of all the individual outcomes, by Equation (1.2.3), so we conclude that this sum is equal to unity: that is,

$$\sum_{i=1, M} P(X_i) = 1. \quad (1.2.4)$$

The previous expression is called the *normalization condition*, and must be satisfied by any complete set of probabilities.

There is another way in which we can combine probabilities. Suppose that we make an observation on a system picked at random from the ensemble, and then pick a second similar system, completely independently, and make another observation. We are assuming that the first observation does not influence the second observation in any way. In other words, the two observations are *statistically independent* of one another. Let us determine the probability of obtaining the outcome  $X$  in the first system and obtaining the outcome  $Y$  in the second system, which we shall denote  $P(X \otimes Y)$ . In order to determine this probability, we have to form an ensemble of all the possible pairs of systems that we could choose from the ensemble  $(\Sigma)$ . Let us denote this ensemble  $\Sigma \otimes \Sigma$ . The number of pairs of systems in this new ensemble is just the square of the number of systems in the original ensemble, so

$$\Omega(\Sigma \otimes \Sigma) = \Omega(\Sigma) \Omega(\Sigma) \quad (1.2.5)$$

Furthermore, the number of pairs of systems in the ensemble  $\Sigma \otimes \Sigma$  that exhibit the outcome  $X$  in the first system and the outcome  $Y$  in the second system is simply the product of the number of systems that exhibit the outcome  $X$  and the number of systems that exhibit the outcome  $Y$  in the original ensemble, so that

$$\Omega(X \otimes Y) = \Omega(X) \Omega(Y). \quad (1.2.6)$$

It follows from the basic definition of probability that

$$P(X \otimes Y) = \lim_{\Omega(\Sigma) \rightarrow \infty} \frac{\Omega(X \otimes Y)}{\Omega(\Sigma \otimes \Sigma)} = P(X) P(Y). \quad (1.2.7)$$

Thus, the probability of obtaining the outcomes  $X$  and  $Y$  in two statistically independent observations is the product of the individual probabilities of  $X$  and  $Y$ . For instance, the probability of throwing a one and then a two on a six-sided die is  $1/6 \times 1/6$ , which equals  $1/36$ .

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [1.2: Combining Probabilities](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 1.3: Mean, Variance, and Standard Deviation

What is meant by the mean or average of a quantity? Suppose that we wish to calculate the average age of undergraduates at the University of Texas at Austin. We could go to the central administration building and find out how many eighteen year-olds, nineteen year-olds, et cetera, were currently enrolled. We would then write something like

$$\text{Average Age} \simeq \frac{N_{18} \times 18 + N_{19} \times 19 + N_{20} \times 20 + \cdots}{N_{18} + N_{19} + N_{20} + \cdots}, \quad (1.3.1)$$

where  $N_{18}$  is the number of enrolled eighteen year-olds, et cetera. The probability that a randomly picked student is eighteen is

$$P_{18} \simeq \frac{N_{18}}{N_{\text{students}}}, \quad (1.3.2)$$

where  $N_{\text{students}} = N_{18} + N_{19} + N_{20} + \cdots$  is the total number of enrolled students. (Actually, this definition is only accurate in the limit that  $N_{\text{students}}$  is very large.) We can now see that the average age takes the form

$$\text{Average Age} \simeq P_{18} \times 18 + P_{19} \times 19 + P_{20} \times 20 + \cdots. \quad (1.3.3)$$

Finally, because there is nothing unique about the age distribution of students at UT Austin, for a general variable  $u$ , which can take on any one of  $M$  possible values  $u_1, u_2, \cdots, u_M$ , with corresponding probabilities  $P(u_1), P(u_2), \cdots, P(u_M)$ , the *mean* or *average* value of  $u$ , which is denoted  $\langle u \rangle$ , is defined

$$\langle u \rangle \equiv \sum_{i=1, M} P(u_i) u_i. \quad (1.3.4)$$

Suppose that  $f(u)$  is some function of  $u$ . Thus, for each of the  $M$  possible values of  $u$ , there is a corresponding value of  $f(u)$  that occurs with the same probability. That is,  $f(u_1)$  corresponds to  $u_1$ , and occurs with the probability  $P(u_1)$ , and so on. It follows from our previous definition that the mean value of  $f(u)$  is given by

$$\langle f(u) \rangle \equiv \sum_{i=1, M} P(u_i) f(u_i). \quad (1.3.5)$$

Suppose that  $f(u)$  and  $g(u)$  are two general functions of  $u$ . It follows that

$$\langle f(u) + g(u) \rangle = \sum_{i=1, M} P(u_i) [f(u_i) + g(u_i)] = \sum_{i=1, M} P(u_i) f(u_i) + \sum_{i=1, M} P(u_i) g(u_i), \quad (1.3.6)$$

so

$$\langle f(u) + g(u) \rangle = \langle f(u) \rangle + \langle g(u) \rangle. \quad (1.3.7)$$

Finally, if  $c$  is a general constant then it is clear that

$$\langle c f(u) \rangle = c \langle f(u) \rangle. \quad (1.3.8)$$

We now know how to define the mean value of the general variable,  $u$ . Let us consider how we might characterize the scatter around the mean value. We could investigate the deviation of  $u$  from its mean value,  $\langle u \rangle$ , which is denoted

$$\Delta u \equiv u - \langle u \rangle. \quad (1.3.9)$$

In fact, this is not a particularly interesting quantity because its average is obviously zero: that is,

$$\langle \Delta u \rangle = \langle (u - \langle u \rangle) \rangle = \langle u \rangle - \langle u \rangle = 0. \quad (1.3.10)$$

This is another way of saying that the average deviation from the mean vanishes. A more interesting quantity is the square of the deviation. The average value of this quantity,

$$\langle (\Delta u)^2 \rangle = \sum_{i=1, M} P(u_i) (u_i - \langle u \rangle)^2, \quad (1.3.11)$$

is usually called the *variance*. The variance is a positive real number, unless there is no scatter at all in the distribution, so that all possible values of  $u$  correspond to the mean value,  $\langle u \rangle$ , in which case it takes the value zero. Note that



$$\langle (u - \langle u \rangle)^2 \rangle = \langle (u^2 - 2u \langle u \rangle + \langle u \rangle^2) \rangle = \langle u^2 \rangle - 2 \langle u \rangle \langle u \rangle + \langle u \rangle^2, \quad (1.3.12)$$

which yields the following useful relationship

$$\langle (\Delta u)^2 \rangle = \langle u^2 \rangle - \langle u \rangle^2. \quad (1.3.13)$$

The variance of  $u$  is proportional to the square of the scatter of  $u$  around its mean value. A more useful measure of the scatter is given by the square root of the variance,

$$\sigma_u = [\langle (\Delta u)^2 \rangle]^{1/2}, \quad (1.3.14)$$

which is usually called the *standard deviation* of  $u$ . The standard deviation is essentially the width of the range over which  $u$  is distributed around its mean value,  $\langle u \rangle$ .

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [1.3: Mean, Variance, and Standard Deviation](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 1.4: Continuous Probability Distributions

---

Suppose that the variable  $u$  can take on a continuous range of possible values. In general, we expect the probability that  $u$  takes on a value in the range  $u$  to  $u + du$  to be directly proportional to  $du$ , in the limit that  $du \rightarrow 0$ . In other words,

$$P(u \in u : u + du) = P(u) du, \quad (1.4.1)$$

where  $P(u)$  is known as the *probability density*. The earlier results (1.2.4), (1.3.4), and (1.3.11) generalize in a straightforward manner to give:

$$\begin{aligned} 1 &= \int_{-\infty}^{\infty} P(u) du, \\ \langle u \rangle &= \int_{-\infty}^{\infty} P(u) u du, \\ \langle (\Delta u)^2 \rangle &= \int_{-\infty}^{\infty} P(u) (u - \langle u \rangle)^2 du = \langle u^2 \rangle - \langle u \rangle^2, \end{aligned}$$

respectively.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [1.4: Continuous Probability Distributions](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 1.5: Exercises

1. Show that the probability of throwing 6 points or less with three (six-sided) dice is  $5/54$ .
2. A battery of total emf  $V$  is connected to a resistor  $R$ . As a result, an amount of power  $P = V^2/R$  is dissipated in the resistor. The battery itself consists of  $N$  individual cells connected in series, so that  $V$  is equal to the sum of the emf's of all these cells. The battery is old, however, so that not all cells are in perfect condition. Thus, there is a probability  $p$  that the emf of any individual cell has its normal value  $v$ ; and a probability  $1 - p$  that the emf of any individual cell is zero because the cell has become internally shorted. The individual cells are statistically independent of each other. Under these conditions, show that the mean power,  $\langle P \rangle$ , dissipated in the resistor, is

$$\langle P \rangle = \frac{p^2 V^2}{R} \left[ 1 + \frac{(1-p)}{Np} \right]. \quad (1.5.1)$$

3. In the “game” of Russian roulette, the player inserts a single bullet into the drum of a revolver, leaving the other five chambers of the drum empty. The player then spins the drum, aims at his/her head, and pulls the trigger.
  1. What is the probability of the player still being alive after playing the game  $N$  times?
  2. What is the probability of the player surviving  $N - 1$  turns in this game, and then being shot the  $N$ th time he/she pulls the trigger?
  3. What is the mean number of times the player gets to pull the trigger?
4. Suppose that the probability density for the speed,  $s$ , of a car on a road is given by

$$P(s) = A s \exp\left(-\frac{s}{s_0}\right), \quad (1.5.2)$$

where  $0 \leq s \leq \infty$ . Here,  $A$  and  $s_0$  are positive constants. More explicitly,  $P(s) ds$  gives the probability that a car has a speed between  $s$  and  $s + ds$ .

1. Determine  $A$  in terms of  $s_0$ .
  2. What is the mean value of the speed?
  3. What is the “most probable” speed: that is, the speed for which the probability density has a maximum?
  4. What is the probability that a car has a speed more than three times as large as the mean value?
5. An radioactive atom has a uniform decay probability per unit time  $w$ . In other words, the probability of decay in a time interval  $dt$  is  $w dt$ . Let  $P(t)$  be the probability of the atom not having decayed at time  $t$ , given that it was created at time  $t = 0$ . Demonstrate that

$$P(t) = e^{-wt}. \quad (1.5.3)$$

What is the mean lifetime of the atom?

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled 1.5: Exercises is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## CHAPTER OVERVIEW

### 2: Wave-Particle Duality

In classical mechanics, waves and particles are two completely distinct types of physical entity. Waves are continuous and spatially extended, whereas particles are discrete and have little or no spatial extent. However, in quantum mechanics, waves sometimes act as particles, and particles sometimes act as waves—this strange behavior is known as *wave-particle duality*. In this chapter, we shall examine how wave-particle duality shapes the general features of quantum mechanics.

- [2.1: Wavefunctions](#)
- [2.2: Plane-Waves](#)
- [2.3: Representation of Waves via Complex Functions](#)
- [2.4: Classical Light-Waves](#)
- [2.5: Photoelectric Effect](#)
- [2.6: Quantum Theory of Light](#)
- [2.7: Classical Interferences of Light Waves](#)
- [2.8: Quantum Interference of Light](#)
- [2.9: Particles](#)
- [2.10: Wave-Packets](#)
- [2.11: Evolution of Wave-Packets](#)
- [2.12: Schrodinger's Equation and Wavefunction Collapse](#)
- [2.13: Exercises](#)

---

This page titled [2: Wave-Particle Duality](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 2.1: Wavefunctions

A wave is defined as a disturbance in some physical system that is periodic in both space and time. In one dimension, a wave is generally represented in terms of a *wavefunction*: for instance,

$$\psi(x, t) = A \cos(kx - \omega t + \varphi), \quad (2.1.1)$$

where  $x$  is a position coordinate,  $t$  represents time, and  $A, k, \omega > 0$ . For example, if we are considering a sound wave then  $\psi(x, t)$  might correspond to the pressure perturbation associated with the wave at position  $x$  and time  $t$ . On the other hand, if we are considering a light-wave then  $\psi(x, t)$  might represent the wave's transverse electric field. As is well known, the cosine function,  $\cos \theta$ , is periodic in its argument,  $\theta$ , with period  $2\pi$ : in other words,  $\cos(\theta + 2\pi) = \cos \theta$  for all  $\theta$ . The function also oscillates between the minimum and maximum values  $-1$  and  $+1$ , respectively, as  $\theta$  varies. It follows that the wavefunction (2.1.1) is periodic in  $x$  with period  $\lambda = 2\pi/k$ . In other words,  $\psi(x + \lambda, t) = \psi(x, t)$  for all  $x$  and  $t$ . Moreover, the wavefunction is periodic in  $t$  with period  $T = 2\pi/\omega$ . In other words,  $\psi(x, t + T) = \psi(x, t)$  for all  $x$  and  $t$ . Finally, the wavefunction oscillates between the minimum and maximum values  $-A$  and  $+A$ , respectively, as  $x$  and  $t$  vary. The spatial period of the wave,  $\lambda$ , is known as its *wavelength*, and the temporal period,  $T$ , is called its *period*. Furthermore, the quantity  $A$  is termed the wave *amplitude*, the quantity  $k$  the *wavenumber*, and the quantity  $\omega$  the wave *angular frequency*. Note that the units of  $\omega$  are radians per second. The conventional wave *frequency*, in cycles per second (otherwise known as hertz), is  $\nu = 1/T = \omega/2\pi$ . Finally, the quantity  $\varphi$ , appearing in expression (2.1.1), is termed the *phase angle*, and determines the exact positions of the wave maxima and minima at a given time. In fact, the maxima are located at  $kx - \omega t + \varphi = j2\pi$ , where  $j$  is an integer. This follows because the maxima of  $\cos \theta$  occur at  $\theta = j2\pi$ . Note that a given maximum satisfies  $x = (j - \varphi/2\pi)\lambda + vt$ , where  $v = \omega/k$ . It follows that the maximum, and, by implication, the whole wave, propagates in the positive  $x$ -direction at the velocity  $\omega/k$ . Analogous reasoning reveals that

$$\psi(x, t) = A \cos(-kx - \omega t + \varphi) = A \cos(kx + \omega t - \varphi), \quad (2.1.2)$$

is the wavefunction of a wave of amplitude  $A$ , wavenumber  $k$ , angular frequency  $\omega$ , and phase angle  $\varphi$ , that propagates in the negative  $x$ -direction at the velocity  $\omega/k$ .

---

This page titled 2.1: Wavefunctions is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 2.2: Plane-Waves

As we have just seen, a wave of amplitude  $A$ , wavenumber  $k$ , angular frequency  $\omega$ , and phase angle  $\varphi$ , propagating in the positive  $x$ -direction, is represented by the following wavefunction:

$$\psi(x, t) = A \cos(kx - \omega t + \varphi). \quad (2.2.1)$$

This type of wave is conventionally termed a *one-dimensional plane-wave*. It is one-dimensional because its associated wavefunction only depends on the single Cartesian coordinate,  $x$ . Furthermore, it is a plane-wave because the wave maxima, which are located at

$$kx - \omega t + \varphi = j2\pi, \quad (2.2.2)$$

where  $j$  is an integer, consist of a series of parallel planes, normal to the  $x$ -axis, that are equally spaced a distance  $\lambda = 2\pi/k$  apart, and propagate along the positive  $x$ -axis at the velocity  $v = \omega/k$ . These conclusions follow because Equation (2.2.2) can be rewritten in the form

$$x = d, \quad (2.2.3)$$

where  $d = (j - \varphi/2\pi)\lambda + vt$ . Moreover, as is well known, Equation (2.2.3) is the equation of a plane, normal to the  $x$ -axis, whose distance of closest approach to the origin is  $d$ .




Figure 1: The solution of  $\mathbf{n} \cdot \mathbf{r} = d$  is a plane.

The previous equation can also be written in the coordinate-free form

$$\mathbf{n} \cdot \mathbf{r} = d, \quad (2.2.4)$$

where  $\mathbf{n} = (1, 0, 0)$  is a unit vector directed along the positive  $x$ -axis, and  $\mathbf{r} = (x, y, z)$  represents the vector displacement of a general point from the origin. Because there is nothing special about the  $x$ -direction, it follows that if  $\mathbf{n}$  is reinterpreted as a unit vector pointing in an arbitrary direction then Equation (2.2.4) can be reinterpreted as the general equation of a plane. As before, the plane is normal to  $\mathbf{n}$ , and its distance of closest approach to the origin is  $d$ . See Figure [f10.1]. This observation allows us to write the three-dimensional equivalent to the wavefunction (2.2.1) as

$$\psi(\mathbf{r}, t) = A \cos(\mathbf{k} \cdot \mathbf{r} - \omega t + \varphi), \quad (2.2.5)$$

where the constant vector  $\mathbf{k} = (k_x, k_y, k_z) = k\mathbf{n}$  is called the *wavevector*. The wave represented previously is conventionally termed a *three-dimensional plane-wave*. It is three-dimensional because its wavefunction,  $\psi(\mathbf{r}, t)$ , depends on all three Cartesian coordinates. Moreover, it is a plane-wave because the wave maxima are located at

$$\mathbf{k} \cdot \mathbf{r} - \omega t + \varphi = j2\pi, \quad (2.2.6)$$

or

$$\mathbf{n} \cdot \mathbf{r} = (j - \varphi/2\pi) \lambda + vt, \quad (2.2.7)$$

where  $\lambda = 2\pi/k$ , and  $v = \omega/k$ . Note that the wavenumber,  $k$ , is the magnitude of the wavevector,  $\mathbf{k}$ : that is,  $k \equiv |\mathbf{k}|$ . It follows, by comparison with Equation (2.2.4), that the wave maxima consist of a series of parallel planes, normal to the wavevector, that are equally spaced a distance  $\lambda$  apart, and that propagate in the  $\mathbf{k}$ -direction at the velocity  $v$ . See Figure [f10.2]. Hence, the direction of the wavevector specifies the wave propagation direction, whereas its magnitude determines the wavenumber,  $k$ , and, thus, the wavelength,  $\lambda = 2\pi/k$ .




Figure 2: Wave maxima associated with a three-dimensional plane wave.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled 2.2: Plane-Waves is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 2.3: Representation of Waves via Complex Functions

In mathematics, the symbol  $i$  is conventionally used to represent the square-root of minus one: in other words, one of the solutions of  $i^2 = -1$ . Now, a *real number*,  $x$  (say), can take any value in a continuum of different values lying between  $-\infty$  and  $+\infty$ . On the other hand, an *imaginary number* takes the general form  $iy$ , where  $y$  is a real number. It follows that the square of a real number is a positive real number, whereas the square of an imaginary number is a negative real number. In addition, a general *complex number* is written

$$z = x + iy, \quad (2.3.1)$$

where  $x$  and  $y$  are real numbers. In fact,  $x$  is termed the *real part* of  $z$ , and  $y$  the *imaginary part* of  $z$ . This is written mathematically as  $x = \text{Re}(z)$  and  $y = \text{Im}(z)$ . Finally, the *complex conjugate* of  $z$  is defined  $z^* = x - iy$ .

Just as we can visualize a real number as a point lying on an infinite straight-line, we can visualize a complex number as a point lying in an infinite plane. The coordinates of the point in question are the real and imaginary parts of the number: that is,  $z \equiv (x, y)$ . This idea is illustrated in Figure [f13.2]. The distance,  $r = (x^2 + y^2)^{1/2}$ , of the representative point from the origin is termed the *modulus* of the corresponding complex number,  $z$ . This is written mathematically as  $|z| = (x^2 + y^2)^{1/2}$ . Incidentally, it follows that  $zz^* = x^2 + y^2 = |z|^2$ . The angle,  $\theta = \tan^{-1}(y/x)$ , that the straight-line joining the representative point to the origin subtends with the real axis is termed the *argument* of the corresponding complex number,  $z$ . This is written mathematically as  $\arg(z) = \tan^{-1}(y/x)$ . It follows from standard trigonometry that  $x = r \cos \theta$ , and  $y = r \sin \theta$ . Hence,  $z = r \cos \theta + ir \sin \theta$ .




Figure 3: Representation of a complex number as a point in a plane.

Complex numbers are often used to represent wavefunctions. All such representations depend ultimately on a fundamental mathematical identity, known as *Euler's theorem*, that takes the form

$$e^{i\phi} \equiv \cos \phi + i \sin \phi, \quad (2.3.2)$$

where  $\phi$  is a real number. Incidentally, given that  $z = r \cos \theta + ir \sin \theta = r(\cos \theta + i \sin \theta)$ , where  $z$  is a general complex number,  $r = |z|$  its modulus, and  $\theta = \arg(z)$  its argument, it follows from Euler's theorem that any complex number,  $z$ , can be written

$$z = r e^{i\theta}, \quad (2.3.3)$$

where  $r = |z|$  and  $\theta = \arg(z)$  are real numbers.

A one-dimensional wavefunction takes the general form

$$\psi(x, t) = A \cos(kx - \omega t + \varphi), \quad (2.3.4)$$

where  $A$  is the wave amplitude,  $k$  the wavenumber,  $\omega$  the angular frequency, and  $\varphi$  the phase angle. Consider the complex wavefunction



$$\psi(x, t) = \psi_0 e^{i(kx - \omega t)}, \quad (2.3.5)$$

where  $\psi_0$  is a complex constant. We can write

$$\psi_0 = A e^{i\varphi}, \quad (2.3.6)$$

where  $A$  is the modulus, and  $\varphi$  the argument, of  $\psi_0$ . Hence, we deduce that

$$\operatorname{Re} \left[ \psi_0 e^{i(kx - \omega t)} \right] = \operatorname{Re} \left[ A e^{i\varphi} e^{i(kx - \omega t)} \right] = \operatorname{Re} \left[ A e^{i(kx - \omega t + \varphi)} \right] = A \operatorname{Re} \left[ e^{i(kx - \omega t + \varphi)} \right].$$

Thus, it follows from Euler's theorem, and Equation (2.3.4), that

$$\operatorname{Re} \left[ \psi_0 e^{i(kx - \omega t)} \right] = A \cos(kx - \omega t + \varphi) = \psi(x, t). \quad (2.3.7)$$

In other words, a general one-dimensional real wavefunction, (2.3.4), can be represented as the real part of a complex wavefunction of the form (2.3.5). For ease of notation, the “take the real part” aspect of the previous expression is usually omitted, and our general one-dimension wavefunction is simply written

$$\psi(x, t) = \psi_0 e^{i(kx - \omega t)}. \quad (2.3.8)$$

The main advantage of the complex representation, (2.3.8), over the more straightforward real representation, (2.3.4), is that the former enables us to combine the amplitude,  $A$ , and the phase angle,  $\varphi$ , of the wavefunction into a single complex amplitude,  $\psi_0$ . Finally, the three-dimensional generalization of the previous expression is

$$\psi(\mathbf{r}, t) = \psi_0 e^{i(\mathbf{k} \cdot \mathbf{r} - \omega t)}, \quad (2.3.9)$$

where  $\mathbf{k}$  is the wavevector.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [2.3: Representation of Waves via Complex Functions](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 2.4: Classical Light-Waves

Consider a classical, monochromatic, linearly-polarized, plane light-wave, propagating through a vacuum in the  $x$ -direction. It is convenient to characterize a light-wave (which is, of course, a type of electromagnetic wave) by specifying its associated electric field. Suppose that the wave is polarized such that this electric field oscillates in the  $y$ -direction. (According to standard electromagnetic theory, the magnetic field oscillates in the  $z$ -direction, in phase with the electric field, with an amplitude which is that of the electric field divided by the velocity of light in vacuum. ) Now, the electric field can be conveniently represented in terms of a *complex wavefunction*:

$$\psi(x, t) = \bar{\psi} e^{i(kx - \omega t)}. \quad (2.4.1)$$

Here,  $i = \sqrt{-1}$ ,  $k$  and  $\omega$  are real parameters, and  $\bar{\psi}$  is a complex wave amplitude. By convention, the physical electric field is the real part of the previous expression. Suppose that

$$\bar{\psi} = |\bar{\psi}| e^{i\varphi}, \quad (2.4.2)$$

where  $\varphi$  is real. It follows that the physical electric field takes the form

$$E_y(x, t) = \text{Re}[\psi(x, t)] = |\bar{\psi}| \cos(kx - \omega t + \varphi), \quad (2.4.3)$$

where  $|\bar{\psi}|$  is the amplitude of the electric oscillation,  $k$  the wavenumber,  $\omega$  the angular frequency, and  $\varphi$  the phase angle. In addition,  $\lambda = 2\pi/k$  is the wavelength, and  $\nu = \omega/2\pi$  the frequency (in hertz).

According to standard electromagnetic theory, the frequency and wavelength of light-waves are related according to the well-known expression

$$c = \nu \lambda, \quad (2.4.4)$$

or, equivalently,

$$\omega = kc, \quad (2.4.5)$$

where  $c = 3 \times 10^8$  m/s is the velocity of light in vacuum. Equations (2.4.3) and (2.4.5) yield

$$E_y(x, t) = |\bar{\psi}| \cos(k[x - (\omega/k)t] + \varphi) = |\bar{\psi}| \cos(k[x - ct] + \varphi). \quad (2.4.6)$$

Note that  $E_y$  depends on  $x$  and  $t$  only via the combination  $x - ct$ . It follows that the wave maxima and minima satisfy

$$x - ct = \text{constant}. \quad (2.4.7)$$

Thus, the wave maxima and minima propagate in the  $x$ -direction at the fixed velocity

$$\frac{dx}{dt} = c. \quad (2.4.8)$$

An expression, such as Equation (2.4.5), that determines the wave angular frequency as a function of the wavenumber, is generally termed a *dispersion relation*. As we have already seen, and as is apparent from Equation (2.4.6), the maxima and minima of a plane-wave propagate at the characteristic velocity

$$v_p = \frac{\omega}{k}, \quad (2.4.9)$$

which is known as the *phase-velocity*. Hence, the dispersion relation (2.4.5) is effectively saying that the phase-velocity of a plane light-wave, propagating through a vacuum, always takes the fixed value  $c$ , irrespective of its wavelength or frequency.

From standard electromagnetic theory, the *energy density* (i.e., the energy per unit volume) of a plane light-wave is

$$U = \frac{E_y^2}{\epsilon_0}, \quad (2.4.10)$$

where  $\epsilon_0 = 8.85 \times 10^{-12}$  F/m is the *electrical permittivity of free space*. Hence, it follows from Equations (2.4.1) and (2.4.3) that

$$U \propto |\bar{\psi}|^2. \quad (2.4.11)$$

Furthermore, a light-wave possesses linear momentum, as well as energy. This momentum is directed along the wave's direction of propagation, and is of density

$$G = \frac{U}{c}. \quad (2.4.12)$$

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [2.4: Classical Light-Waves](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 2.5: Photoelectric Effect

The so-called *photoelectric effect*, by which a polished metal surface emits electrons when illuminated by visible and ultra-violet light, was discovered by Heinrich Hertz in 1887. The following facts regarding this effect can be established via careful observation. First, a given surface only emits electrons when the frequency of the light with which it is illuminated exceeds a certain threshold value, which is a property of the metal. Second, the current of photoelectrons, when it exists, is proportional to the intensity of the light falling on the surface. Third, the energy of the photoelectrons is independent of the light intensity, but varies linearly with the light frequency. These facts are inexplicable within the framework of classical physics.

In 1905, Albert Einstein proposed a radical new theory of light in order to account for the photoelectric effect. According to this theory, light of fixed frequency  $\nu$  consists of a collection of indivisible discrete packages, called *quanta*,<sup>1</sup> whose energy is

$$E = h \nu. \quad (2.5.1)$$

Here,  $h = 6.6261 \times 10^{-34}$  J s is a new constant of nature, known as *Planck's constant*. Incidentally,  $h$  is called Planck's constant, rather than Einstein's constant, because Max Planck first introduced the concept of the quantization of light, in 1900, while trying to account for the electromagnetic spectrum of a black body (i.e., a perfect emitter and absorber of electromagnetic radiation).

Suppose that the electrons at the surface of a metal lie in a potential well of depth  $W$ . In other words, the electrons have to acquire an energy  $W$  in order to be emitted from the surface. Here,  $W$  is generally called the *work function* of the surface, and is a property of the metal. Suppose that an electron absorbs a single quantum of light. Its energy therefore increases by  $h \nu$ . If  $h \nu$  is greater than  $W$  then the electron is emitted from the surface with residual kinetic energy

$$K = h \nu - W. \quad (2.5.2)$$

Otherwise, the electron remains trapped in the potential well, and is not emitted. Here, we are assuming that the probability of an electron simultaneously absorbing two or more light quanta is negligibly small compared to the probability of it absorbing a single light quantum (as is, indeed, the case for sufficiently low-intensity illumination). Incidentally, we can calculate Planck's constant, and the work function of the metal, by simply plotting the kinetic energy of the emitted photoelectrons as a function of the wave frequency, as shown in Figure [f1]. This plot is a straight-line whose slope is  $h$ , and whose intercept with the  $\nu$ -axis is  $W/h$ . Finally, the number of emitted electrons increases with the intensity of the light because the more intense the light, the larger the flux of light quanta onto the surface. Thus, Einstein's quantum theory is capable of accounting for all three of the previously mentioned observational facts regarding the photoelectric effect.




Figure 4: Variation of the kinetic energy  $K$  of photoelectrons with the wave-frequency  $\nu$ .

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled 2.5: Photoelectric Effect is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 2.6: Quantum Theory of Light

---

According to Einstein's quantum theory of light, a monochromatic light-wave of angular frequency  $\omega$ , propagating through a vacuum, can be thought of as a stream of particles, called *photons*, of energy

$$E = \hbar \omega, \quad (2.6.1)$$

where  $\hbar = h/2\pi = 1.0546 \times 10^{-34}$  J s. Because classical light-waves propagate at the fixed velocity  $c$ , it stands to reason that photons must also move at this velocity. According to Einstein's special theory of relativity, only massless particles can move at the speed of light in vacuum. Hence, photons must be massless. Special relativity also gives the following relationship between the energy  $E$  and the momentum  $p$  of a massless particle,

$$p = \frac{E}{c}. \quad (2.6.2)$$

Note that the previous relation is consistent with Equation (2.4.12), because if light is made up of a stream of photons, for which  $E/p = c$ , then the momentum density of light must be the energy density divided by  $c$ . It follows, from the previous two equations, that photons carry momentum

$$p = \hbar k \quad (2.6.3)$$

along their direction of motion, because  $\omega/c = k$  for a light-wave. [See Equation (2.4.5).]

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled 2.6: Quantum Theory of Light is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 2.7: Classical Interferences of Light Waves

Let us now consider the classical interference of light-waves. Figure [f2] shows a standard double-slit interference experiment in which monochromatic plane light-waves are normally incident on two narrow parallel slits that are situated a distance  $d$  apart. The light from the two slits is projected onto a screen a distance  $D$  behind them, where  $D \gg d$ .




Figure 5: Classical double-slit interference of light.

Consider some point on the screen that is located a distance  $y$  from the centre-line, as shown in the figure. Light from the first slit travels a distance  $x_1$  to get to this point, whereas light from the second slit travels a slightly different distance  $x_2$ . It is easily demonstrated that

$$\Delta x = x_2 - x_1 \simeq \frac{d}{D} y, \quad (2.7.1)$$

provided  $d \ll D$ . It follows from Equation (2.4.1), and the well-known fact that light-waves are superposable, that the wavefunction at the point in question can be written

$$\psi(y, t) \propto \psi_1(t) e^{i k x_1} + \psi_2(t) e^{i k x_2}, \quad (2.7.2)$$

where  $\psi_1$  and  $\psi_2$  are the wavefunctions at the first and second slits, respectively. However,

$$\psi_1 = \psi_2, \quad (2.7.3)$$

because the two slits are assumed to be illuminated by in-phase light-waves of equal amplitude. (Note that we are ignoring the difference in amplitude of the waves from the two slits at the screen, due to the slight difference between  $x_1$  and  $x_2$ , compared to the difference in their phases. This is reasonable provided  $D \gg \lambda$ .) The intensity (that is, the energy flux) of the light at some point on the projection screen is approximately equal to the energy density of the light at this point times the velocity of light (provided that  $y \ll D$ ). Hence, it follows from Equation (2.4.11) that the light intensity on the screen a distance  $y$  from the center-line is


$$I(y) \propto |\psi(y, t)|^2. \quad (2.7.4)$$

Using Equations (2.7.1)–(2.7.4), we obtain

$$I(y) \propto \cos^2 \left( \frac{k \Delta x}{2} \right) \simeq \cos^2 \left( \pi \frac{d}{D \lambda} y \right). \quad (2.7.5)$$

Figure [f3] shows the characteristic interference pattern corresponding to the previous expression. This pattern consists of equally-spaced light and dark bands of characteristic width

$$\Delta y = \frac{D \lambda}{d}. \quad (2.7.6)$$



### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

2.7: Classical Interferences of Light Waves is shared under a [not declared](#) license and was authored, remixed, and/or curated by LibreTexts.

## 2.8: Quantum Interference of Light

Let us now consider double-slit light interference from a quantum-mechanical point of view. According to quantum theory, light-waves consist of a stream of massless photons moving at the speed of light. Hence, we expect the two slits in Figure [f2] to be spraying photons in all directions at the same rate. Suppose, however, that we reduce the intensity of the light source illuminating the slits until the source is so weak that only a single photon is present between the slits and the projection screen at any given time. Let us also replace the projection screen by a photographic film that records the position where it is struck by each photon. If we wait a sufficiently long time that a great many photons have passed through the slits and struck the photographic film, and then develop the film, do we see an interference pattern which looks like that shown in Figure [f3]? The answer to this question, as determined by experiment, is that we see exactly the same interference pattern.

According to the previous discussion, the interference pattern is built up one photon at a time. In other words, the pattern is not due to the interaction of different photons. Moreover, the point at which a given photon strikes the film is not influenced by the points at which previous photons struck the film, given that there is only one photon in the apparatus at any given time. Hence, the only way in which the classical interference pattern can be reconstructed, after a great many photons have passed through the apparatus, is if each photon has a greater probability of striking the film at points where the classical interference pattern is bright, and a lesser probability of striking the film at points where the interference pattern is dark.

Suppose, then, that we allow  $N$  photons to pass through our apparatus, and then count the number of photons that strike the recording film between  $y$  and  $y + \Delta y$ , where  $\Delta y$  is a relatively small division. Let us call this number  $n(y)$ . The number of photons that strike a region of the film in a given time interval is equivalent to the intensity of the light illuminating that region of the film multiplied by the area of the region, because each photon carries a fixed amount of energy. Hence, in order to reconcile the classical and quantum viewpoints, we need

$$P_y(y) \equiv \lim_{N \rightarrow \infty} \left[ \frac{n(y)}{N} \right] \propto I(y) \Delta y, \quad (2.8.1)$$

where  $I(y)$  is given in Equation (2.7.5). Here,  $P_y(y)$  is the probability that a given photon strikes the film between  $y$  and  $y + \Delta y$ . Note that  $P_y \propto \Delta y$ . In other words, the probability of a photon striking a region of the film of width  $\Delta y$  is directly proportional to this width. Actually, this is only true as long as  $\Delta y$  is relatively small. It is convenient to define a probability density,  $P(y)$ , which is such that the probability of a photon striking a region of the film of infinitesimal width  $dy$  is  $P_y(y) = P(y) dy$ . Now, Equation (2.8.1) yields  $P_y(y) \propto I(y) dy$ , which gives  $P(y) \propto I(y)$ . However, according to Equation (2.7.4),  $I(y) \propto |\psi(y, t)|^2$ . Thus, we obtain

$$P(y) \propto |\psi(y, t)|^2. \quad (2.8.2)$$

In other words, the probability density of a photon striking a given point on the film is proportional to the modulus squared of the wavefunction at that point. Another way of saying this is that the probability of a measurement of the photon's distance from the centerline, at the location of the film, yielding a result between  $y$  and  $y + dy$  is proportional to  $|\psi(y, t)|^2 dy$ .

Note that, in the quantum-mechanical picture, we can only predict the probability that a given photon strikes a given point on the film. If photons behaved classically then we could, in principle, solve their equations of motion and predict exactly where each photon was going to strike the film, given its initial position and velocity. This loss of determinacy in quantum mechanics is a direct consequence of wave-particle duality. In other words, we can only reconcile the wave-like and particle-like properties of light in a statistical sense. It is impossible to reconcile them on the individual particle level.

In principle, each photon that passes through our apparatus is equally likely to pass through one of the two slits. Can we determine through which slit a given photon passed? Suppose that our original interference experiment involves sending  $N \gg 1$  photons through our apparatus. We know that we get an interference pattern in this experiment. Suppose that we perform a modified interference experiment in which we close off one slit, send  $N/2$  photons through the apparatus, and then open the slit and close off the other slit, and send  $N/2$  photons through the apparatus. In this second experiment, which is virtually identical to the first on the individual photon level, we know exactly which slit each photon passed through. However, the wave theory of light (which we expect to agree with the quantum theory in the limit  $N \gg 1$ ) tells us that our modified interference experiment will not result in the formation of an interference pattern. After all, according to conventional wave theory, it is impossible to obtain a two-slit interference pattern from a single slit. Hence, we conclude that any attempt to measure through which slit each photon passes in our two-slit interference experiment results in the destruction of the interference pattern. It follows that, in the quantum-mechanical



version of the two-slit interference experiment, we must think of each photon as essentially passing through both slits simultaneously.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

2.8: Quantum Interference of Light is shared under a [not declared](#) license and was authored, remixed, and/or curated by LibreTexts.

## 2.9: Particles

### Classical Particles

In this book, we are going to concentrate, almost exclusively, on the behavior of non-relativistic particles of non-zero mass (e.g., electrons). In the absence of external forces, such particles, of mass  $m$ , energy  $E$ , and momentum  $p$ , move classically in a straight-line with velocity

$$v = \frac{p}{m}, \quad (2.9.1)$$

and satisfy

$$E = \frac{p^2}{2m}. \quad (2.9.2)$$

### Quantum Particles

Just as light-waves sometimes exhibit particle-like properties, it turns out that massive particles sometimes exhibit wave-like properties. For instance, it is possible to obtain a double-slit interference pattern from a stream of mono-energetic electrons passing through two closely-spaced narrow slits. The effective wavelength of the electrons can be determined by measuring the width of the light and dark bands in the interference pattern. [See Equation (2.7.6).] It is found that

$$\lambda = \frac{h}{p}. \quad (2.9.3)$$

The same relation is found for other types of particles. The previous wavelength is called the *de Broglie wavelength*, after Louis de Broglie, who first suggested that particles should have wave-like properties in 1923. Note that the de Broglie wavelength is generally very small. For instance, that of an electron is

$$\lambda_e = 1.2 \times 10^{-9} [E(\text{eV})]^{-1/2} \text{ m}, \quad (2.9.4)$$

where the electron energy is conveniently measured in units of electron-volts (eV). (An electron accelerated from rest through a potential difference of 1000 V acquires an energy of 1000 eV, and so on.) The de Broglie wavelength of a proton is

$$\lambda_p = 2.9 \times 10^{-11} [E(\text{eV})]^{-1/2} \text{ m}. \quad (2.9.5)$$

Given the smallness of the de Broglie wavelengths of common particles, it is actually quite difficult to perform particle interference experiments. In general, in order to perform an effective interference experiment, the spacing of the slits must not be too much greater than the wavelength of the wave. Hence, particle interference experiments require either very low-energy particles (because  $\lambda \propto E^{-1/2}$ ), or very closely-spaced slits. Usually the “slits” consist of crystals, which act a bit like diffraction gratings with a characteristic spacing of order the inter-atomic spacing (which is generally about  $10^{-9}$  m).

Equation (2.9.3) can be rearranged to give

$$p = \hbar k, \quad (2.9.6)$$

which is exactly the same as the relation between momentum and wavenumber that we obtained earlier for photons. [See Equation (e2.19b).] For the case of a particle moving the three dimensions, the previous relation generalizes to give

$$\mathbf{p} = \hbar \mathbf{k}, \quad (2.9.7)$$

where  $\mathbf{p}$  is the particle’s vector momentum, and  $\mathbf{k}$  its wavevector. It follows that the momentum of a quantum particle, and, hence, its velocity, is always parallel to its wavevector.

Because the relation (e2.19b) between momentum and wavenumber applies to both photons and massive particles, it seems plausible that the closely-related relation (2.6.1) between energy and wave angular frequency should also apply to both photons and particles. If this is the case, and we can write

$$E = \hbar \omega \quad (2.9.8)$$

for particle waves, then Equations (2.9.2) and (2.9.6) yield the following dispersion relation for such waves:

$$\omega = \frac{\hbar k^2}{2m}. \quad (2.9.9)$$

We saw earlier that a plane-wave propagates at the so-called phase-velocity,

$$v_p = \frac{\omega}{k}. \quad (2.9.10)$$

However, according to the previous dispersion relation, a particle plane-wave propagates at

$$v_p = \frac{p}{2m}. \quad (2.9.11)$$

Note, from Equation (2.9.1), that this is only half of the classical particle velocity. Does this imply that the dispersion relation (2.9.9) is incorrect? Let us investigate further.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

2.9: Particles is shared under a [not declared](#) license and was authored, remixed, and/or curated by LibreTexts.

## 2.10: Wave-Packets

The previous discussion suggests that the wavefunction of a massive particle of momentum  $p$  and energy  $E$ , moving in the positive  $x$ -direction, can be written

$$\psi(x, t) = \bar{\psi} e^{i(kx - \omega t)}, \quad (2.10.1)$$

where  $k = p/\hbar > 0$  and  $\omega = E/\hbar > 0$ . Here,  $\omega$  and  $k$  are linked via the dispersion relation ([e2.38]). Expression ([e2.41]) represents a plane-wave whose maxima and minima propagate in the positive  $x$ -direction with the phase-velocity  $v_p = \omega/k$ . As we have seen, this phase-velocity is only half of the classical velocity of a massive particle.

From before, the most reasonable physical interpretation of the wavefunction is that  $|\psi(x, t)|^2$  is proportional to the probability density of finding the particle at position  $x$  at time  $t$ . However, the modulus squared of the wavefunction ([e2.41]) is  $|\bar{\psi}|^2$ , which depends on neither  $x$  nor  $t$ . In other words, this wavefunction represents a particle that is equally likely to be found anywhere on the  $x$ -axis at all times. Hence, the fact that the maxima and minima of the wavefunction propagate at a phase-velocity that does not correspond to the classical particle velocity does not have any real physical consequences.

How can we write the wavefunction of a particle that is localized in  $x$ : that is, a particle that is more likely to be found at some positions on the  $x$ -axis than at others? It turns out that we can achieve this goal by forming a linear combination of plane-waves of different wavenumbers: in other words,

$$\psi(x, t) = \int_{-\infty}^{\infty} \bar{\psi}(k) e^{i(kx - \omega t)} dk. \quad (2.10.2)$$

Here,  $\bar{\psi}(k)$  represents the complex amplitude of plane-waves of wavenumber  $k$  in this combination. In writing the previous expression, we are relying on the assumption that particle waves are superposable: that is, that it is always possible to add two valid wave solutions to form a third valid wave solution. The ultimate justification for this assumption is that particle waves satisfy a differential wave equation that is linear in  $\psi$ . As we shall see, in Section 1.15, this is indeed the case. Incidentally, a plane-wave that varies as  $\exp[i(kx - \omega t)]$  and has a negative  $k$  (but positive  $\omega$ ) propagates in the negative  $x$ -direction at the phase-velocity  $\omega/|k|$ . Hence, the superposition ([e2.42]) includes both forward and backward propagating waves.

There is a useful mathematical theorem, known as *Fourier's theorem*, which states that if

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \bar{f}(k) e^{ikx} dk, \quad (2.10.3)$$

then

$$\bar{f}(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-ikx} dx. \quad (2.10.4)$$

Here,  $\bar{f}(k)$  is known as the *Fourier transform* of the function  $f(x)$ . We can use Fourier's theorem to find the  $k$ -space function  $\bar{\psi}(k)$  that generates any given  $x$ -space wavefunction  $\psi(x)$  at a given time.

For instance, suppose that at  $t = 0$  the wavefunction of our particle takes the form

$$\psi(x, 0) \propto \exp\left[ik_0 x - \frac{(x - x_0)^2}{4(\Delta x)^2}\right]. \quad (2.10.5)$$

Thus, the initial probability density of the particle is written

$$|\psi(x, 0)|^2 \propto \exp\left[-\frac{(x - x_0)^2}{2(\Delta x)^2}\right]. \quad (2.10.6)$$

This particular probability distribution is called a *Gaussian* distribution, and is plotted in Figure [f4]. It can be seen that a measurement of the particle's position is most likely to yield the value  $x_0$ , and very unlikely to yield a value which differs from  $x_0$  by more than  $3\Delta x$ . Thus, Equation ([e2.45]) is the wavefunction of a particle that is initially localized around  $x = x_0$  in some region whose width is of order  $\Delta x$ . This type of wavefunction is known as a *wave-packet*.




Figure 7: A Gaussian probability distribution in  $x$ -space.

According to Equation ([e2.42]),

$$\psi(x, 0) = \int_{-\infty}^{\infty} \bar{\psi}(k) e^{i k x} dk. \quad (2.10.7)$$

Hence, we can employ Fourier's theorem to invert this expression to give

$$\bar{\psi}(k) \propto \int_{-\infty}^{\infty} \psi(x, 0) e^{-i k x} dx. \quad (2.10.8)$$

Making use of Equation ([e2.45]), we obtain

$$\bar{\psi}(k) \propto e^{-i(k-k_0)x_0} \int_{-\infty}^{\infty} \exp \left[ -i(k-k_0)(x-x_0) - \frac{(x-x_0)^2}{4(\Delta x)^2} \right] dx. \quad (2.10.9)$$

Changing the variable of integration to  $y = (x - x_0)/(2 \Delta x)$ , this reduces to

$$\bar{\psi}(k) \propto e^{-i k x_0} \int_{-\infty}^{\infty} \exp(-i \beta y - y^2) dy, \quad (2.10.10)$$

where  $\beta = 2(k - k_0) \Delta x$ . The previous equation can be rearranged to give

$$\bar{\psi}(k) \propto e^{-i k x_0 - \beta^2/4} \int_{-\infty}^{\infty} e^{-(y-y_0)^2} dy, \quad (2.10.11)$$

where  $y_0 = -i \beta/2$ . The integral now just reduces to a number, as can easily be seen by making the change of variable  $z = y - y_0$ . Hence, we obtain

$$\bar{\psi}(k) \propto \exp \left[ -i k x_0 - \frac{(k - k_0)^2}{4(\Delta k)^2} \right], \quad (2.10.12)$$

where

$$\Delta k = \frac{1}{2 \Delta x}. \quad (2.10.13)$$

If  $|\psi(x)|^2$  is proportional to the probability density of a measurement of the particle's position yielding the value  $x$  then it stands to reason that  $|\bar{\psi}(k)|^2$  is proportional to the probability density of a measurement of the particle's wavenumber yielding the value  $k$ . (Recall that  $p = \hbar k$ , so a measurement of the particle's wavenumber,  $k$ , is equivalent to a measurement of the particle's momentum,  $p$ ). According to Equation ([e2.51]),

$$|\bar{\psi}(k)|^2 \propto \exp \left[ -\frac{(k - k_0)^2}{2(\Delta k)^2} \right]. \quad (2.10.14)$$

Note that this probability distribution is a Gaussian in  $k$ -space. [See Equation ([e2.46]) and Figure [f4].] Hence, a measurement of  $k$  is most likely to yield the value  $k_0$ , and very unlikely to yield a value which differs from  $k_0$  by more than  $3 \Delta k$ . Incidentally, a Gaussian is the only simple mathematical function in  $x$ -space that has the same form as its Fourier transform in  $k$ -space.

We have just seen that a Gaussian probability distribution of characteristic width  $\Delta x$  in  $x$ -space [see Equation ([e2.46])] transforms to a Gaussian probability distribution of characteristic width  $\Delta k$  in  $k$ -space [see Equation ([e2.53])], where

$$\Delta x \Delta k = \frac{1}{2}. \quad (2.10.15)$$

This illustrates an important property of wave-packets. Namely, if we wish to construct a packet that is very localized in  $x$ -space (i.e., if  $\Delta x$  is small) then we need to combine plane-waves with a very wide range of different  $k$ -values (i.e.,  $\Delta k$  will be large). Conversely, if we only combine plane-waves whose wavenumbers differ by a small amount (i.e., if  $\Delta k$  is small) then the resulting wave-packet will be very extended in  $x$ -space (i.e.,  $\Delta x$  will be large).

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

2.10: Wave-Packets is shared under a [not declared](#) license and was authored, remixed, and/or curated by LibreTexts.

## 2.11: Evolution of Wave-Packets

We have seen, in Equation ([e2.45]), how to write the wavefunction of a particle that is initially localized in  $x$ -space. Let us examine how this wavefunction evolves in time. According to Equation ([e2.42]), we have

$$\psi(x, t) = \int_{-\infty}^{\infty} \bar{\psi}(k) e^{i\phi(k)} dk, \quad (2.11.1)$$

where

$$\phi(k) = kx - \omega(k)t. \quad (2.11.2)$$

The function  $\bar{\psi}(k)$  is obtained by Fourier transforming the wavefunction at  $t = 0$ . [See Equations ([e2.42a]) and ([e2.51]).] According to Equation ([e2.53]),  $|\bar{\psi}(k)|$  is strongly peaked around  $k = k_0$ . Thus, it is a reasonable approximation to Taylor expand  $\phi(k)$  about  $k_0$ . Keeping terms up to second order in  $k - k_0$ , we obtain

$$\psi(x, t) \propto \int_{-\infty}^{\infty} \bar{\psi}(k) \exp \left[ i \left\{ \phi_0 + \phi'_0 (k - k_0) + \frac{1}{2} \phi''_0 (k - k_0)^2 \right\} \right], \quad (2.11.3)$$

where

$$\begin{aligned} \phi_0 &= \phi(k_0) = k_0 x - \omega_0 t, \\ \phi'_0 &= \frac{d\phi(k_0)}{dk} = x - v_g t, \\ \phi''_0 &= \frac{d^2\phi(k_0)}{dk^2} = -\alpha t, \end{aligned}$$

with

$$\begin{aligned} \omega_0 &= \omega(k_0), \\ v_g &= \frac{d\omega(k_0)}{dk}, \\ \alpha &= \frac{d^2\omega(k_0)}{dk^2}. \end{aligned}$$

Substituting from Equation ([e2.51]), rearranging, and then changing the variable of integration to  $y = (k - k_0)/(2\Delta k)$ , we get

$$\psi(x, t) \propto e^{i(k_0 x - \omega_0 t)} \int_{-\infty}^{\infty} e^{i\beta_1 y - (1+i\beta_2)y^2} dy, \quad (2.11.4)$$

where

$$\begin{aligned} \beta_1 &= 2\Delta k (x - x_0 - v_g t), \\ \beta_2 &= 2\alpha (\Delta k)^2 t. \end{aligned}$$

Incidentally,  $\Delta k = 1/(2\Delta x)$ , where  $\Delta x$  is the initial width of the wave-packet. The previous expression can be rearranged to give

$$\psi(x, t) \propto e^{i(k_0 x - \omega_0 t) - (1+i\beta_2)\beta^2/4} \int_{-\infty}^{\infty} e^{-(1+i\beta_2)(y-y_0)^2} dy, \quad (2.11.5)$$

where  $y_0 = i\beta/2$  and  $\beta = \beta_1/(1+i\beta_2)$ . Again changing the variable of integration to  $z = (1+i\beta_2)^{1/2}(y-y_0)$ , we get

$$\psi(x, t) \propto (1+i\beta_2)^{-1/2} e^{i(k_0 x - \omega_0 t) - (1+i\beta_2)\beta^2/4} \int_{-\infty}^{\infty} e^{-z^2} dz. \quad (2.11.6)$$

The integral now just reduces to a number. Hence, we obtain

$$\psi(x, t) \propto \frac{\exp \left[ i(k_0 x - \omega_0 t) - (x - x_0 - v_g t)^2 \{1 - i2\alpha(\Delta k)^2 t\} / (4\sigma^2) \right]}{[1 + i2\alpha(\Delta k)^2 t]^{1/2}}, \quad (2.11.7)$$

where

$$\sigma^2(t) = (\Delta x)^2 + \frac{\alpha^2 t^2}{4(\Delta x)^2}. \quad (2.11.8)$$

Note that the previous wavefunction is identical to our original wavefunction ([e2.45]) at  $t = 0$ . This justifies the approximation that we made earlier by Taylor expanding the phase factor  $\phi(k)$  about  $k = k_0$ .

According to Equation ([exxx]), the probability density of our particle as a function of time is written

$$|\psi(x, t)|^2 \propto \sigma^{-1}(t) \exp \left[ -\frac{(x - x_0 - v_g t)^2}{2\sigma^2(t)} \right]. \quad (2.11.9)$$

Hence, the probability distribution is a Gaussian, of characteristic width  $\sigma(t)$ , that peaks at  $x = x_0 + v_g t$ . The most likely position of our particle coincides with the peak of the distribution function. Thus, the particle's most likely position is given by

$$x = x_0 + v_g t. \quad (2.11.10)$$

It can be seen that the particle effectively moves at the uniform velocity

$$v_g = \frac{d\omega}{dk}, \quad (2.11.11)$$

which is known as the *group-velocity*. In other words, a plane-wave travels at the phase-velocity,  $v_p = \omega/k$ , whereas a wave-packet travels at the group-velocity,  $v_g = d\omega/dt$ . It follows from the dispersion relation ([e2.38]) for particle waves that

$$v_g = \frac{p}{m}. \quad (2.11.12)$$

However, it can be seen from Equation ([e2.31]) that this is identical to the classical particle velocity. Hence, the dispersion relation ([e2.38]) turns out to be consistent with classical physics, after all, as soon as we realize that individual particles must be identified with wave-packets rather than plane-waves. In fact, a plane-wave is usually interpreted as a continuous stream of particles propagating in the same direction as the wave.

According to Equation ([e2.70]), the width of our wave-packet grows as time progresses. Indeed, it follows from Equations ([e2.38]) and ([e2.64]) that the characteristic time for a wave-packet of original width  $\Delta x$  to double in spatial extent is

$$t_2 \sim \frac{m(\Delta x)^2}{\hbar}. \quad (2.11.13)$$

For instance, if an electron is originally localized in a region of atomic scale (i.e.,  $\Delta x \sim 10^{-10}$  m) then the doubling time is only about  $10^{-16}$  s. Evidently, particle wave-packets (for freely-moving particles) spread very rapidly.

Note, from the previous analysis, that the rate of spreading of a wave-packet is ultimately governed by the second derivative of  $\omega(k)$  with respect to  $k$ . [See Equations ([e2.64]) and ([e2.70]).] This explains why a functional relationship between  $\omega$  and  $k$  is generally known as a dispersion relation—it governs how fast wave-packets disperse as time progresses. However, for the special case where  $\omega$  is a linear function of  $k$ , the second derivative of  $\omega$  with respect to  $k$  is zero, and, hence, there is no dispersion of wave-packets: that is, wave-packets propagate without changing shape. The dispersion relation ([e2.7]) for light-waves is linear in  $k$ . It follows that light pulses propagate through a vacuum without spreading. Another property of linear dispersion relations is that the phase-velocity,  $v_p = \omega/k$ , and the group-velocity,  $v_g = d\omega/dk$ , are identical. Thus, plane light-waves and light pulses both propagate through a vacuum at the characteristic speed  $c = 3 \times 10^8$  m/s. Of course, the dispersion relation ([e2.38]) for particle waves is not linear in  $k$ . Hence, particle plane-waves and particle wave-packets propagate at different velocities, and particle wave-packets also gradually disperse as time progresses.

## Heisenberg's Uncertainty Principle

According to the analysis contained in the previous two sections, a particle wave-packet that is initially localized in  $x$ -space with characteristic width  $\Delta x$  is also localized in  $k$ -space with characteristic width  $\Delta k = 1/(2\Delta x)$ . However, as time progresses, the width of the wave-packet in  $x$ -space increases, while that of the wave-packet in  $k$ -space stays the same. [After all, our previous analysis obtained  $\psi(x, t)$  from Equation ([e2.56]), but assumed that  $\bar{\psi}(k)$  was given by Equation ([e2.51]) at all times.] Hence, in general, we can say that



$$\Delta x \Delta k \gtrsim \frac{1}{2} \quad (2.11.14)$$

Furthermore, we can think of  $\Delta x$  and  $\Delta k$  as characterizing our uncertainty regarding the values of the particle's position and wavenumber, respectively.

A measurement of a particle's wavenumber,  $k$ , is equivalent to a measurement of its momentum,  $p$ , because  $p = \hbar k$ . Hence, an uncertainty in  $k$  of order  $\Delta k$  translates to an uncertainty in  $p$  of order  $\Delta p = \hbar \Delta k$ . It follows from the previous inequality that

$$\Delta x \Delta p \gtrsim \frac{\hbar}{2} \quad (2.11.15)$$

This is the famous *Heisenberg uncertainty principle*, first proposed by Werner Heisenberg in 1927. According to this principle, it is impossible to simultaneously measure the position and momentum of a particle (exactly). Indeed, a good knowledge of the particle's position implies a poor knowledge of its momentum, and vice versa. Note that the uncertainty principle is a direct consequence of representing particles as waves.

It can be seen from Equations ([e2.38]), ([e2.64]), and ([e2.70]) that, at large  $t$ , a particle wavefunction of original width  $\Delta x$  (at  $t = 0$ ) spreads out such that its spatial extent becomes

$$\sigma \sim \frac{\hbar t}{m \Delta x}. \quad (2.11.16)$$

It is easily demonstrated that this spreading is a consequence of the uncertainty principle. Because the initial uncertainty in the particle's position is  $\Delta x$ , it follows that the uncertainty in its momentum is of order  $\hbar/\Delta x$ . This translates to an uncertainty in velocity of  $\Delta v = \hbar/(m \Delta x)$ . Thus, if we imagine that parts of the wavefunction propagate at  $v_0 + \Delta v/2$ , and others at  $v_0 - \Delta v/2$ , where  $v_0$  is the mean propagation velocity, then the wavefunction will spread as time progresses. Indeed, at large  $t$ , we expect the width of the wavefunction to be

$$\sigma \sim \Delta v t \sim \frac{\hbar t}{m \Delta x}, \quad (2.11.17)$$

which is identical to Equation ([espread]). Evidently, the spreading of a particle wavefunction must be interpreted as an increase in our uncertainty regarding the particle's position, rather than an increase in the spatial extent of the particle itself.




Figure 8: *Heisenberg's microscope*.

Figure [fh] illustrates a famous thought experiment known as *Heisenberg's microscope*. Suppose that we try to image an electron using a simple optical system in which the objective lens is of diameter  $D$  and focal-length  $f$ . (In practice, this would only be possible using extremely short-wavelength light.) It is a well-known result in optics that such a system has a minimum angular resolving power of  $\lambda/D$ , where  $\lambda$  is the wavelength of the light illuminating the electron. If the electron is placed at the focus of the lens, which is where the minimum resolving power is achieved, then this translates to a uncertainty in the electron's transverse position of

$$\Delta x \simeq f \frac{\lambda}{D}. \quad (2.11.18)$$

However,

$$\tan \alpha = \frac{D/2}{f}, \quad (2.11.19)$$

where  $\alpha$  is the half-angle subtended by the lens at the electron. Assuming that  $\alpha$  is small, we can write

$$\alpha \simeq \frac{D}{2f}, \quad (2.11.20)$$

so

$$\Delta x \simeq \frac{\lambda}{2\alpha}. \quad (2.11.21)$$

It follows that we can reduce the uncertainty in the electron's position by minimizing the ratio  $\lambda/\alpha$ : that is, by employing short-wavelength radiation, and a wide-angle lens.

Let us now examine Heisenberg's microscope from a quantum-mechanical point of view. According to quantum mechanics, the electron is imaged when it scatters an incoming photon towards the objective lens. Let the wavevector of the incoming photon have the  $(x, y)$  components  $(k, 0)$ . See Figure [fh]. If the scattered photon subtends an angle  $\theta$  with the center-line of the optical system, as shown in the figure, then its wavevector is written  $(k \sin \theta, k \cos \theta)$ . Here, we are ignoring any shift in wavelength of the photon on scattering—in other words, the magnitude of the  $\mathbf{k}$ -vector is assumed to be the same before and after scattering. Thus, the change in the  $x$ -component of the photon's wavevector is  $\Delta k_x = k(\sin \theta - 1)$ . This translates to a change in the photon's  $x$ -component of momentum of  $\Delta p_x = \hbar k(\sin \theta - 1)$ . By momentum conservation, the electron's  $x$ -momentum will change by an equal and opposite amount. However,  $\theta$  can range all the way from  $-\alpha$  to  $+\alpha$ , and the scattered photon will still be collected by the imaging system. It follows that the uncertainty in the electron's momentum is

$$\Delta p \simeq 2 \hbar k \sin \alpha \simeq \frac{4\pi \hbar \alpha}{\lambda}. \quad (2.11.22)$$

Note that in order to reduce the uncertainty in the momentum we need to maximize the ratio  $\lambda/\alpha$ . This is exactly the opposite of what we need to do to reduce the uncertainty in the position. Multiplying the previous two equations, we obtain

$$\Delta x \Delta p \sim h, \quad (2.11.23)$$

which is essentially the uncertainty principle.

According to Heisenberg's microscope, the uncertainty principle follows from two facts. First, it is impossible to measure any property of a microscopic dynamical system without disturbing the system somewhat. Second, particle and light energy and momentum are quantized. Hence, there is a limit to how small we can make the aforementioned disturbance. Thus, there is an irreducible uncertainty in certain measurements that is a consequence of the act of measurement itself.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

2.11: Evolution of Wave-Packets is shared under a [not declared](#) license and was authored, remixed, and/or curated by LibreTexts.

## 2.12: Schrodinger's Equation and Wavefunction Collapse

We have seen that the wavefunction of a free particle of mass  $m$  satisfies

$$\psi(x, t) = \int_{-\infty}^{\infty} \bar{\psi}(k) e^{i(kx - \omega t)} dk, \quad (2.12.1)$$

where  $\bar{\psi}(k)$  is determined by  $\psi(x, 0)$ , and

$$\omega(k) = \frac{\hbar k^2}{2m}. \quad (2.12.2)$$

It follows from Equation ([e2.78]) that

$$\frac{\partial \psi}{\partial x} = \int_{-\infty}^{\infty} (ik) \bar{\psi}(k) e^{i(kx - \omega t)} dk, \quad (2.12.3)$$

and

$$\frac{\partial^2 \psi}{\partial x^2} = \int_{-\infty}^{\infty} (-k^2) \bar{\psi}(k) e^{i(kx - \omega t)} dk, \quad (2.12.4)$$

whereas

$$\frac{\partial \psi}{\partial t} = \int_{-\infty}^{\infty} (-i\omega) \bar{\psi}(k) e^{i(kx - \omega t)} dk. \quad (2.12.5)$$

Thus,

$$i \frac{\partial \psi}{\partial t} + \frac{\hbar}{2m} \frac{\partial^2 \psi}{\partial x^2} = \int_{-\infty}^{\infty} \left( \omega - \frac{\hbar k^2}{2m} \right) \bar{\psi}(k) e^{i(kx - \omega t)} dk = 0, \quad (2.12.6)$$

where use has been made of the dispersion relation ([e2.79]). Multiplying through by  $\hbar$ , we obtain

$$i \hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2}. \quad (2.12.7)$$

This expression is known as *Schrödinger's equation*, because it was first introduced by Erwin Schrödinger in 1926. Schrödinger's equation is a linear, second-order, partial differential equation that governs the time evolution of a particle wavefunction, and is generally easier to solve than the integral equation ([e2.78]).

Of course, Equation ([e2.84]) is only applicable to freely-moving particles. Fortunately, it is fairly easy to guess the generalization of this equation for particles moving in some potential  $V(x)$ . It is plausible, from Equation ([e2.80]), that we can identify  $k$  with the differential operator  $-i\partial/\partial x$ . Hence, the differential operator on the right-hand side of Equation ([e2.84]) is equivalent to  $\hbar^2 k^2/(2m)$ . But,  $p = \hbar k$ . Thus, the operator is also equivalent to  $p^2/(2m)$ , which is just the energy of a freely-moving particle. However, in the presence of a potential  $V(x)$ , the particle's energy is written  $p^2/(2m) + V$ . Thus, it seems reasonable to make the substitution

$$-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} \rightarrow -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + V(x). \quad (2.12.8)$$

This leads to the general (one-dimensional) form of Schrödinger's equation:

$$i \hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2} + V(x) \psi. \quad (2.12.9)$$

### Wavefunction Collapse

Consider an extended wavefunction,  $\psi(x, t)$ . According to our usual interpretation,  $|\psi(x, t)|^2$  is proportional to the probability density of a measurement of the particle's position yielding the value  $x$  at time  $t$ . If the wavefunction is extended then there is a wide range of likely values that this measurement could give. Suppose that we make such a measurement, and obtain the value  $x_0$ . We now know that the particle is located at  $x = x_0$ . If we make another measurement immediately after the first one then common

sense tells us that we must obtain the same value,  $x_0$ , because the particle cannot have shifted position appreciably in an infinitesimal time interval. Thus, immediately after the first measurement, a measurement of the particle's position is certain to give the value  $x_0$ , and has no chance of giving any other value. This implies that the wavefunction must have collapsed to some sort of "spike" function located at  $x = x_0$ . This is illustrated in Figure [coll]. Of course, as soon as the wavefunction has collapsed, it starts to expand again, as discussed in Section 1.13. Thus, the second measurement must be made reasonably quickly after the first, in order to guarantee that the same result will be obtained.




Figure 9: Collapse of the wavefunction upon measurement of  $x$ .

The previous discussion illustrates an important point in quantum mechanics. Namely, that the wavefunction of a particle changes discontinuously (in time) whenever a measurement is made. We conclude that there are two types of time evolution of the wavefunction in quantum mechanics. First, there is a smooth evolution that is governed by Schrödinger's equation. This evolution takes place between measurements. Second, there is a discontinuous evolution that takes place each time a measurement is made.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

2.12: Schrodinger's Equation and Wavefunction Collapse is shared under a [not declared](#) license and was authored, remixed, and/or curated by LibreTexts.

## 2.13: Exercises

1. A He-Ne laser emits radiation of wavelength  $\lambda = 633 \text{ nm}$ . How many photons are emitted per second by a laser with a power of 1 mW? What force does such laser exert on a body which completely absorbs its radiation?
2. The ionization energy of a hydrogen atom in its ground state is  $E_{\text{ion}} = 13.60 \text{ eV}$  (1 eV is the energy acquired by an electron accelerated through a potential difference of 1 V). Calculate the frequency, wavelength, and wavenumber of the electromagnetic radiation that will just ionize the atom.
3. The maximum energy of photoelectrons from aluminium is 2.3 eV for radiation of wavelength 2000 Angstrom, and 0.90 eV for radiation of wavelength 2580 Angstrom. Use this data to calculate Planck's constant, as well as the work function of aluminium.
4. Show that the de Broglie wavelength of an electron accelerated from rest across a potential difference  $V$  is given by

$$\lambda = 1.23 \times 10^{-9} V^{-1/2} \text{ m}, \quad (2.13.1)$$

where  $V$  is measured in volts.

5. If the atoms in a regular crystal are separated by  $3 \times 10^{-10} \text{ m}$  demonstrate that an accelerating voltage of about 1.5 kV would be required to produce an electron diffraction pattern from the crystal.
6. The relationship between wavelength and frequency for electromagnetic waves in a waveguide is

$$\lambda = \frac{c}{\sqrt{\nu^2 - \nu_0^2}}, \quad (2.13.2)$$

where  $c$  is the velocity of light in vacuum. What are the group- and phase-velocities of such waves as functions of  $\nu_0$  and  $\lambda$ ?

7. Nuclei, typically of size  $10^{-14} \text{ m}$ , frequently emit electrons with energies of 1–10 MeV. Use the uncertainty principle to show that electrons of energy 1 MeV could not be contained in the nucleus before the decay.
8. A particle of mass  $m$  has a wavefunction

$$\psi(x, t) = A \exp \left[ -a \left( \frac{m x^2}{\hbar} + i t \right) \right], \quad (2.13.3)$$

where  $A$  and  $a$  are positive real constants. For what potential function  $V(x)$  does  $\psi$  satisfy the Schrödinger equation?

1. Plural of *quantum*: Latin neuter of *quantus*: how much?↩

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

2.13: Exercises is shared under a [not declared](#) license and was authored, remixed, and/or curated by LibreTexts.

## CHAPTER OVERVIEW

### 3: Fundamentals of Quantum Mechanics

The previous chapter serves as a useful introduction to many of the basic concepts of quantum mechanics. In this chapter, we shall examine these concepts in a more systematic fashion. For the sake of simplicity, we shall concentrate on one-dimensional systems.

- 3.1: Schrodinger's Equation
- 3.2: Normalization of the Wavefunction
- 3.3: Expectation Values (Averages) and Variances
- 3.4: Ehrenfest's Theorem
- 3.5: Operators
- 3.6: Momentum Representation
- 3.7: Heisenberg's Uncertainty Principle
- 3.8: Eigenstates and Eigenvalues
- 3.9: Measurement
- 3.10: Stationary States
- 3.11: Exercises

---

This page titled [3: Fundamentals of Quantum Mechanics](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by Richard Fitzpatrick.

### 3.1: Schrodinger's Equation

---

Consider a dynamical system consisting of a single non-relativistic particle of mass  $m$  moving along the  $x$ -axis in some real potential  $V(x)$ . In quantum mechanics, the instantaneous state of the system is represented by a complex wavefunction  $\psi(x, t)$ . This wavefunction evolves in time according to Schrödinger's equation:

$$i \hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2} + V(x) \psi. \quad (3.1.1)$$

The wavefunction is interpreted as follows:  $|\psi(x, t)|^2$  is the probability density of a measurement of the particle's displacement yielding the value  $x$ . Thus, the probability of a measurement of the displacement giving a result between  $a$  and  $b$  (where  $a < b$ ) is

$$P_{x \in a:b}(t) = \int_a^b |\psi(x, t)|^2 dx. \quad (3.1.2)$$

Note that this quantity is real and positive definite.

---

This page titled [3.1: Schrodinger's Equation](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 3.2: Normalization of the Wavefunction

Now, a probability is a real number lying between 0 and 1. An outcome of a measurement that has a probability 0 is an impossible outcome, whereas an outcome that has a probability 1 is a certain outcome. According to Equation ([e3.2]), the probability of a measurement of  $x$  yielding a result lying between  $-\infty$  and  $+\infty$  is

$$P_{x \in -\infty:\infty}(t) = \int_{-\infty}^{\infty} |\psi(x, t)|^2 dx. \quad (3.2.1)$$

However, a measurement of  $x$  must yield a value lying between  $-\infty$  and  $+\infty$ , because the particle has to be located somewhere. It follows that  $P_{x \in -\infty:\infty} = 1$ , or

$$\int_{-\infty}^{\infty} |\psi(x, t)|^2 dx = 1, \quad (3.2.2)$$

which is generally known as the *normalization condition* for the wavefunction.

For example, suppose that we wish to normalize the wavefunction of a Gaussian wave-packet, centered on  $x = x_0$ , and of characteristic width  $\sigma$  (see Section [s2.9]): that is,

$$\psi(x) = \psi_0 e^{-(x-x_0)^2/(4\sigma^2)}. \quad (3.2.3)$$

In order to determine the normalization constant  $\psi_0$ , we simply substitute Equation ([e3.5]) into Equation ([e3.4]) to obtain

$$|\psi_0|^2 \int_{-\infty}^{\infty} e^{-(x-x_0)^2/(2\sigma^2)} dx = 1. \quad (3.2.4)$$

Changing the variable of integration to  $y = (x - x_0)/(\sqrt{2}\sigma)$ , we get

$$|\psi_0|^2 \sqrt{2}\sigma \int_{-\infty}^{\infty} e^{-y^2} dy = 1. \quad (3.2.5)$$

However,

$$\int_{-\infty}^{\infty} e^{-y^2} dy = \sqrt{\pi}, \quad (3.2.6)$$

which implies that

$$|\psi_0|^2 = \frac{1}{(2\pi\sigma^2)^{1/2}}. \quad (3.2.7)$$

Hence, a general normalized Gaussian wavefunction takes the form

$$\psi(x) = \frac{e^{i\varphi}}{(2\pi\sigma^2)^{1/4}} e^{-(x-x_0)^2/(4\sigma^2)}, \quad (3.2.8)$$

where  $\varphi$  is an arbitrary real phase-angle.

It is important to demonstrate that if a wavefunction is initially normalized then it stays normalized as it evolves in time according to Schrödinger's equation. If this is not the case then the probability interpretation of the wavefunction is untenable, because it does not make sense for the probability that a measurement of  $x$  yields any possible outcome (which is, manifestly, unity) to change in time. Hence, we require that

$$\frac{d}{dt} \int_{-\infty}^{\infty} |\psi(x, t)|^2 dx = 0, \quad (3.2.9)$$

for wavefunctions satisfying Schrödinger's equation. The previous equation gives

$$\frac{d}{dt} \int_{-\infty}^{\infty} \psi^* \psi dx = \int_{-\infty}^{\infty} \left( \frac{\partial \psi^*}{\partial t} \psi + \psi^* \frac{\partial \psi}{\partial t} \right) dx = 0. \quad (3.2.10)$$

Now, multiplying Schrödinger's equation by  $\psi^*/(i\hbar)$ , we obtain



$$\psi^* \frac{\partial \psi}{\partial t} = \frac{i \hbar}{2m} \psi^* \frac{\partial^2 \psi}{\partial x^2} - \frac{i}{\hbar} V |\psi|^2. \quad (3.2.11)$$

The complex conjugate of this expression yields

$$\psi \frac{\partial \psi^*}{\partial t} = -\frac{i \hbar}{2m} \psi \frac{\partial^2 \psi^*}{\partial x^2} + \frac{i}{\hbar} V |\psi|^2 \quad (3.2.12)$$

[because  $(AB)^* = A^* B^*$ ,  $A^{**} = A$ , and  $i^* = -i$ ].

Summing the previous two equations, we get

$$\frac{\partial \psi^*}{\partial t} \psi + \psi^* \frac{\partial \psi}{\partial t} = \frac{i \hbar}{2m} \left( \psi^* \frac{\partial^2 \psi}{\partial x^2} - \psi \frac{\partial^2 \psi^*}{\partial x^2} \right) = \frac{i \hbar}{2m} \frac{\partial}{\partial x} \left( \psi^* \frac{\partial \psi}{\partial x} - \psi \frac{\partial \psi^*}{\partial x} \right). \quad (3.2.13)$$

Equations (3.2.12) and (3.2.13) can be combined to produce

$$\frac{d}{dt} \int_{-\infty}^{\infty} |\psi|^2 dx = \frac{i \hbar}{2m} \left[ \psi^* \frac{\partial \psi}{\partial x} - \psi \frac{\partial \psi^*}{\partial x} \right]_{-\infty}^{\infty} = 0. \quad (3.2.14)$$

The previous equation is satisfied provided

$$|\psi| \rightarrow 0 \quad \text{as} \quad |x| \rightarrow \infty. \quad (3.2.15)$$

However, this is a necessary condition for the integral on the left-hand side of Equation (3.2.14) to converge. Hence, we conclude that all wavefunctions that are *square-integrable* [i.e., are such that the integral in Equation (3.2.14) converges] have the property that if the normalization condition (3.2.14) is satisfied at one instant in time then it is satisfied at all subsequent times.

It is also possible to demonstrate, via very similar analysis to that just described, that

$$\frac{dP_{x \in a:b}}{dt} + j(b, t) - j(a, t) = 0, \quad (3.2.16)$$

where  $P_{x \in a:b}$  is defined in Equation (3.2.2), and

$$j(x, t) = \frac{i \hbar}{2m} \left( \psi \frac{\partial \psi^*}{\partial x} - \psi^* \frac{\partial \psi}{\partial x} \right) \quad (3.2.17)$$

is known as the *probability current*. Note that  $j$  is real. Equation (3.2.16) is a *probability conservation equation*. According to this equation, the probability of a measurement of  $x$  lying in the interval  $a$  to  $b$  evolves in time due to the difference between the flux of probability into the interval [i.e.,  $j(a, t)$ ], and that out of the interval [i.e.,  $j(b, t)$ ]. Here, we are interpreting  $j(x, t)$  as the flux of probability in the  $+x$ -direction at position  $x$  and time  $t$ .

Note, finally, that not all wavefunctions can be normalized according to the scheme set out in Equation (3.2.14). For instance, a plane-wave wavefunction

$$\psi(x, t) = \psi_0 e^{i(kx - \omega t)} \quad (3.2.18)$$

is not square-integrable, and, thus, cannot be normalized. For such wavefunctions, the best we can say is that

$$P_{x \in a:b}(t) \propto \int_a^b |\psi(x, t)|^2 dx. \quad (3.2.19)$$

In the following, all wavefunctions are assumed to be square-integrable and normalized, unless otherwise stated.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled 3.2: Normalization of the Wavefunction is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

### 3.3: Expectation Values (Averages) and Variances

We have seen that  $|\psi(x, t)|^2$  is the probability density of a measurement of a particle's displacement yielding the value  $x$  at time  $t$ . Suppose that we make a large number of independent measurements of the displacement on an equally large number of identical quantum systems. In general, measurements made on different systems will yield different results. However, from the definition of probability (see Chapter [s2]), the mean of all these results is simply

$$\langle x \rangle = \int_{-\infty}^{\infty} x |\psi|^2 dx. \quad (3.3.1)$$

Here,  $\langle x \rangle$  is called the *expectation value* of  $x$ . (See Chapter [s2].) Similarly the expectation value of any function of  $x$  is

$$\langle f(x) \rangle = \int_{-\infty}^{\infty} f(x) |\psi|^2 dx. \quad (3.3.2)$$

In general, the results of the various different measurements of  $x$  will be scattered around the expectation value,  $\langle x \rangle$ . The degree of scatter is parameterized by the quantity

$$\sigma_x^2 = \int_{-\infty}^{\infty} (x - \langle x \rangle)^2 |\psi|^2 dx \equiv \langle x^2 \rangle - \langle x \rangle^2, \quad (3.3.3)$$

which is known as the *variance* of  $x$ . (See Chapter [s2].) The square-root of this quantity,  $\sigma_x$ , is called the *standard deviation* of  $x$ . (See Chapter [s2].) We generally expect the results of measurements of  $x$  to lie within a few standard deviations of the expectation value.

For instance, consider the normalized Gaussian wave-packet [see Equation ([eng])]

$$\psi(x) = \frac{e^{i\varphi}}{(2\pi\sigma^2)^{1/4}} e^{-(x-x_0)^2/(4\sigma^2)}. \quad (3.3.4)$$

The expectation value of  $x$  associated with this wavefunction is

$$\langle x \rangle = \frac{1}{\sqrt{2\pi}\sigma^2} \int_{-\infty}^{\infty} x e^{-(x-x_0)^2/(2\sigma^2)} dx. \quad (3.3.5)$$

Let  $y = (x - x_0)/(\sqrt{2}\sigma)$ . It follows that

$$\langle x \rangle = \frac{x_0}{\sqrt{\pi}} \int_{-\infty}^{\infty} e^{-y^2} dy + \frac{\sqrt{2}\sigma}{\sqrt{\pi}} \int_{-\infty}^{\infty} y e^{-y^2} dy. \quad (3.3.6)$$

However, the second integral on the right-hand side is zero, by symmetry. Hence, making use of Equation ([e3.8]), we obtain

$$\langle x \rangle = x_0. \quad (3.3.7)$$

Evidently, the expectation value of  $x$  for a Gaussian wave-packet is equal to the most likely value of  $x$  (i.e., the value of  $x$  that maximizes  $|\psi|^2$ ).

The variance of  $x$  associated with the Gaussian wave-packet ([e3.24]) is

$$\sigma_x^2 = \frac{1}{\sqrt{2\pi}\sigma^2} \int_{-\infty}^{\infty} (x - x_0)^2 e^{-(x-x_0)^2/(2\sigma^2)} dx. \quad (3.3.8)$$

Let  $y = (x - x_0)/(\sqrt{2}\sigma)$ . It follows that

$$\sigma_x^2 = \frac{2\sigma^2}{\sqrt{\pi}} \int_{-\infty}^{\infty} y^2 e^{-y^2} dy. \quad (3.3.9)$$

However,

$$\int_{-\infty}^{\infty} y^2 e^{-y^2} dy = \frac{\sqrt{\pi}}{2}, \quad (3.3.10)$$

giving

$$\sigma_x^2 = \sigma^2. \quad (3.3.11)$$

This result is consistent with our earlier interpretation of  $\sigma$  as a measure of the spatial extent of the wave-packet. (See Section [\[s2.9\]](#).) It follows that we can rewrite the Gaussian wave-packet ([\[e3.24\]](#)) in the convenient form

$$\psi(x) = \frac{e^{i\varphi}}{(2\pi\sigma_x^2)^{1/4}} e^{-(x-\langle x \rangle)^2/(4\sigma_x^2)}. \quad (3.3.12)$$

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [3.3: Expectation Values \(Averages\) and Variances](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

### 3.4: Ehrenfest's Theorem

A simple way to calculate the expectation value of momentum is to evaluate the time derivative of  $\langle x \rangle$ , and then multiply by the mass  $m$ : that is,

$$\langle p \rangle = m \frac{d\langle x \rangle}{dt} = m \frac{d}{dt} \int_{-\infty}^{\infty} x |\psi|^2 dx = m \int_{-\infty}^{\infty} x \frac{\partial |\psi|^2}{\partial t} dx. \quad (3.4.1)$$

However, it is easily demonstrated that

$$\frac{\partial |\psi|^2}{\partial t} + \frac{\partial j}{\partial x} = 0 \quad (3.4.2)$$

[this is just the differential form of Equation ([\[epc\]](#))], where  $j$  is the probability current defined in Equation ([\[eprobcl\]](#)). Thus,

$$\langle p \rangle = -m \int_{-\infty}^{\infty} x \frac{\partial j}{\partial x} dx = m \int_{-\infty}^{\infty} j dx, \quad (3.4.3)$$

where we have integrated by parts. It follows from Equation ([\[eprobcl\]](#)) that

$$\langle p \rangle = -\frac{i\hbar}{2} \int_{-\infty}^{\infty} \left( \psi^* \frac{\partial \psi}{\partial x} - \frac{\partial \psi^*}{\partial x} \psi \right) dx = -i\hbar \int_{-\infty}^{\infty} \psi^* \frac{\partial \psi}{\partial x} dx, \quad (3.4.4)$$

where we have again integrated by parts. Hence, the expectation value of the momentum can be written

$$\langle p \rangle = m \frac{d\langle x \rangle}{dt} = -i\hbar \int_{-\infty}^{\infty} \psi^* \frac{\partial \psi}{\partial x} dx. \quad (3.4.5)$$

It follows from the previous equation that

$$\frac{d\langle p \rangle}{dt} = -i\hbar \int_{-\infty}^{\infty} \left( \frac{\partial \psi^*}{\partial t} \frac{\partial \psi}{\partial x} + \psi^* \frac{\partial^2 \psi}{\partial t \partial x} \right) dx = \int_{-\infty}^{\infty} \left[ \left( i\hbar \frac{\partial \psi}{\partial t} \right)^* \frac{\partial \psi}{\partial x} + \frac{\partial \psi^*}{\partial x} \left( i\hbar \frac{\partial \psi}{\partial t} \right) \right] dx,$$

where we have integrated by parts. Substituting from Schrödinger's equation ([\[e3.1\]](#)), and simplifying, we obtain

$$\frac{d\langle p \rangle}{dt} = \int_{-\infty}^{\infty} \left[ -\frac{\hbar^2}{2m} \frac{\partial}{\partial x} \left( \frac{\partial \psi^*}{\partial x} \frac{\partial \psi}{\partial x} \right) + V(x) \frac{\partial |\psi|^2}{\partial x} \right] dx = \int_{-\infty}^{\infty} V(x) \frac{\partial |\psi|^2}{\partial x} dx. \quad (3.4.6)$$

Integration by parts yields

$$\frac{d\langle p \rangle}{dt} = - \int_{-\infty}^{\infty} \frac{dV}{dx} |\psi|^2 dx = - \left\langle \frac{dV}{dx} \right\rangle. \quad (3.4.7)$$

Hence, according to Equations ([\[e4.34x\]](#)) and ([\[e3.41\]](#)),

$$\begin{aligned} m \frac{d\langle x \rangle}{dt} &= \langle p \rangle, \\ \frac{d\langle p \rangle}{dt} &= - \left\langle \frac{dV}{dx} \right\rangle. \end{aligned}$$

Evidently, the expectation values of displacement and momentum obey time evolution equations that are analogous to those of classical mechanics. This result is known as *Ehrenfest's theorem*.

Suppose that the potential  $V(x)$  is slowly varying. In this case, we can expand  $dV/dx$  as a Taylor series about  $\langle x \rangle$ . Keeping terms up to second order, we obtain

$$\frac{dV(x)}{dx} = \frac{dV(\langle x \rangle)}{d\langle x \rangle} + \frac{d^2 V(\langle x \rangle)}{d\langle x \rangle^2} (x - \langle x \rangle) + \frac{1}{2} \frac{d^3 V(\langle x \rangle)}{d\langle x \rangle^3} (x - \langle x \rangle)^2. \quad (3.4.8)$$

Substitution of the previous expansion into Equation ([\[e3.43\]](#)) yields

$$\frac{d\langle p \rangle}{dt} = -\frac{dV(\langle x \rangle)}{d\langle x \rangle} - \frac{\sigma_x^2}{2} \frac{d^3V(\langle x \rangle)}{d\langle x \rangle^3}, \quad (3.4.9)$$

because  $\langle 1 \rangle = 1$ , and  $\langle x - \langle x \rangle \rangle = 0$ , and  $\langle (x - \langle x \rangle)^2 \rangle = \sigma_x^2$ . The final term on the right-hand side of the previous equation can be neglected when the spatial extent of the particle wavefunction,  $\sigma_x$ , is much smaller than the variation length-scale of the potential. In this case, Equations ([e3.42]) and ([e3.43]) reduce to

$$m \frac{d\langle x \rangle}{dt} = \langle p \rangle,$$

$$\frac{d\langle p \rangle}{dt} = -\frac{dV(\langle x \rangle)}{d\langle x \rangle}.$$

These equations are exactly equivalent to the equations of classical mechanics, with  $\langle x \rangle$  playing the role of the particle displacement. Of course, if the spatial extent of the wavefunction is negligible then a measurement of  $x$  is almost certain to yield a result that lies very close to  $\langle x \rangle$ . Hence, we conclude that quantum mechanics corresponds to classical mechanics in the limit that the spatial extent of the wavefunction (which is typically of order the de Broglie wavelength) is negligible. This is an important result, because we know that classical mechanics gives the correct answer in this limit.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [3.4: Ehrenfest's Theorem](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 3.5: Operators

An operator,  $O$  (say), is a mathematical entity that transforms one function into another: that is,

$$O(f(x)) \rightarrow g(x). \quad (3.5.1)$$

For instance,  $x$  is an operator, because  $x f(x)$  is a different function to  $f(x)$ , and is fully specified once  $f(x)$  is given. Furthermore,  $d/dx$  is also an operator, because  $df(x)/dx$  is a different function to  $f(x)$ , and is fully specified once  $f(x)$  is given. Now,

$$x \frac{df}{dx} \neq \frac{d}{dx}(x f). \quad (3.5.2)$$

This can also be written

$$x \frac{d}{dx} \neq \frac{d}{dx} x, \quad (3.5.3)$$

where the operators are assumed to act on everything to their right, and a final  $f(x)$  is understood [where  $f(x)$  is a general function]. The previous expression illustrates an important point. Namely, in general, operators do not commute with one another. Of course, some operators do commute. For instance,

$$x x^2 = x^2 x. \quad (3.5.4)$$

Finally, an operator,  $O$ , is termed linear if

$$O(c f(x)) = c O(f(x)), \quad (3.5.5)$$

where  $f$  is a general function, and  $c$  a general complex number. All of the operators employed in quantum mechanics are linear.

Now, from Equations ([e3.22]) and ([e3.38]),

$$\begin{aligned} \langle x \rangle &= \int_{-\infty}^{\infty} \psi^* x \psi dx, \\ \langle p \rangle &= \int_{-\infty}^{\infty} \psi^* \left( -i \hbar \frac{\partial}{\partial x} \right) \psi dx. \end{aligned}$$

These expressions suggest a number of things. First, classical dynamical variables, such as  $x$  and  $p$ , are represented in quantum mechanics by linear operators that act on the wavefunction. Second, displacement is represented by the algebraic operator  $x$ , and momentum by the differential operator  $-i \hbar \partial/\partial x$ : that is,  $p \equiv -i \hbar \partial/\partial x$ .

Finally, the expectation value of some dynamical variable represented by the operator  $O(x)$  is simply

$$\langle O \rangle = \int_{-\infty}^{\infty} \psi^*(x, t) O(x) \psi(x, t) dx. \quad (3.5.6)$$

Clearly, if an operator is to represent a dynamical variable that has physical significance then its expectation value must be real. In other words, if the operator  $O$  represents a physical variable then we require that  $\langle O \rangle = \langle O \rangle^*$ , or

$$\int_{-\infty}^{\infty} \psi^* (O \psi) dx = \int_{-\infty}^{\infty} (O \psi)^* \psi dx, \quad (3.5.7)$$

where  $O^*$  is the complex conjugate of  $O$ . An operator that satisfies the previous constraint is called an *Hermitian* operator. It is easily demonstrated that  $x$  and  $p$  are both Hermitian. The *Hermitian conjugate*,  $O^\dagger$ , of a general operator,  $O$ , is defined as follows:

$$\int_{-\infty}^{\infty} \psi^* (O \psi) dx = \int_{-\infty}^{\infty} (O^\dagger \psi)^* \psi dx. \quad (3.5.8)$$

The Hermitian conjugate of an Hermitian operator is the same as the operator itself: that is,  $p^\dagger = p$ . For a non-Hermitian operator,  $O$  (say), it is easily demonstrated that  $(O^\dagger)^\dagger = O$ , and that the operator  $O + O^\dagger$  is Hermitian. Finally, if  $A$  and  $B$  are two operators, then  $(AB)^\dagger = B^\dagger A^\dagger$ .

Suppose that we wish to find the operator that corresponds to the classical dynamical variable  $x p$ . In classical mechanics, there is no difference between  $x p$  and  $p x$ . However, in quantum mechanics, we have already seen that  $x p \neq p x$ . So, should we choose

$x p$  or  $p x$ ? Actually, neither of these combinations is Hermitian. However,  $(1/2) [x p + (x p)^\dagger]$  is Hermitian. Moreover,  $(1/2) [x p + (x p)^\dagger] = (1/2) (x p + p^\dagger x^\dagger) = (1/2) (x p + p x)$ , which neatly resolves our problem of the order in which to place  $x$  and  $p$ .

It is a reasonable guess that the operator corresponding to energy (which is called the Hamiltonian, and conventionally denoted  $H$ ) takes the form

$$H \equiv \frac{p^2}{2m} + V(x). \quad (3.5.9)$$

Note that  $H$  is Hermitian. Now, it follows from Equation ([e3.54]) that

$$H \equiv -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + V(x). \quad (3.5.10)$$

However, according to Schrödinger's equation, ([e3.1]), we have

$$-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + V(x) = i \hbar \frac{\partial}{\partial t}, \quad (3.5.11)$$

so

$$H \equiv i \hbar \frac{\partial}{\partial t}. \quad (3.5.12)$$

Thus, the time-dependent Schrödinger equation can be written

$$i \hbar \frac{\partial \psi}{\partial t} = H \psi. \quad (3.5.13)$$

Finally, if  $O(x, p, E)$  is a classical dynamical variable that is a function of displacement, momentum, and energy then a reasonable guess for the corresponding operator in quantum mechanics is  $(1/2) [O(x, p, H) + O^\dagger(x, p, H)]$ , where  $p = -i \hbar \partial / \partial x$ , and  $H = i \hbar \partial / \partial t$ .

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [3.5: Operators](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

### 3.6: Momentum Representation

Fourier's theorem (see Section [s2.9]), applied to one-dimensional wavefunctions, yields

$$\psi(x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \bar{\psi}(k, t) e^{+i k x} dk,$$

$$\bar{\psi}(k, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \psi(x, t) e^{-i k x} dx,$$

where  $k$  represents wavenumber. However,  $p = \hbar k$ . Hence, we can also write

$$\psi(x, t) = \frac{1}{\sqrt{2\pi \hbar}} \int_{-\infty}^{\infty} \phi(p, t) e^{+i p x / \hbar} dp,$$

$$\phi(p, t) = \frac{1}{\sqrt{2\pi \hbar}} \int_{-\infty}^{\infty} \psi(x, t) e^{-i p x / \hbar} dx,$$

where  $\phi(p, t) = \bar{\psi}(k, t) / \sqrt{\hbar}$  is the momentum-space equivalent to the real-space wavefunction  $\psi(x, t)$ .

At this stage, it is convenient to introduce a useful function called the *Dirac delta-function*. This function, denoted  $\delta(x)$ , was first devised by Paul Dirac, and has the following rather unusual properties:  $\delta(x)$  is zero for  $x \neq 0$ , and is infinite at  $x = 0$ . However, the singularity at  $x = 0$  is such that

$$\int_{-\infty}^{\infty} \delta(x) dx = 1. \quad (3.6.1)$$

The delta-function is an example of what is known as a *generalized function*: that is, its value is not well defined at all  $x$ , but its integral is well defined. Consider the integral

$$\int_{-\infty}^{\infty} f(x) \delta(x) dx. \quad (3.6.2)$$

Because  $\delta(x)$  is only non-zero infinitesimally close to  $x = 0$ , we can safely replace  $f(x)$  by  $f(0)$  in the previous integral (assuming  $f(x)$  is well behaved at  $x = 0$ ), to give

$$\int_{-\infty}^{\infty} f(x) \delta(x) dx = f(0) \int_{-\infty}^{\infty} \delta(x) dx = f(0), \quad (3.6.3)$$

where use has been made of Equation ([e3.64a]). A simple generalization of this result yields

$$\int_{-\infty}^{\infty} f(x) \delta(x - x_0) dx = f(x_0), \quad (3.6.4)$$

which can also be thought of as an alternative definition of a delta-function.

Suppose that  $\psi(x) = \delta(x - x_0)$ . It follows from Equations ([e3.65]) and ([e3.69]) that

$$\phi(p) = \frac{e^{-i p x_0 / \hbar}}{\sqrt{2\pi \hbar}} \quad (3.6.5)$$

Hence, Equation ([e3.64]) yields the important result

$$\delta(x - x_0) = \frac{1}{2\pi \hbar} \int_{-\infty}^{\infty} e^{+i p (x - x_0) / \hbar} dp. \quad (3.6.6)$$

Similarly,

$$\delta(p - p_0) = \frac{1}{2\pi \hbar} \int_{-\infty}^{\infty} e^{+i (p - p_0) x / \hbar} dx. \quad (3.6.7)$$

It turns out that we can just as easily formulate quantum mechanics using the momentum-space wavefunction,  $\phi(p, t)$ , as the real-space wavefunction,  $\psi(x, t)$ . The former scheme is known as the *momentum representation* of quantum mechanics. In the momentum representation, wavefunctions are the Fourier transforms of the equivalent real-space wavefunctions, and dynamical



variables are represented by different operators. Furthermore, by analogy with Equation ([e3.55]), the expectation value of some operator  $O(p)$  takes the form

$$\langle O \rangle = \int_{-\infty}^{\infty} \phi^*(p, t) O(p) \phi(p, t) dp. \quad (3.6.8)$$

Consider momentum. We can write

$$\begin{aligned} \langle p \rangle &= \int_{-\infty}^{\infty} \psi^*(x, t) \left( -i\hbar \frac{\partial}{\partial x} \right) \psi(x, t) dx \\ &= \frac{1}{2\pi\hbar} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \phi^*(p', t) \phi(p, t) p e^{+i(p-p')x/\hbar} dx dp dp', \end{aligned}$$

where use has been made of Equation ([e3.64]). However, it follows from Equation ([e3.72]) that

$$\langle p \rangle = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \phi^*(p', t) \phi(p, t) p \delta(p-p') dp dp'. \quad (3.6.9)$$

Hence, using Equation ([e3.69]), we obtain

$$\langle p \rangle = \int_{-\infty}^{\infty} \phi^*(p, t) p \phi(p, t) dp = \int_{-\infty}^{\infty} p |\phi|^2 dp. \quad (3.6.10)$$

Evidently, momentum is represented by the operator  $p$  in the momentum representation. The previous expression also strongly suggests [by comparison with Equation ([e3.22])] that  $|\phi(p, t)|^2$  can be interpreted as the probability density of a measurement of momentum yielding the value  $p$  at time  $t$ . It follows that  $\phi(p, t)$  must satisfy an analogous normalization condition to Equation ([e3.4]): that is,

$$\int_{-\infty}^{\infty} |\phi(p, t)|^2 dp = 1. \quad (3.6.11)$$

Consider displacement. We can write

$$\begin{aligned} \langle x \rangle &= \int_{-\infty}^{\infty} \psi^*(x, t) x \psi(x, t) dx \\ &= \frac{1}{2\pi\hbar} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \phi^*(p', t) \phi(p, t) \left( -i\hbar \frac{\partial}{\partial p} \right) e^{+i(p-p')x/\hbar} dx dp dp'. \end{aligned}$$

Integration by parts yields

$$\langle x \rangle = \frac{1}{2\pi\hbar} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \phi^*(p', t) e^{+i(p-p')x/\hbar} \left( i\hbar \frac{\partial}{\partial p} \right) \phi(p, t) dx dp dp'. \quad (3.6.12)$$

Hence, making use of Equations ([e3.72]) and ([e3.69]), we obtain

$$\langle x \rangle = \frac{1}{2\pi\hbar} \int_{-\infty}^{\infty} \phi^*(p) \left( i\hbar \frac{\partial}{\partial p} \right) \phi(p) dp. \quad (3.6.13)$$

Evidently, displacement is represented by the operator

$$x \equiv i\hbar \frac{\partial}{\partial p} \quad (3.6.14)$$

in the momentum representation.

Finally, let us consider the normalization of the momentum-space wavefunction  $\phi(p, t)$ . We have

$$\int_{-\infty}^{\infty} \psi^*(x, t) \psi(x, t) dx = \frac{1}{2\pi\hbar} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \phi^*(p', t) \phi(p, t) e^{+i(p-p')x/\hbar} dx dp dp'. \quad (3.6.15)$$

Thus, it follows from Equations ([e3.69]) and ([e3.72]) that

$$\int_{-\infty}^{\infty} |\psi(x, t)|^2 dx = \int_{-\infty}^{\infty} |\phi(p, t)|^2 dp. \quad (3.6.16)$$

Hence, if  $\psi(x, t)$  is properly normalized [see Equation ([\[e3.4\]](#))] then  $\phi(p, t)$ , as defined in Equation ([\[e3.65\]](#)), is also properly normalized [see Equation ([\[enormp\]](#))].

The existence of the momentum representation illustrates an important point. Namely, there are many different, but entirely equivalent, ways of mathematically formulating quantum mechanics. For instance, it is also possible to represent wavefunctions as row and column vectors, and dynamical variables as matrices that act upon these vectors.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [3.6: Momentum Representation](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

### 3.7: Heisenberg's Uncertainty Principle

Consider a real-space Hermitian operator,  $O(x)$ . A straightforward generalization of Equation ([e3.55a]) yields

$$\int_{-\infty}^{\infty} \psi_1^* (O \psi_2) dx = \int_{-\infty}^{\infty} (O \psi_1)^* \psi_2 dx, \quad (3.7.1)$$

where  $\psi_1(x)$  and  $\psi_2(x)$  are general functions.

Let  $f = (A - \langle A \rangle) \psi$ , where  $A(x)$  is an Hermitian operator, and  $\psi(x)$  a general wavefunction. We have

$$\int_{-\infty}^{\infty} |f|^2 dx = \int_{-\infty}^{\infty} f^* f dx = \int_{-\infty}^{\infty} [(A - \langle A \rangle) \psi]^* [(A - \langle A \rangle) \psi] dx. \quad (3.7.2)$$

Making use of Equation ([e3.84]), we obtain

$$\int_{-\infty}^{\infty} |f|^2 dx = \int_{-\infty}^{\infty} \psi^* (A - \langle A \rangle)^2 \psi dx = \sigma_A^2, \quad (3.7.3)$$

where  $\sigma_A^2$  is the variance of  $A$ . [See Equation ([e3.24a]).] Similarly, if  $g = (B - \langle B \rangle) \psi$ , where  $B$  is a second Hermitian operator, then

$$\int_{-\infty}^{\infty} |g|^2 dx = \sigma_B^2, \quad (3.7.4)$$

Now, there is a standard result in mathematics, known as the *Schwartz inequality*, which states that

$$\left| \int_a^b f^*(x) g(x) dx \right|^2 \leq \int_a^b |f(x)|^2 dx \int_a^b |g(x)|^2 dx, \quad (3.7.5)$$

where  $f$  and  $g$  are two general functions. Furthermore, if  $z$  is a complex number then

$$|z|^2 = [\text{Re}(z)]^2 + [\text{Im}(z)]^2 \geq [\text{Im}(z)]^2 = \left[ \frac{1}{2i} (z - z^*) \right]^2. \quad (3.7.6)$$

Hence, if  $z = \int_{-\infty}^{\infty} f^* g dx$  then Equations ([e3.86])–([e3.89]) yield

$$\sigma_A^2 \sigma_B^2 \geq \left[ \frac{1}{2i} (z - z^*) \right]^2. \quad (3.7.7)$$

However,

$$z = \int_{-\infty}^{\infty} [(A - \langle A \rangle) \psi]^* [(B - \langle B \rangle) \psi] dx = \int_{-\infty}^{\infty} \psi^* (A - \langle A \rangle) (B - \langle B \rangle) \psi dx, \quad (3.7.8)$$

where use has been made of Equation ([e3.84]). The previous equation reduces to

$$z = \int_{-\infty}^{\infty} \psi^* A B \psi dx - \langle A \rangle \langle B \rangle. \quad (3.7.9)$$

Furthermore, it is easily demonstrated that

$$z^* = \int_{-\infty}^{\infty} \psi^* B A \psi dx - \langle A \rangle \langle B \rangle. \quad (3.7.10)$$

Hence, Equation ([e3.90]) gives

$$\sigma_A^2 \sigma_B^2 \geq \left( \frac{1}{2i} \langle [A, B] \rangle \right)^2, \quad (3.7.11)$$

where

$$[A, B] \equiv A B - B A. \quad (3.7.12)$$

Equation (e3.94) is the general form of *Heisenberg's uncertainty principle* in quantum mechanics. It states that if two dynamical variables are represented by the two Hermitian operators  $A$  and  $B$ , and these operators do not commute (i.e.,  $AB \neq BA$ ), then it is impossible to simultaneously (exactly) measure the two variables. Instead, the product of the variances in the measurements is always greater than some critical value, which depends on the extent to which the two operators do not commute.

For instance, displacement and momentum are represented (in real-space) by the operators  $x$  and  $p \equiv -i\hbar\partial/\partial x$ , respectively. Now, it is easily demonstrated that

$$[x, p] = i\hbar. \quad (3.7.13)$$

Thus,

$$\sigma_x \sigma_p \geq \frac{\hbar}{2}, \quad (3.7.14)$$

which can be recognized as the standard displacement-momentum uncertainty principle (see Section [sun]). It turns out that the minimum uncertainty (i.e.,  $\sigma_x \sigma_p = \hbar/2$ ) is only achieved by Gaussian wave-packets (see Section [s2.9]): that is,

$$\psi(x) = \frac{e^{+ip_0 x/\hbar}}{(2\pi\sigma_x^2)^{1/4}} e^{-(x-x_0)^2/4\sigma_x^2} \quad (3.7.15)$$

$$\phi(p) = \frac{e^{-ipx_0/\hbar}}{(2\pi\sigma_p^2)^{1/4}} e^{-(p-p_0)^2/4\sigma_p^2} \quad (3.7.16)$$

where  $\phi(p)$  is the momentum-space equivalent of  $\psi(x)$ .

Energy and time are represented by the operators  $H \equiv i\hbar\partial/\partial t$  and  $t$ , respectively. These operators do not commute, indicating that energy and time cannot be measured simultaneously. In fact,

$$[H, t] = i\hbar, \quad (3.7.17)$$

so

$$\sigma_E \sigma_t \geq \frac{\hbar}{2}. \quad (3.7.18)$$

This can be written, somewhat less exactly, as

$\Delta E \Delta t \gtrsim \hbar$  are the uncertainties in energy and time, respectively. The previous expression is generally known as the *energy-time uncertainty principle*.

For instance, suppose that a particle passes some fixed point on the  $x$ -axis. Because the particle is, in reality, an extended wave-packet, it takes a certain amount of time,  $\Delta t$ , for the particle to pass. Thus, there is an uncertainty,  $\Delta t$ , in the arrival time of the particle. Moreover, because  $E = \hbar\omega$ , the only wavefunctions that have unique energies are those with unique frequencies: that is, plane-waves. Because a wave-packet of finite extent is made up of a combination of plane-waves of different wavenumbers, and, hence, different frequencies, there will be an uncertainty  $\Delta E$  in the particle's energy that is proportional to the range of frequencies of the plane-waves making up the wave-packet. The more compact the wave-packet (and, hence, the smaller  $\Delta t$ ), the larger the range of frequencies of the constituent plane-waves (and, hence, the large  $\Delta E$ ), and vice versa.

To be more exact, if  $\psi(t)$  is the wavefunction measured at the fixed point as a function of time then we can write

$$\psi(t) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \chi(E) e^{-iEt/\hbar} dE \quad (3.7.19)$$

In other words, we can express  $\psi(t)$  as a linear combination of plane-waves of definite energy  $E$ . Here,  $\chi(E)$  is the complex amplitude of plane-waves of energy  $E$  in this combination.

By Fourier's theorem, we also have

$$\chi(E) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \psi(t) e^{+iEt/\hbar} dt \quad (3.7.20)$$

For instance, if  $\psi(t)$  is a Gaussian then it is easily shown that  $\chi(E)$  is also a Gaussian: that is,

$$\psi(t) = \frac{e^{-iE_0 t/\hbar}}{(2\pi\sigma_t^2)^{1/4}} e^{-(t-t_0)^2/4\sigma_t^2} \quad (3.7.21)$$

$$\chi(E) = \frac{e^{+iEt_0/\hbar}}{(2\pi\sigma_E^2)^{1/4}} e^{-(E-E_0)^2/4\sigma_E^2} \quad (3.7.22)$$

where  $\sigma_E \sigma_t = \hbar/2$ . As before, Gaussian wave-packets satisfy the minimum uncertainty principle  $\sigma_E \sigma_t = \hbar/2$ . Conversely, non-Gaussian wave-packets are characterized by  $\sigma_E \sigma_t > \hbar/2$ .

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [3.7: Heisenberg's Uncertainty Principle](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 3.8: Eigenstates and Eigenvalues

Consider a general real-space operator,  $A(x)$ . When this operator acts on a general wavefunction  $\psi(x)$  the result is usually a wavefunction with a completely different shape. However, there are certain special wavefunctions which are such that when  $A$  acts on them the result is just a multiple of the original wavefunction. These special wavefunctions are called *eigenstates*, and the multiples are called *eigenvalues*. Thus, if

$$A \psi_a(x) = a \psi_a(x), \quad (3.8.1)$$

where  $a$  is a complex number, then  $\psi_a$  is called an eigenstate of  $A$  corresponding to the eigenvalue  $a$ .

Suppose that  $A$  is an Hermitian operator corresponding to some physical dynamical variable. Consider a particle whose wavefunction is  $\psi_a$ . The expectation of value  $A$  in this state is simply [see Equation ([e3.55])]

$$\langle A \rangle = \int_{-\infty}^{\infty} \psi_a^* A \psi_a dx = a \int_{-\infty}^{\infty} \psi_a^* \psi_a dx = a, \quad (3.8.2)$$

where use has been made of Equation ([e3.107]) and the normalization condition ([e3.4]). Moreover,

$$\langle A^2 \rangle = \int_{-\infty}^{\infty} \psi_a^* A^2 \psi_a dx = a \int_{-\infty}^{\infty} \psi_a^* A \psi_a dx = a^2 \int_{-\infty}^{\infty} \psi_a^* \psi_a dx = a^2, \quad (3.8.3)$$

so the variance of  $A$  is [cf., Equation ([e3.24a])]

$$\sigma_A^2 = \langle A^2 \rangle - \langle A \rangle^2 = a^2 - a^2 = 0. \quad (3.8.4)$$

The fact that the variance is zero implies that every measurement of  $A$  is bound to yield the same result: namely,  $a$ . Thus, the eigenstate  $\psi_a$  is a state that is associated with a unique value of the dynamical variable corresponding to  $A$ . This unique value is simply the associated eigenvalue.

It is easily demonstrated that the eigenvalues of an Hermitian operator are all real. Recall [from Equation ([e3.84])] that an Hermitian operator satisfies

$$\int_{-\infty}^{\infty} \psi_1^* (A \psi_2) dx = \int_{-\infty}^{\infty} (A \psi_1)^* \psi_2 dx. \quad (3.8.5)$$

Hence, if  $\psi_1 = \psi_2 = \psi_a$  then

$$\int_{-\infty}^{\infty} \psi_a^* (A \psi_a) dx = \int_{-\infty}^{\infty} (A \psi_a)^* \psi_a dx, \quad (3.8.6)$$

which reduces to [see Equation ([e3.107])]

$$a = a^*, \quad (3.8.7)$$

assuming that  $\psi_a$  is properly normalized.

Two wavefunctions,  $\psi_1(x)$  and  $\psi_2(x)$ , are said to be *orthogonal* if

$$\int_{-\infty}^{\infty} \psi_1^* \psi_2 dx = 0. \quad (3.8.8)$$

Consider two eigenstates of  $A$ ,  $\psi_a$  and  $\psi_{a'}$ , which correspond to the two different eigenvalues  $a$  and  $a'$ , respectively. Thus,

$$\begin{aligned} A \psi_a &= a \psi_a, \\ A \psi_{a'} &= a' \psi_{a'}. \end{aligned}$$

Multiplying the complex conjugate of the first equation by  $\psi_{a'}$ , and the second equation by  $\psi_a^*$ , and then integrating over all  $x$ , we obtain

$$\int_{-\infty}^{\infty} (A \psi_a)^* \psi_{a'} dx = a \int_{-\infty}^{\infty} \psi_a^* \psi_{a'} dx,$$

$$\int_{-\infty}^{\infty} \psi_a^* (A \psi_{a'}) dx = a' \int_{-\infty}^{\infty} \psi_a^* \psi_{a'} dx.$$

However, from Equation ([e3.111]), the left-hand sides of the previous two equations are equal. Hence, we can write

$$(a - a') \int_{-\infty}^{\infty} \psi_a^* \psi_{a'} dx = 0. \quad (3.8.9)$$

By assumption,  $a \neq a'$ , yielding

$$\int_{-\infty}^{\infty} \psi_a^* \psi_{a'} dx = 0. \quad (3.8.10)$$

In other words, eigenstates of an Hermitian operator corresponding to different eigenvalues are automatically orthogonal.

Consider two eigenstates of  $A$ ,  $\psi_a$  and  $\psi_{a'}$ , that correspond to the same eigenvalue,  $a$ . Such eigenstates are termed *degenerate*. The previous proof of the orthogonality of different eigenstates fails for degenerate eigenstates. Note, however, that any linear combination of  $\psi_a$  and  $\psi_{a'}$  is also an eigenstate of  $A$  corresponding to the eigenvalue  $a$ . Thus, even if  $\psi_a$  and  $\psi_{a'}$  are not orthogonal, we can always choose two linear combinations of these eigenstates that are orthogonal. For instance, if  $\psi_a$  and  $\psi_{a'}$  are properly normalized, and

$$\int_{-\infty}^{\infty} \psi_a^* \psi_{a'} dx = c, \quad (3.8.11)$$

then it is easily demonstrated that

$$\psi_a'' = \frac{|c|}{\sqrt{1 - |c|^2}} (\psi_a - c^{-1} \psi_{a'}) \quad (3.8.12)$$

is a properly normalized eigenstate of  $A$ , corresponding to the eigenvalue  $a$ , that is orthogonal to  $\psi_a$ . It is straightforward to generalize the previous argument to three or more degenerate eigenstates. Hence, we conclude that the eigenstates of an Hermitian operator are, or can be chosen to be, mutually orthogonal.

It is also possible to demonstrate that the eigenstates of an Hermitian operator form a complete set : that is, any general wavefunction can be written as a linear combination of these eigenstates. However, the proof is quite difficult, and we shall not attempt it here.

In summary, given an Hermitian operator  $A$ , any general wavefunction,  $\psi(x)$ , can be written

$$\psi = \sum_i c_i \psi_i, \quad (3.8.13)$$

where the  $c_i$  are complex weights, and the  $\psi_i$  are the properly normalized (and mutually orthogonal) eigenstates of  $A$ : that is,

$$A \psi_i = a_i \psi_i, \quad (3.8.14)$$

where  $a_i$  is the eigenvalue corresponding to the eigenstate  $\psi_i$ , and

$$\int_{-\infty}^{\infty} \psi_i^* \psi_j dx = \delta_{ij}. \quad (3.8.15)$$

Here,  $\delta_{ij}$  is called the *Kronecker delta-function*, and takes the value unity when its two indices are equal, and zero otherwise.

It follows from Equations ([e3.123]) and ([e3.125]) that

$$c_i = \int_{-\infty}^{\infty} \psi_i^* \psi dx. \quad (3.8.16)$$

Thus, the expansion coefficients in Equation ([e3.123]) are easily determined, given the wavefunction  $\psi$  and the eigenstates  $\psi_i$ . Moreover, if  $\psi$  is a properly normalized wavefunction then Equations ([e3.123]) and ([e3.125]) yield

$$\sum_i |c_i|^2 = 1. \quad (3.8.17)$$

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [3.8: Eigenstates and Eigenvalues](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).



### 3.9: Measurement

Suppose that  $A$  is an Hermitian operator corresponding to some dynamical variable. By analogy with the discussion in Section [\[scoil\]](#), we expect that if a measurement of  $A$  yields the result  $a$  then the act of measurement will cause the wavefunction to collapse to a state in which a measurement of  $A$  is bound to give the result  $a$ . What sort of wavefunction,  $\psi$ , is such that a measurement of  $A$  is bound to yield a certain result,  $a$ ? Well, expressing  $\psi$  as a linear combination of the eigenstates of  $A$ , we have

$$\psi = \sum_i c_i \psi_i, \quad (3.9.1)$$

where  $\psi_i$  is an eigenstate of  $A$  corresponding to the eigenvalue  $a_i$ . If a measurement of  $A$  is bound to yield the result  $a$  then

$$\langle A \rangle = a, \quad (3.9.2)$$

and

$$\sigma_A^2 = \langle A^2 \rangle - \langle A \rangle^2 = 0. \quad (3.9.3)$$

Now, it is easily seen that

$$\begin{aligned} \langle A \rangle &= \sum_i |c_i|^2 a_i, \\ \langle A^2 \rangle &= \sum_i |c_i|^2 a_i^2. \end{aligned}$$

Thus, Equation ([\[e4.130\]](#)) gives

$$\sum_i a_i^2 |c_i|^2 - \left( \sum_i a_i |c_i|^2 \right)^2 = 0. \quad (3.9.4)$$

Furthermore, the normalization condition yields

$$\sum_i |c_i|^2 = 1. \quad (3.9.5)$$

For instance, suppose that there are only two eigenstates. The previous two equations then reduce to  $|c_1|^2 = x$ , and  $|c_2|^2 = 1 - x$ , where  $0 \leq x \leq 1$ , and

$$(a_1 - a_2)^2 x (1 - x) = 0. \quad (3.9.6)$$

The only solutions are  $x = 0$  and  $x = 1$ . This result can easily be generalized to the case where there are more than two eigenstates. It follows that a state associated with a definite value of  $A$  is one in which one of the  $|c_i|^2$  is unity, and all of the others are zero. In other words, the only states associated with definite values of  $A$  are the eigenstates of  $A$ . It immediately follows that the result of a measurement of  $A$  must be one of the eigenvalues of  $A$ . Moreover, if a general wavefunction is expanded as a linear combination of the eigenstates of  $A$ , like in Equation ([\[e4.128\]](#)), then it is clear from Equation ([\[e4.131\]](#)), and the general definition of a mean, that the probability of a measurement of  $A$  yielding the eigenvalue  $a_i$  is simply  $|c_i|^2$ , where  $c_i$  is the coefficient in front of the  $i$ th eigenstate in the expansion. Note, from Equation ([\[e4.134\]](#)), that these probabilities are properly normalized: that is, the probability of a measurement of  $A$  resulting in any possible answer is unity. Finally, if a measurement of  $A$  results in the eigenvalue  $a_i$  then immediately after the measurement the system will be left in the eigenstate corresponding to  $a_i$ .

Consider two physical dynamical variables represented by the two Hermitian operators  $A$  and  $B$ . Under what circumstances is it possible to simultaneously measure these two variables (exactly)? Well, the possible results of measurements of  $A$  and  $B$  are the eigenvalues of  $A$  and  $B$ , respectively. Thus, to simultaneously measure  $A$  and  $B$  (exactly) there must exist states which are simultaneous eigenstates of  $A$  and  $B$ . In fact, in order for  $A$  and  $B$  to be simultaneously measurable under all circumstances, we need all of the eigenstates of  $A$  to also be eigenstates of  $B$ , and vice versa, so that all states associated with unique values of  $A$  are also associated with unique values of  $B$ , and vice versa.

Now, we have already seen, in Section [1.8](#), that if  $A$  and  $B$  do not commute (i.e., if  $AB \neq BA$ ) then they cannot be simultaneously measured. This suggests that the condition for simultaneous measurement is that  $A$  and  $B$  should commute.

Suppose that this is the case, and that the  $\psi_i$  and  $a_i$  are the normalized eigenstates and eigenvalues of  $A$ , respectively. It follows that

$$(A B - B A) \psi_i = (A B - B a_i) \psi_i = (A - a_i) B \psi_i = 0, \quad (3.9.7)$$

or

$$A (B \psi_i) = a_i (B \psi_i). \quad (3.9.8)$$

Thus,  $B \psi_i$  is an eigenstate of  $A$  corresponding to the eigenvalue  $a_i$  (though not necessarily a normalized one). In other words,  $B \psi_i \propto \psi_i$ , or

$$B \psi_i = b_i \psi_i, \quad (3.9.9)$$

where  $b_i$  is a constant of proportionality. Hence,  $\psi_i$  is an eigenstate of  $B$ , and, thus, a simultaneous eigenstate of  $A$  and  $B$ . We conclude that if  $A$  and  $B$  commute then they possess simultaneous eigenstates, and are thus simultaneously measurable (exactly).

## Continuous Eigenvalues

In the previous two sections, it was tacitly assumed that we were dealing with operators possessing discrete eigenvalues and square-integrable eigenstates. Unfortunately, some operators—most notably,  $x$  and  $p$ —possess eigenvalues that lie in a continuous range and non-square-integrable eigenstates (in fact, these two properties go hand in hand). Let us, therefore, investigate the eigenstates and eigenvalues of the displacement and momentum operators.

Let  $\psi_x(x, x')$  be the eigenstate of  $x$  corresponding to the eigenvalue  $x'$ . It follows that

$$x \psi_x(x, x') = x' \psi_x(x, x') \quad (3.9.10)$$

for all  $x$ . Consider the Dirac delta-function  $\delta(x - x')$ . We can write

$$x \delta(x - x') = x' \delta(x - x'), \quad (3.9.11)$$

because  $\delta(x - x')$  is only non-zero infinitesimally close to  $x = x'$ . Evidently,  $\psi_x(x, x')$  is proportional to  $\delta(x - x')$ . Let us make the constant of proportionality unity, so that

$$\psi_x(x, x') = \delta(x - x'). \quad (3.9.12)$$

It is easily demonstrated that

$$\int_{-\infty}^{\infty} \delta(x - x') \delta(x - x'') dx = \delta(x' - x''). \quad (3.9.13)$$

Hence,  $\psi_x(x, x')$  satisfies the orthonormality condition

$$\int_{-\infty}^{\infty} \psi_x^*(x, x') \psi_x(x, x'') dx = \delta(x' - x''). \quad (3.9.14)$$

This condition is analogous to the orthonormality condition ([e3.125]) satisfied by square-integrable eigenstates. Now, by definition,  $\delta(x - x')$  satisfies

$$\int_{-\infty}^{\infty} f(x) \delta(x - x') dx = f(x'), \quad (3.9.15)$$

where  $f(x)$  is a general function. We can thus write

$$\psi(x) = \int_{-\infty}^{\infty} c(x') \psi_x(x, x') dx', \quad (3.9.16)$$

where  $c(x') = \psi(x')$ , or

$$c(x') = \int_{-\infty}^{\infty} \psi_x^*(x, x') \psi(x) dx. \quad (3.9.17)$$

In other words, we can expand a general wavefunction  $\psi(x)$  as a linear combination of the eigenstates,  $\psi_x(x, x')$ , of the displacement operator. Equations ([e4.144]) and ([e4.145]) are analogous to Equations ([e3.123]) and ([e3.126]), respectively, for

square-integrable eigenstates. Finally, by analogy with the results in Section 1.9, the probability density of a measurement of  $x$  yielding the value  $x'$  is  $|c(x')|^2$ , which is equivalent to the standard result  $|\psi(x')|^2$ . Moreover, these probabilities are properly normalized provided  $\psi(x)$  is properly normalized [cf., Equation ([e3.127])]: that is,

$$\int_{-\infty}^{\infty} |c(x')|^2 dx' = \int_{-\infty}^{\infty} |\psi(x')|^2 dx' = 1. \quad (3.9.18)$$

Finally, if a measurement of  $x$  yields the value  $x'$  then the system is left in the corresponding displacement eigenstate,  $\psi_x(x, x')$ , immediately after the measurement. That is, the wavefunction collapses to a “spike-function”,  $\delta(x - x')$ , as discussed in Section [scol].

Now, an eigenstate of the momentum operator  $p \equiv -i\hbar \partial/\partial x$  corresponding to the eigenvalue  $p'$  satisfies

$$-i\hbar \frac{\partial \psi_p(x, p')}{\partial x} = p' \psi_p(x, p'). \quad (3.9.19)$$

It is evident that

$$\psi_p(x, p') \propto e^{+ip'x/\hbar}. \quad (3.9.20)$$

We require  $\psi_p(x, p')$  to satisfy an analogous orthonormality condition to Equation ([e4.143]): that is,

$$\int_{-\infty}^{\infty} \psi_p^*(x, p') \psi_p(x, p'') dx = \delta(p' - p''). \quad (3.9.21)$$

Thus, it follows from Equation ([e3.72]) that the constant of proportionality in Equation ([e4.148]) should be  $(2\pi\hbar)^{-1/2}$ ; that is,

$$\psi_p(x, p') = \frac{e^{+ip'x/\hbar}}{(2\pi\hbar)^{1/2}}. \quad (3.9.22)$$

Furthermore, according to Equations ([e3.64]) and ([e3.65]),

$$\psi(x) = \int_{-\infty}^{\infty} c(p') \psi_p(x, p') dp', \quad (3.9.23)$$

where  $c(p') = \phi(p')$  [see Equation ([e3.65])], or

$$c(p') = \int_{-\infty}^{\infty} \psi_p^*(x, p') \psi(x) dx. \quad (3.9.24)$$

In other words, we can expand a general wavefunction  $\psi(x)$  as a linear combination of the eigenstates,  $\psi_p(x, p')$ , of the momentum operator. Equations ([e4.152]) and ([e4.153]) are again analogous to Equations ([e3.123]) and ([e3.126]), respectively, for square-integrable eigenstates. Likewise, the probability density of a measurement of  $p$  yielding the result  $p'$  is  $|c(p')|^2$ , which is equivalent to the standard result  $|\phi(p')|^2$ . The probabilities are also properly normalized provided  $\psi(x)$  is properly normalized [cf., Equation ([e3.83])]: that is,

$$\int_{-\infty}^{\infty} |c(p')|^2 dp' = \int_{-\infty}^{\infty} |\phi(p')|^2 dp' = \int_{-\infty}^{\infty} |\psi(x')|^2 dx' = 1. \quad (3.9.25)$$

Finally, if a measurement of  $p$  yields the value  $p'$  then the system is left in the corresponding momentum eigenstate,  $\psi_p(x, p')$ , immediately after the measurement.

## Contributors and Attributions

- Richard Fitzpatrick (Professor of Physics, The University of Texas at Austin)

This page titled 3.9: Measurement is shared under a not declared license and was authored, remixed, and/or curated by Richard Fitzpatrick.

## 3.10: Stationary States

An eigenstate of the energy operator  $H \equiv i \hbar \partial / \partial t$  corresponding to the eigenvalue  $E_i$  satisfies

$$i \hbar \frac{\partial \psi_E(x, t, E_i)}{\partial t} = E_i \psi_E(x, t, E_i). \quad (3.10.1)$$

It is evident that this equation can be solved by writing

$$\psi_E(x, t, E_i) = \psi_i(x) e^{-i E_i t / \hbar}, \quad (3.10.2)$$

where  $\psi_i(x)$  is a properly normalized stationary (i.e., non-time-varying) wavefunction. The wavefunction  $\psi_E(x, t, E_i)$  corresponds to a so-called *stationary state*, because the probability density  $|\psi_E|^2$  is non-time-varying. Note that a stationary state is associated with a unique value for the energy. Substitution of the previous expression into Schrödinger's equation ([e3.1]) yields the equation satisfied by the stationary wavefunction:

$$\frac{\hbar^2}{2m} \frac{d^2 \psi_i}{dx^2} = [V(x) - E_i] \psi_i. \quad (3.10.3)$$

This is known as the *time-independent Schrödinger equation*. More generally, this equation takes the form

$$H \psi_i = E_i \psi_i, \quad (3.10.4)$$

where  $H$  is assumed not to be an explicit function of  $t$ . Of course, the  $\psi_i$  satisfy the usual orthonormality condition:

$$\int_{-\infty}^{\infty} \psi_i^* \psi_j dx = \delta_{ij}. \quad (3.10.5)$$

Moreover, we can express a general wavefunction as a linear combination of energy eigenstates:

$$\psi(x, t) = \sum_i c_i \psi_i(x) e^{-i E_i t / \hbar}, \quad (3.10.6)$$

where

$$c_i = \int_{-\infty}^{\infty} \psi_i^*(x) \psi(x, 0) dx. \quad (3.10.7)$$

Here,  $|c_i|^2$  is the probability that a measurement of the energy will yield the eigenvalue  $E_i$ . Furthermore, immediately after such a measurement, the system is left in the corresponding energy eigenstate. The generalization of the previous results to the case where  $H$  has continuous eigenvalues is straightforward.

If a dynamical variable is represented by some Hermitian operator  $A$  that commutes with  $H$  (so that it has simultaneous eigenstates with  $H$ ), and contains no specific time dependence, then it is evident from Equations ([e4.157]) and ([e4.158]) that the expectation value and variance of  $A$  are time independent. In this sense, the dynamical variable in question is a constant of the motion.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled 3.10: Stationary States is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

### 3.11: Exercises

1. Monochromatic light with a wavelength of  $6000\text{\AA}$  passes through a fast shutter that opens for  $10^{-9}$  sec. What is the subsequent spread in wavelengths of the no longer monochromatic light?
2. Calculate  $\langle x \rangle$ ,  $\langle x^2 \rangle$ , and  $\sigma_x$ , as well as  $\langle p \rangle$ ,  $\langle p^2 \rangle$ , and  $\sigma_p$ , for the normalized wavefunction

$$\psi(x) = \sqrt{\frac{2a^3}{\pi}} \frac{1}{x^2 + a^2}. \quad (3.11.1)$$

Use these to find  $\sigma_x \sigma_p$ . Note that  $\int_{-\infty}^{\infty} dx/(x^2 + a^2) = \pi/a$ .

3. Classically, if a particle is not observed then the probability of finding it in a one-dimensional box of length  $L$ , which extends from  $x = 0$  to  $x = L$ , is a constant  $1/L$  per unit length. Show that the classical expectation value of  $x$  is  $L/2$ , the expectation value of  $x^2$  is  $L^2/3$ , and the standard deviation of  $x$  is  $L/\sqrt{12}$ .
4. Demonstrate that if a particle in a one-dimensional stationary state is bound then the expectation value of its momentum must be zero.
5. Suppose that  $V(x)$  is complex. Obtain an expression for  $\partial P(x, t)/\partial t$  and  $d/dt \int P(x, t) dx$  from Schrödinger's equation. What does this tell us about a complex  $V(x)$ ?
6.  $\psi_1(x)$  and  $\psi_2(x)$  are normalized eigenfunctions corresponding to the same eigenvalue. If

$$\int_{-\infty}^{\infty} \psi_1^* \psi_2 dx = c, \quad (3.11.2)$$

where  $c$  is real, find normalized linear combinations of  $\psi_1$  and  $\psi_2$  that are orthogonal to (a)  $\psi_1$ , (b)  $\psi_1 + \psi_2$ .

7. Demonstrate that  $p = -i\hbar \partial/\partial x$  is an Hermitian operator. Find the Hermitian conjugate of  $a = x + ip$ .
8. An operator  $A$ , corresponding to a physical quantity  $\alpha$ , has two normalized eigenfunctions  $\psi_1(x)$  and  $\psi_2(x)$ , with eigenvalues  $a_1$  and  $a_2$ . An operator  $B$ , corresponding to another physical quantity  $\beta$ , has normalized eigenfunctions  $\phi_1(x)$  and  $\phi_2(x)$ , with eigenvalues  $b_1$  and  $b_2$ . The eigenfunctions are related via

$$\begin{aligned} \psi_1 &= (2\phi_1 + 3\phi_2)/\sqrt{13}, \\ \psi_2 &= (3\phi_1 - 2\phi_2)/\sqrt{13}. \end{aligned}$$

$\alpha$  is measured and the value  $a_1$  is obtained. If  $\beta$  is then measured and then  $\alpha$  again, show that the probability of obtaining  $a_1$  a second time is  $97/169$ .

9. Demonstrate that an operator that commutes with the Hamiltonian, and contains no explicit time dependence, has an expectation value that is constant in time.
10. For a certain system, the operator corresponding to the physical quantity  $A$  does not commute with the Hamiltonian. It has eigenvalues  $a_1$  and  $a_2$ , corresponding to properly normalized eigenfunctions

$$\begin{aligned} \phi_1 &= (u_1 + u_2)/\sqrt{2}, \\ \phi_2 &= (u_1 - u_2)/\sqrt{2}, \end{aligned}$$

where  $u_1$  and  $u_2$  are properly normalized eigenfunctions of the Hamiltonian with eigenvalues  $E_1$  and  $E_2$ . If the system is in the state  $\psi = \phi_1$  at time  $t = 0$ , show that the expectation value of  $A$  at time  $t$  is

$$\langle A \rangle = \left( \frac{a_1 + a_2}{2} \right) + \left( \frac{a_1 - a_2}{2} \right) \cos \left( \frac{[E_1 - E_2] t}{\hbar} \right). \quad (3.11.3)$$

#### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [3.11: Exercises](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## CHAPTER OVERVIEW

### 4: One-Dimensional Potentials

In this chapter, we shall investigate the interaction of a non-relativistic particle of mass  $m$  and energy  $E$  with various one-dimensional potentials,  $V(x)$ . Because we are searching for stationary solutions with unique energies, we can write the wavefunction in the form (see Section [\[sstat\]](#))

$$\psi(x, t) = \psi(x) e^{-i E t / \hbar}, \quad (4.1)$$

where  $\psi(x)$  satisfies the time-independent Schrödinger equation:

$$\frac{d^2 \psi}{dx^2} = \frac{2m}{\hbar^2} [V(x) - E] \psi. \quad (4.2)$$

In general, the solution,  $\psi(x)$ , to the previous equation must be finite, otherwise the probability density  $|\psi|^2$  would become infinite (which is unphysical). Likewise, the solution must be continuous, otherwise the probability current ([\[eprob\]](#)) would become infinite (which is also unphysical).

[4.1: Infinite Potential Well](#)

[4.2: Square Potential Barrier](#)

[4.3: WKB Approximation](#)

[4.4: Cold Emission](#)

[4.5: Alpha Decay](#)

[4.6: Square Potential Well](#)

[4.7: Simple Harmonic Oscillator](#)

[4.E: One-Dimensional Potentials \(Exercises\)](#)

---

This page titled [4: One-Dimensional Potentials](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 4.1: Infinite Potential Well

Consider a particle of mass  $m$  and energy  $E$  moving in the following simple potential:

$$V(x) = \begin{cases} 0 & \text{for } 0 \leq x \leq a \\ \infty & \text{otherwise} \end{cases}. \quad (4.1.1)$$

It follows from Equation ([e5.2]) that if  $d^2\psi/dx^2$  (and, hence,  $\psi$ ) is to remain finite then  $\psi$  must go to zero in regions where the potential is infinite. Hence,  $\psi = 0$  in the regions  $x \leq 0$  and  $x \geq a$ . Evidently, the problem is equivalent to that of a particle trapped in a one-dimensional box of length  $a$ . The boundary conditions on  $\psi$  in the region  $0 < x < a$  are

$$\psi(0) = \psi(a) = 0. \quad (4.1.2)$$

Furthermore, it follows from Equation ([e5.2]) that  $\psi$  satisfies

$$\frac{d^2\psi}{dx^2} = -k^2 \psi \quad (4.1.3)$$

in this region, where

$$k^2 = \frac{2mE}{\hbar^2}. \quad (4.1.4)$$

Here, we are assuming that  $E > 0$ . It is easily demonstrated that there are no solutions with  $E < 0$  which are capable of satisfying the boundary conditions ([e5.4]).

The solution to Equation ([e5.5]), subject to the boundary conditions ([e5.4]), is

$$\psi_n(x) = A_n \sin(k_n x), \quad (4.1.5)$$

where the  $A_n$  are arbitrary (real) constants, and

$$k_n = \frac{n\pi}{a}, \quad (4.1.6)$$

for  $n = 1, 2, 3, \dots$ . Now, it can be seen from Equations ([e5.6]) and ([e5.8]) that the energy  $E$  is only allowed to take certain discrete values: that is,

$$E_n = \frac{n^2 \pi^2 \hbar^2}{2ma^2}. \quad (4.1.7)$$

In other words, the eigenvalues of the energy operator are discrete. This is a general feature of bounded solutions: that is, solutions for which  $|\psi| \rightarrow 0$  as  $|x| \rightarrow \infty$ . According to the discussion in Section [sstat], we expect the stationary eigenfunctions  $\psi_n(x)$  to satisfy the orthonormality constraint

$$\int_0^a \psi_n(x) \psi_m(x) dx = \delta_{nm}. \quad (4.1.8)$$

It is easily demonstrated that this is the case, provided  $A_n = \sqrt{2/a}$ . Hence,

$$\psi_n(x) = \sqrt{\frac{2}{a}} \sin\left(n\pi \frac{x}{a}\right) \quad (4.1.9)$$

for  $n = 1, 2, 3, \dots$ .

Finally, again from Section [sstat], the general time-dependent solution can be written as a linear superposition of stationary solutions:

$$\psi(x, t) = \sum_{n=0, \infty} c_n \psi_n(x) e^{-i E_n t / \hbar}, \quad (4.1.10)$$

where

$$c_n = \int_0^a \psi_n(x) \psi(x, 0) dx. \quad (4.1.11)$$

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [4.1: Infinite Potential Well](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).



## 4.2: Square Potential Barrier

Consider a particle of mass  $m$  and energy  $E > 0$  interacting with the simple square potential barrier

$$V(x) = \begin{cases} V_0 & \text{for } 0 \leq x \leq a \\ 0 & \text{otherwise} \end{cases} \quad (4.2.1)$$

where  $V_0 > 0$ . In the regions to the left and to the right of the barrier,  $\psi(x)$  satisfies

$$\frac{d^2\psi}{dx^2} = -k^2 \psi, \quad (4.2.2)$$

where  $k$  is given by Equation ([e5.6]).

Let us adopt the following solution of the previous equation to the left of the barrier (i.e.,  $x < 0$ ):

$$\psi(x) = e^{ikx} + R e^{-ikx}. \quad (4.2.3)$$

This solution consists of a plane-wave of unit amplitude traveling to the right [because the time-dependent wavefunction is multiplied by  $\exp(-i\omega t)$ , where  $\omega = E/\hbar > 0$ ], and a plane wave of complex amplitude  $R$  traveling to the left. We interpret the first plane wave as an incoming particle (or, rather, a stream of incoming particles), and the second as a particle (or stream of particles) reflected by the potential barrier. Hence,  $|R|^2$  is the probability of reflection. This can be seen by calculating the probability current ([e5.15]) in the region  $x < 0$ , which takes the form

$$j_l = v(1 - |R|^2), \quad (4.2.4)$$

where  $v = p/m = \hbar k/m$  is the classical particle velocity.

Let us adopt the following solution to Equation ([e5.15]) to the right of the barrier (i.e.  $x > a$ ):

$$\psi(x) = T e^{ikx}. \quad (4.2.5)$$

This solution consists of a plane-wave of complex amplitude  $T$  traveling to the right. We interpret this as a particle (or stream of particles) transmitted through the barrier. Hence,  $|T|^2$  is the probability of transmission. The probability current in the region  $x > a$  takes the form

$$j_r = v|T|^2. \quad (4.2.6)$$

Now, according to Equation ([e5.15]), in a stationary state (i.e.,  $\partial|\psi|^2/\partial t = 0$ ), the probability current is a spatial constant (i.e.,  $\partial j/\partial x = 0$ ). Hence, we must have  $j_l = j_r$ , or

$$|R|^2 + |T|^2 = 1. \quad (4.2.7)$$

In other words, the probabilities of reflection and transmission sum to unity, as must be the case, because reflection and transmission are the only possible outcomes for a particle incident on the barrier.

Inside the barrier (i.e.,  $0 \leq x \leq a$ ),  $\psi(x)$  satisfies

$$\frac{d^2\psi}{dx^2} = -q^2 \psi, \quad (4.2.8)$$

where

$$q^2 = \frac{2m(E - V_0)}{\hbar^2}. \quad (4.2.9)$$

Let us, first of all, consider the case where  $E > V_0$ . In this case, the general solution to Equation ([e5.21]) inside the barrier takes the form

$$\psi(x) = A e^{iqx} + B e^{-iqx}, \quad (4.2.10)$$

where  $q = \sqrt{2m(E - V_0)}/\hbar$ .

Now, the boundary conditions at the edges of the barrier (i.e., at  $x = 0$  and  $x = a$ ) are that  $\psi$  and  $d\psi/dx$  are both continuous. These boundary conditions ensure that the probability current ([\[e5.28\]](#)) remains finite and continuous across the edges of the boundary, as must be the case if it is to be a spatial constant.

Continuity of  $\psi$  and  $d\psi/dx$  at the left edge of the barrier (i.e.,  $x = 0$ ) yields

$$\begin{aligned} 1 + R &= A + B, \\ k(1 - R) &= q(A - B). \end{aligned}$$

Likewise, continuity of  $\psi$  and  $d\psi/dx$  at the right edge of the barrier (i.e.,  $x = a$ ) gives

$$\begin{aligned} A e^{iqa} + B e^{-iqa} &= T e^{ika}, \\ q(A e^{iqa} - B e^{-iqa}) &= k T e^{ika}. \end{aligned}$$

After considerable algebra, the previous four equations yield

$$|R|^2 = \frac{(k^2 - q^2)^2 \sin^2(qa)}{4k^2 q^2 + (k^2 - q^2)^2 \sin^2(qa)}, \quad (4.2.11)$$

and

$$|T|^2 = \frac{4k^2 q^2}{4k^2 q^2 + (k^2 - q^2)^2 \sin^2(qa)}. \quad (4.2.12)$$

Note that the previous two expressions satisfy the constraint ([\[e5.20\]](#)).

It is instructive to compare the quantum mechanical probabilities of reflection and transmission—([\[e5.28\]](#)) and ([\[e5.29\]](#)), respectively—with those derived from classical physics. Now, according to classical physics, if a particle of energy  $E$  is incident on a potential barrier of height  $V_0 < E$  then the particle slows down as it passes through the barrier, but is otherwise unaffected. In other words, the classical probability of reflection is zero, and the classical probability of transmission is unity.




Figure 10: Transmission (solid-curve) and reflection (dashed-curve) probabilities for a square potential barrier of width  $a = 1.25\lambda$ , where  $\lambda$  the free-space de Broglie wavelength, as a function of the ratio of the height of the barrier,  $V_0$ , to the energy,  $E$ , of the incident particle.




Figure 11: Transmission (solid-curve) and reflection (dashed-curve) probabilities for a particle of energy  $E$  incident on a square potential barrier of height  $V_0 = 0.75E$ , as a function of the ratio of the width of the barrier,  $a$ , to the free-space de Broglie wavelength,  $\lambda$

The reflection and transmission probabilities obtained from Equations (e5.28) and (e5.29), respectively, are plotted in Figures [fb1] and [fb2]. It can be seen, from Figure [fb1], that the classical result,  $|R|^2 = 0$  and  $|T|^2 = 1$ , is obtained in the limit where the height of the barrier is relatively small (i.e.,  $V_0 \ll E$ ). However, when  $V_0$  is of order  $E$ , there is a substantial probability that the incident particle will be reflected by the barrier. According to classical physics, reflection is impossible when  $V_0 < E$ .

It can also be seen, from Figure [fb2], that at certain barrier widths the probability of reflection goes to zero. It turns out that this is true irrespective of the energy of the incident particle. It is evident, from Equation (e5.28), that these special barrier widths correspond to

$$qa = n\pi, \quad (4.2.13)$$

where  $n = 1, 2, 3, \dots$ . In other words, the special barriers widths are integer multiples of half the de Broglie wavelength of the particle inside the barrier. There is no reflection at the special barrier widths because, at these widths, the backward traveling wave reflected from the left edge of the barrier interferes destructively with the similar wave reflected from the right edge of the barrier to give zero net reflected wave.

Let us, now, consider the case  $E < V_0$ . In this case, the general solution to Equation (e5.21) inside the barrier takes the form

$$\psi(x) = Ae^{qx} + Be^{-qx}, \quad (4.2.14)$$

where  $q = \sqrt{2m(V_0 - E)/\hbar^2}$ . Continuity of  $\psi$  and  $d\psi/dx$  at the left edge of the barrier (i.e.,  $x = 0$ ) yields

$$\begin{aligned} 1 + R &= A + B, \\ ik(1 - R) &= q(A - B). \end{aligned}$$

Likewise, continuity of  $\psi$  and  $d\psi/dx$  at the right edge of the barrier (i.e.,  $x = a$ ) gives

$$\begin{aligned} Ae^{qa} + Be^{-qa} &= Te^{ika}, \\ q(Ae^{qa} - Be^{-qa}) &= ikTe^{ika}. \end{aligned}$$

After considerable algebra, the previous four equations yield

$$|R|^2 = \frac{(k^2 + q^2)^2 \sinh^2(qa)}{4k^2q^2 + (k^2 + q^2)^2 \sinh^2(qa)}, \quad (4.2.15)$$

and

$$|T|^2 = \frac{4 k^2 q^2}{4 k^2 q^2 + (k^2 + q^2)^2 \sinh^2(qa)}. \quad (4.2.16)$$

These expressions can also be obtained from Equations ([e5.28]) and ([e5.29]) by making the substitution  $q \rightarrow -iq$ . Note that Equations ([e5.36]) and ([e5.37]) satisfy the constraint ([e5.20]).

It is again instructive to compare the quantum mechanical probabilities of reflection and trans/-mission—([e5.36]) and ([e5.37]), respectively—with those derived from classical physics. Now, according to classical physics, if a particle of energy  $E$  is incident on a potential barrier of height  $V_0 > E$  then the particle is reflected. In other words, the classical probability of reflection is unity, and the classical probability of transmission is zero.




Figure 12: Transmission (solid-curve) and reflection (dashed-curve) probabilities for a square potential barrier of width  $a = 0.5\lambda$ , where  $\lambda$  is the free-space de Broglie wavelength, as a function of the ratio of the energy,  $E$ , of the incoming particle to the height,  $V_0$ , of the barrier.




Figure 13: Transmission (solid-curve) and reflection (dashed-curve) probabilities for a particle of energy  $E$  incident on a square potential barrier of height  $V_0 = (4/3)E$ , as a function of the ratio of the width of the barrier,  $a$ , to the free-space de Broglie wavelength,  $\lambda$

The reflection and transmission probabilities obtained from Equations (e5.36) and (e5.37), respectively, are plotted in Figures [fb3] and [fb4]. It can be seen, from Figure [fb3], that the classical result,  $|R|^2 = 1$  and  $|T|^2 = 0$ , is obtained for relatively thin barriers (i.e.,  $qa \sim 1$ ) in the limit where the height of the barrier is relatively large (i.e.,  $V_0 \gg E$ ). However, when  $V_0$  is of order  $E$ , there is a substantial probability that the incident particle will be transmitted by the barrier. According to classical physics, transmission is impossible when  $V_0 > E$ .

It can also be seen, from Figure [fb4], that the transmission probability decays exponentially as the width of the barrier increases. Nevertheless, even for very wide barriers (i.e.,  $qa \gg 1$ ), there is a small but finite probability that a particle incident on the barrier will be transmitted. This phenomenon, which is inexplicable within the context of classical physics, is called *tunneling*.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled 4.2: Square Potential Barrier is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

### 4.3: WKB Approximation

Consider a particle of mass  $m$  and energy  $E > 0$  moving through some slowly varying potential  $V(x)$ . The particle's wavefunction satisfies

$$\frac{d^2 \psi(x)}{dx^2} = -k^2(x) \psi(x), \quad (4.3.1)$$

where

$$k^2(x) = \frac{2m[E - V(x)]}{\hbar^2}. \quad (4.3.2)$$

Let us try a solution to Equation (4.3.1) of the form

$$\psi(x) = \psi_0 \exp\left(\int_0^x i k(x') dx'\right), \quad (4.3.3)$$

where  $\psi_0$  is a complex constant. Note that this solution represents a particle propagating in the positive  $x$ -direction [because the full wavefunction is multiplied by  $\exp(-i\omega t)$ , where  $\omega = E/\hbar > 0$ ] with the continuously varying wavenumber  $k(x)$ . It follows that

$$\frac{d\psi(x)}{dx} = i k(x) \psi(x), \quad (4.3.4)$$

and

$$\frac{d^2 \psi(x)}{dx^2} = i k'(x) \psi(x) - k^2(x) \psi(x), \quad (4.3.5)$$

where  $k' \equiv dk/dx$ . A comparison of Equations (4.3.3) and (4.3.5) reveals that Equation (4.3.3) represents an approximate solution to Equation (4.3.1) provided that the first term on its right-hand side is negligible compared to the second. This yields the validity criterion  $|k'| \ll k^2$ , or

$$\frac{k}{|k'|} \gg k^{-1}. \quad (4.3.6)$$

In other words, the variation length-scale of  $k(x)$ , which is approximately the same as the variation length-scale of  $V(x)$ , must be much greater than the particle's de Broglie wavelength (which is of order  $k^{-1}$ ). Let us suppose that this is the case. Incidentally, the approximation involved in dropping the first term on the right-hand side of Equation (4.3.5) is generally known as the *WKB approximation*, after G. Wentzel, H.A. Kramers, and L. Brillouin. Similarly, Equation (4.3.3) is termed a WKB solution.

According to the WKB solution (4.3.3), the probability density remains constant: that is,

$$|\psi(x)|^2 = |\psi_0|^2, \quad (4.3.7)$$

as long as the particle moves through a region in which  $E > V(x)$ , and  $k(x)$  is consequently real (i.e., an allowed region according to classical physics). Suppose, however, that the particle encounters a potential barrier (i.e., a region from which the particle is excluded according to classical physics). By definition,  $E < V(x)$  inside such a barrier, and  $k(x)$  is consequently imaginary. Let the barrier extend from  $x = x_1$  to  $x_2$ , where  $0 < x_1 < x_2$ . The WKB solution inside the barrier is written

$$\psi(x) = \psi_1 \exp\left(-\int_{x_1}^x |k(x')| dx'\right), \quad (4.3.8)$$

where

$$\psi_1 = \psi_0 \exp\left(\int_0^{x_1} i k(x') dx'\right). \quad (4.3.9)$$

Here, we have neglected the unphysical exponentially growing solution.

According to the WKB solution (4.3.9), the probability density decays exponentially inside the barrier: that is,

$$|\psi(x)|^2 = |\psi_1|^2 \exp\left(-2 \int_{x_1}^x |k(x')| dx'\right), \quad (4.3.10)$$

where  $|\psi_1|^2$  is the probability density at the left-hand side of the barrier (i.e.,  $x = x_1$ ). It follows that the probability density at the right-hand side of the barrier (i.e.,  $x = x_2$ ) is

$$|\psi_2|^2 = |\psi_1|^2 \exp\left(-2 \int_{x_1}^{x_2} |k(x')| dx'\right). \quad (4.3.11)$$

Note that  $|\psi_2|^2 < |\psi_1|^2$ . Of course, in the region to the right of the barrier (i.e.,  $x > x_2$ ), the probability density takes the constant value  $|\psi_2|^2$ .

We can interpret the ratio of the probability densities to the right and to the left of the potential barrier as the probability,  $|T|^2$ , that a particle incident from the left will tunnel through the barrier and emerge on the other side: that is,

$$|T|^2 = \frac{|\psi_2|^2}{|\psi_1|^2} = \exp\left(-2 \int_{x_1}^{x_2} |k(x')| dx'\right). \quad (4.3.12)$$

(See Section 1.3.) It is easily demonstrated that the probability of a particle incident from the right tunneling through the barrier is the same.

Note that the criterion ([e5.43]) for the validity of the WKB approximation implies that the previous transmission probability is very small. Hence, the WKB approximation only applies to situations in which there is very little chance of a particle tunneling through the potential barrier in question. Unfortunately, the validity criterion ([e5.43]) breaks down completely at the edges of the barrier (i.e., at  $x = x_1$  and  $x_2$ ), because  $k(x) = 0$  at these points. However, it can be demonstrated that the contribution of those regions, around  $x = x_1$  and  $x_2$ , in which the WKB approximation breaks down, to the integral in Equation ([e5.49]) is fairly negligible. Hence, the previous expression for the tunneling probability is a reasonable approximation provided that the incident particle's de Broglie wavelength is much smaller than the spatial extent of the potential barrier.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [4.3: WKB Approximation](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 4.4: Cold Emission

Suppose that an unheated metal surface is subject to a large uniform external electric field of strength  $\mathcal{E}$ , which is directed such that it accelerates electrons away from the surface. We have already seen (in Section [s3.3]) that electrons just below the surface of a metal can be regarded as being in a potential well of depth  $W$ , where  $W$  is called the workfunction of the surface. Adopting a simple one-dimensional treatment of the problem, let the metal lie at  $x < 0$ , and the surface at  $x = 0$ . Now, the applied electric field is shielded from the interior of the metal. Hence, the energy,  $E$ , say, of an electron just below the surface is unaffected by the field. In the absence of the electric field, the potential barrier just above the surface is simply  $V(x) - E = W$ . The electric field modifies this to  $V(x) - E = W - e\mathcal{E}x$ . The potential barrier is sketched in Figure [fcold].




Figure 14: The potential barrier for an electron in a metal surface subject to an external electric field.

It can be seen, from Figure [fcold], that an electron just below the surface of the metal is confined by a triangular potential barrier which extends from  $x = x_1$  to  $x_2$ , where  $x_1 = 0$  and  $x_2 = W/e\mathcal{E}$ . Making use of the WKB approximation (see the previous section), the probability of such an electron tunneling through the barrier, and consequently being emitted from the surface, is

$$|T|^2 = \exp\left(-\frac{2\sqrt{2m}}{\hbar} \int_{x_1}^{x_2} \sqrt{V(x) - E} dx\right), \quad (4.4.1)$$

or

$$|T|^2 = \exp\left(-\frac{2\sqrt{2m}}{\hbar} \int_0^{W/e\mathcal{E}} \sqrt{W - e\mathcal{E}x} dx\right). \quad (4.4.2)$$

This reduces to

$$|T|^2 = \exp\left(-2\sqrt{2} \frac{m^{1/2} W^{3/2}}{\hbar e \mathcal{E}} \int_0^1 \sqrt{1-y} dy\right), \quad (4.4.3)$$

or

$$|T|^2 = \exp\left(-\frac{4\sqrt{2}}{3} \frac{m^{1/2} W^{3/2}}{\hbar e \mathcal{E}}\right). \quad (4.4.4)$$

The previous result is known as the *Fowler-Nordheim formula*. Note that the probability of emission increases exponentially as the electric field-strength above the surface of the metal increases.

The cold emission of electrons from a metal surface is the basis of an important device known as a *scanning tunneling microscope*, or an STM. An STM consists of a very sharp conducting probe which is scanned over the surface of a metal (or any other solid conducting medium). A large voltage difference is applied between the probe and the surface. Now, the surface electric field-strength immediately below the probe tip is proportional to the applied potential difference, and inversely proportional to the



spacing between the tip and the surface. Electrons tunneling between the surface and the probe tip give rise to a weak electric current. The magnitude of this current is proportional to the tunneling probability ([e5.53]). It follows that the current is an extremely sensitive function of the surface electric field-strength, and, hence, of the spacing between the tip and the surface (assuming that the potential difference is held constant). An STM can thus be used to construct a very accurate contour map of the surface under investigation. In fact, STMs are capable of achieving sufficient resolution to image individual atoms

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [4.4: Cold Emission](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 4.5: Alpha Decay

Many types of heavy atomic nucleus spontaneously decay to produce daughter nuclei via the emission of  $\alpha$ -particles (i.e., helium nuclei) of some characteristic energy. This process is known as  $\alpha$ -decay. Let us investigate the  $\alpha$ -decay of a particular type of atomic nucleus of radius  $R$ , charge-number  $Z$ , and mass-number  $A$ . Such a nucleus thus decays to produce a daughter nucleus of charge-number  $Z_1 = Z - 2$  and mass-number  $A_1 = A - 4$ , and an  $\alpha$ -particle of charge-number  $Z_2 = 2$  and mass-number  $A_2 = 4$ . Let the characteristic energy of the  $\alpha$ -particle be  $E$ . Incidentally, nuclear radii are found to satisfy the empirical formula

$$R = 1.5 \times 10^{-15} A^{1/3} \text{ m} = 2.0 \times 10^{-15} Z_1^{1/3} \text{ m} \quad (4.5.1)$$

for  $Z \gg 1$ .

In 1928, George Gamow proposed a very successful theory of  $\alpha$ -decay, according to which the  $\alpha$ -particle moves freely inside the nucleus, and is emitted after tunneling through the potential barrier between itself and the daughter nucleus. In other words, the  $\alpha$ -particle, whose energy is  $E$ , is trapped in a potential well of radius  $R$  by the potential barrier

$$V(r) = \frac{Z_1 Z_2 e^2}{4\pi \epsilon_0 r} \quad (4.5.2)$$

for  $r > R$ .

Making use of the WKB approximation (and neglecting the fact that  $r$  is a radial, rather than a Cartesian, coordinate), the probability of the  $\alpha$ -particle tunneling through the barrier is

$$|T|^2 = \exp\left(-\frac{2\sqrt{2m}}{\hbar} \int_{r_1}^{r_2} \sqrt{V(r) - E} dr\right), \quad (4.5.3)$$

where  $r_1 = R$  and  $r_2 = Z_1 Z_2 e^2 / (4\pi \epsilon_0 E)$ . Here,  $m = 4 m_p$  is the  $\alpha$ -particle mass. The previous expression reduces to

$$|T|^2 = \exp\left(-2\sqrt{2}\beta \int_1^{E_c/E} \left[\frac{1}{y} - \frac{E}{E_c}\right]^{1/2} dy\right), \quad (4.5.4)$$

where

$$\beta = \left(\frac{Z_1 Z_2 e^2 m R}{4\pi \epsilon_0 \hbar^2}\right)^{1/2} = 0.74 Z_1^{2/3} \quad (4.5.5)$$

is a dimensionless constant, and

$$E_c = \frac{Z_1 Z_2 e^2}{4\pi \epsilon_0 R} = 1.44 Z_1^{2/3} \text{ MeV} \quad (4.5.6)$$

is the characteristic energy the  $\alpha$ -particle would need in order to escape from the nucleus without tunneling. Of course,  $E \ll E_c$ . It is easily demonstrated that

$$\int_1^{1/\epsilon} \left(\frac{1}{y} - \epsilon\right)^{1/2} dy \simeq \frac{\pi}{2\sqrt{\epsilon}} - 2 \quad (4.5.7)$$

when  $\epsilon \ll 1$ . Hence,

$$|T|^2 \simeq \exp\left(-2\sqrt{2}\beta \left[\frac{\pi}{2} \sqrt{\frac{E_c}{E}} - 2\right]\right). \quad (4.5.8)$$

Now, the  $\alpha$ -particle moves inside the nucleus with the characteristic velocity  $v = \sqrt{2E/m}$ . It follows that the particle bounces backward and forward within the nucleus at the frequency  $\nu \simeq v/R$ , giving

$$\nu \simeq 2 \times 10^{28} \text{ yr}^{-1} \quad (4.5.9)$$

for a 1 MeV  $\alpha$ -particle trapped inside a typical heavy nucleus of radius  $10^{-14}$  m. Thus, the  $\alpha$ -particle effectively attempts to tunnel through the potential barrier  $\nu$  times a second. If each of these attempts has a probability  $|T|^2$  of succeeding then the probability of

decay per unit time is  $\nu |T|^2$ . Hence, if there are  $N(t) \gg 1$  undecayed nuclei at time  $t$  then there are only  $N + dN$  at time  $t + dt$ , where

$$dN = -N \nu |T|^2 dt. \quad (4.5.10)$$

This expression can be integrated to give

$$N(t) = N(0) \exp(-\nu |T|^2 t). \quad (4.5.11)$$

Now, the *half-life*,  $\tau$ , is defined as the time which must elapse in order for half of the nuclei originally present to decay. It follows from the previous formula that

$$\tau = \frac{\ln 2}{\nu |T|^2}. \quad (4.5.12)$$

Note that the half-life is independent of  $N(0)$ .

Finally, making use of the previous results, we obtain

$$\log_{10}[\tau(\text{yr})] = -C_1 - C_2 Z_1^{2/3} + C_3 \frac{Z_1}{\sqrt{E(\text{MeV})}}, \quad (4.5.13)$$

where

$$C_1 = 28.5,$$

$$C_2 = 1.83,$$

$$C_3 = 1.73.$$




Figure 15: The experimentally determined half-life,  $\tau_{\text{ex}}$  of various atomic nuclei which decay via  $\alpha$  emission versus the best-fit theoretical half-life  $\log_{10}(\tau_{\text{th}}) = -28.9 - 1.60 Z_1^{2/3} + 1.61 Z_1 / \sqrt{E}$ . Both half-lives are measured in years. Here,  $Z_1 = Z - 2$ . Both half-lives are measured in years. Here,  $Z_1 = Z - 2$ , where  $Z$  is the charge number of the nucleus, and  $E$  the characteristic energy of the emitted  $\alpha$ -particle in MeV. In order of increasing half-life, the points correspond to the following nuclei: Rn 215, Po 214, Po 216, Po 197, Fm 250, Ac 225, U 230, U 232, U 234, Gd 150, U 236, U 238, Pt 190, Gd 152, Nd 144. Data obtained from IAEA Nuclear Data Centre.

Equation (4.5.64) is known as the *Geiger-Nuttall formula*, because it was discovered empirically by H. Geiger and J.M. Nuttall in 1911.

The half-life,  $\tau$ , the daughter charge-number,  $Z_1 = Z - 2$ , and the  $\alpha$ -particle energy,  $E$ , for atomic nuclei which undergo  $\alpha$ -decay are indeed found to satisfy a relationship of the form ([e5.64](#)). The best fit to the data (see Figure [fal](#)) is obtained using

$$C_1 = 28.9,$$

$$C_2 = 1.60,$$

$$C_3 = 1.61.$$

Note that these values are remarkably similar to those calculated previously.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [4.5: Alpha Decay](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 4.6: Square Potential Well

Consider a particle of mass  $m$  and energy  $E$  interacting with the simple square potential well

$$V(x) = \begin{cases} -V_0 & \text{for } -a/2 \leq x \leq a/2 \\ 0 & \text{otherwise} \end{cases} \quad (4.6.1)$$

Now, if  $E > 0$  then the particle is unbounded. Thus, when the particle encounters the well it is either reflected or transmitted. As is easily demonstrated, the reflection and transmission probabilities are given by Equations ([e5.28]) and ([e5.29]), respectively, where

$$k^2 = \frac{2mE}{\hbar^2},$$

$$q^2 = \frac{2m(E + V_0)}{\hbar^2}.$$

Suppose, however, that  $E < 0$ . In this case, the particle is bounded (i.e.,  $|\psi|^2 \rightarrow 0$  as  $|x| \rightarrow \infty$ ). Is it possible to find bounded solutions of Schrödinger's equation in the finite square potential well ([e5.71])?

Now, it is easily seen that independent solutions of Schrödinger's equation ([e5.2]) in the symmetric [i.e.,  $V(-x) = V(x)$ ] potential ([e5.71]) must be either totally symmetric [i.e.,  $\psi(-x) = \psi(x)$ ], or totally anti-symmetric [i.e.,  $\psi(-x) = -\psi(x)$ ]. Moreover, the solutions must satisfy the boundary condition

$$\psi \rightarrow 0 \quad \text{as } |x| \rightarrow \infty \quad (4.6.2)$$

Let us, first of all, search for a totally symmetric solution. In the region to the left of the well (i.e.  $x < -a/2$ ), the solution of Schrödinger's equation which satisfies the boundary condition  $\psi \rightarrow 0$  and  $x \rightarrow -\infty$  is

$$\psi(x) = A e^{kx}, \quad (4.6.3)$$

where

$$k^2 = \frac{2m|E|}{\hbar^2}. \quad (4.6.4)$$

By symmetry, the solution in the region to the right of the well (i.e.,  $x > a/2$ ) is

$$\psi(x) = A e^{-kx}. \quad (4.6.5)$$

The solution inside the well (i.e.,  $|x| \leq a/2$ ) which satisfies the symmetry constraint  $\psi(-x) = \psi(x)$  is

$$\psi(x) = B \cos(qx), \quad (4.6.6)$$

where

$$q^2 = \frac{2m(V_0 + E)}{\hbar^2}. \quad (4.6.7)$$

Here, we have assumed that  $E > -V_0$ . The constraint that  $\psi(x)$  and its first derivative be continuous at the edges of the well (i.e., at  $x = \pm a/2$ ) yields

$$k = q \tan(qa/2). \quad (4.6.8)$$

Let  $y = qa/2$ . It follows that

$$E = E_0 y^2 - V_0, \quad (4.6.9)$$

where

$$E_0 = \frac{2\hbar^2}{ma^2}. \quad (4.6.10)$$

Moreover, Equation ([e5.81]) becomes

$$\frac{\sqrt{\lambda - y^2}}{y} = \tan y, \quad (4.6.11)$$

with

$$\lambda = \frac{V_0}{E_0}. \quad (4.6.12)$$

Here,  $y$  must lie in the range  $0 < y < \sqrt{\lambda}$ : that is,  $E$  must lie in the range  $-V_0 < E < 0$ .

Now, the solutions to Equation (e5.84) correspond to the intersection of the curve  $\sqrt{\lambda - y^2}/y$  with the curve  $\tan y$ . Figure [well] shows these two curves plotted for a particular value of  $\lambda$ . In this case, the curves intersect twice, indicating the existence of two totally symmetric bound states in the well. Moreover, it is evident, from the figure, that as  $\lambda$  increases (i.e., as the well becomes deeper) there are more and more bound states. However, it is also evident that there is always at least one totally symmetric bound state, no matter how small  $\lambda$  becomes (i.e., no matter how shallow the well becomes). In the limit  $\lambda \gg 1$  (i.e., the limit in which the well becomes very deep), the solutions to Equation (e5.84) asymptote to the roots of  $\tan y = \infty$ . This gives  $y = (2j - 1)\pi/2$ , where  $j$  is a positive integer, or

$$q = \frac{(2j - 1)\pi}{a}. \quad (4.6.13)$$

These solutions are equivalent to the odd- $n$  infinite square well solutions specified by Equation (e5.8).

For the case of a totally anti-symmetric bound state, similar analysis to the preceding yields

$$-\frac{y}{\sqrt{\lambda - y^2}} = \tan y. \quad (4.6.14)$$

The solutions of this equation correspond to the intersection of the curve  $\tan y$  with the curve  $-y/\sqrt{\lambda - y^2}$ . Figure [well1] shows these two curves plotted for the same value of  $\lambda$  as that used in Figure [well]. In this case, the curves intersect once, indicating the existence of a single totally anti-symmetric bound state in the well. It is, again, evident, from the figure, that as  $\lambda$  increases (i.e., as the well becomes deeper) there are more and more bound states. However, it is also evident that when  $\lambda$  becomes sufficiently small [i.e.,  $\lambda < (\pi/2)^2$ ] then there is no totally anti-symmetric bound state. In other words, a very shallow potential well always possesses a totally symmetric bound state, but does not generally possess a totally anti-symmetric bound state. In the limit  $\lambda \gg 1$  (i.e., the limit in which the well becomes very deep), the solutions to Equation (e5.85) asymptote to the roots of  $\tan y = 0$ . This gives  $y = j\pi$ , where  $j$  is a positive integer, or

$$q = \frac{2j\pi}{a}. \quad (4.6.15)$$

These solutions are equivalent to the even- $n$  infinite square well solutions specified by Equation (e5.8).

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled 4.6: Square Potential Well is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 4.7: Simple Harmonic Oscillator

The classical Hamiltonian of a simple harmonic oscillator is

$$H = \frac{p^2}{2m} + \frac{1}{2} K x^2, \quad (4.7.1)$$

where  $K > 0$  is the so-called force constant of the oscillator. Assuming that the quantum mechanical Hamiltonian has the same form as the classical Hamiltonian, the time-independent Schrödinger equation for a particle of mass  $m$  and energy  $E$  moving in a simple harmonic potential becomes

$$\frac{d^2 \psi}{dx^2} = \frac{2m}{\hbar^2} \left( \frac{1}{2} K x^2 - E \right) \psi. \quad (4.7.2)$$

Let  $\omega = \sqrt{K/m}$ , where  $\omega$  is the oscillator's classical angular frequency of oscillation. Furthermore, let

$$y = \sqrt{\frac{m\omega}{\hbar}} x, \quad (4.7.3)$$

and

$$\epsilon = \frac{2E}{\hbar\omega}. \quad (4.7.4)$$

Equation (4.7.2) reduces to

$$\frac{d^2 \psi}{dy^2} - (y^2 - \epsilon) \psi = 0. \quad (4.7.5)$$

We need to find solutions to the previous equation which are bounded at infinity: that is, solutions which satisfy the boundary condition  $\psi \rightarrow 0$  as  $|y| \rightarrow \infty$ .

Consider the behavior of the solution to Equation (4.7.5) in the limit  $|y| \gg 1$ . As is easily seen, in this limit the equation simplifies somewhat to give

$$\frac{d^2 \psi}{dy^2} - y^2 \psi \simeq 0. \quad (4.7.6)$$

The approximate solutions to the previous equation are

$$\psi(y) \simeq A(y) e^{\pm y^2/2}, \quad (4.7.7)$$

where  $A(y)$  is a relatively slowly varying function of  $y$ . Clearly, if  $\psi(y)$  is to remain bounded as  $|y| \rightarrow \infty$  then we must choose the exponentially decaying solution. This suggests that we should write

$$\psi(y) = h(y) e^{-y^2/2}, \quad (4.7.8)$$

where we would expect  $h(y)$  to be an algebraic, rather than an exponential, function of  $y$ .

Substituting Equation (4.7.8) into Equation (4.7.5), we obtain

$$\frac{d^2 h}{dy^2} - 2y \frac{dh}{dy} + (\epsilon - 1) h = 0. \quad (4.7.9)$$

Let us attempt a power-law solution of the form

$$h(y) = \sum_{i=0, \infty} c_i y^i. \quad (4.7.10)$$

Inserting this test solution into Equation (4.7.9), and equating the coefficients of  $y^i$ , we obtain the recursion relation

$$c_{i+2} = \frac{(2i - \epsilon + 1)}{(i+1)(i+2)} c_i. \quad (4.7.11)$$

Consider the behavior of  $h(y)$  in the limit  $|y| \rightarrow \infty$ . The previous recursion relation simplifies to

$$c_{i+2} \simeq \frac{2}{i} c_i. \quad (4.7.12)$$

Hence, at large  $|y|$ , when the higher powers of  $y$  dominate, we have

$$h(y) \sim C \sum_j \frac{y^{2j}}{j!} \sim C e^{y^2}. \quad (4.7.13)$$

It follows that  $\psi(y) = h(y) \exp(-y^2/2)$  varies as  $\exp(y^2/2)$  as  $|y| \rightarrow \infty$ . This behavior is unacceptable, because it does not satisfy the boundary condition  $\psi \rightarrow 0$  as  $|y| \rightarrow \infty$ . The only way in which we can prevent  $\psi$  from blowing up as  $|y| \rightarrow \infty$  is to demand that the power series ([e5.98]) terminate at some finite value of  $i$ . This implies, from the recursion relation ([e5.99]), that

$$\epsilon = 2n + 1, \quad (4.7.14)$$

where  $n$  is a non-negative integer. Note that the number of terms in the power series ([e5.98]) is  $n + 1$ . Finally, using Equation ([e5.92]), we obtain

$$E = (n + 1/2) \hbar \omega, \quad (4.7.15)$$

for  $n = 0, 1, 2, \dots$ .

Hence, we conclude that a particle moving in a harmonic potential has quantized energy levels that are equally spaced. The spacing between successive energy levels is  $\hbar \omega$ , where  $\omega$  is the classical oscillation frequency. Furthermore, the lowest energy state ( $n = 0$ ) possesses the finite energy  $(1/2) \hbar \omega$ . This is sometimes called *zero-point energy*. It is easily demonstrated that the (normalized) wavefunction of the lowest energy state takes the form

$$\psi_0(x) = \frac{e^{-x^2/2d^2}}{\pi^{1/4} \sqrt{d}} \quad (4.7.16)$$

Let  $\psi_n(x)$  be an energy eigenstate of the harmonic oscillator corresponding to the eigenvalue

$$E_n = (n + 1/2) \hbar \omega. \quad (4.7.17)$$

Assuming that the  $\psi_n$  are properly normalized (and real), we have

$$\int_{-\infty}^{\infty} \psi_n \psi_m dx = \delta_{nm}. \quad (4.7.18)$$

Now, Equation ([e5.93]) can be written

$$\left( -\frac{d^2}{dy^2} + y^2 \right) \psi_n = (2n + 1) \psi_n, \quad (4.7.19)$$

where  $x = d y$ , and  $d = \sqrt{\hbar/m\omega}$ . It is helpful to define the operators

$$a_{\pm} = \frac{1}{\sqrt{2}} \left( \mp \frac{d}{dy} + y \right). \quad (4.7.20)$$

As is easily demonstrated, these operators satisfy the commutation relation

$$[a_+, a_-] = -1. \quad (4.7.21)$$

Using these operators, Equation ([e5.108]) can also be written in the forms

$$a_+ a_- \psi_n = n \psi_n, \quad (4.7.22)$$

or

$$a_- a_+ \psi_n = (n + 1) \psi_n. \quad (4.7.23)$$

The previous two equations imply that



$$\begin{aligned} a_+ \psi_n &= \sqrt{n+1} \psi_{n+1}, \\ a_- \psi_n &= \sqrt{n} \psi_{n-1}. \end{aligned}$$

We conclude that  $a_+$  and  $a_-$  are *raising and lowering operators*, respectively, for the harmonic oscillator: that is, operating on the wavefunction with  $a_+$  causes the quantum number  $n$  to increase by unity, and vice versa. The Hamiltonian for the harmonic oscillator can be written in the form

$$H = \hbar \omega \left( a_+ a_- + \frac{1}{2} \right), \quad (4.7.24)$$

from which the result

$$H \psi_n = (n + 1/2) \hbar \omega \psi_n = E_n \psi_n \quad (4.7.25)$$

is readily deduced. Finally, Equations ([e5.107]), ([e5.113]), and ([e5.114]) yield the useful expression

$$\int_{-\infty}^{\infty} \psi_m x \psi_n dx = \frac{d}{\sqrt{2}} \int_{-\infty}^{\infty} \psi_m (a_+ + a_-) \psi_n dx = \sqrt{\frac{\hbar}{2m\omega}} (\sqrt{m} \delta_{m,n+1} + \sqrt{n} \delta_{m,n-1}).$$

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [4.7: Simple Harmonic Oscillator](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 4.E: One-Dimensional Potentials (Exercises)

1. Show that the wavefunction of a particle of mass  $m$  in an infinite one-dimensional square-well of width  $a$  returns to its original form after a quantum revival time  $T = 4 m a^2 / \pi \hbar$ .
2. A particle of mass  $m$  moves freely in one dimension between impenetrable walls located at  $x = 0$  and  $a$ . Its initial wavefunction is

$$\psi(x, 0) = \sqrt{2/a} \sin(3\pi x/a). \quad (4.E.1)$$

What is the subsequent time evolution of the wavefunction? Suppose that the initial wavefunction is

$$\psi(x, 0) = \sqrt{1/a} \sin(\pi x/a) [1 + 2 \cos(\pi x/a)]. \quad (4.E.2)$$

What now is the subsequent time evolution? Calculate the probability of finding the particle between 0 and  $a/2$  as a function of time in each case.

3. A particle of mass  $m$  is in the ground-state of an infinite one-dimensional square-well of width  $a$ . Suddenly the well expands to twice its original size, as the right wall moves from  $a$  to  $2a$ , leaving the wavefunction momentarily undisturbed. The energy of the particle is now measured. What is the most probable result? What is the probability of obtaining this result? What is the next most probable result, and what is its probability of occurrence? What is the expectation value of the energy?
4. A stream of particles of mass  $m$  and energy  $E > 0$  encounter a potential step of height  $W (< E)$ : that is,  $V(x) = 0$  for  $x < 0$  and  $V(x) = W$  for  $x > 0$  with the particles incident from  $-\infty$ . Show that the fraction reflected is

$$R = \left( \frac{k - q}{k + q} \right)^2, \quad (4.E.3)$$

where  $k^2 = (2m/\hbar^2) E$  and  $q^2 = (2m/\hbar^2) (E - W)$ .

5. A stream of particles of mass  $m$  and energy  $E > 0$  encounter the delta-function potential  $V(x) = -\alpha \delta(x)$ , where  $\alpha > 0$ . Show that the fraction reflected is

$$R = \beta^2 / (1 + \beta^2), \quad (4.E.4)$$

where  $\beta = m\alpha/\hbar^2 k$ , and  $k^2 = (2m/\hbar^2) E$ . Does such a potential have a bound state? If so, what is its energy?

6. Two potential wells of width  $a$  are separated by a distance  $L \gg a$ . A particle of mass  $m$  and energy  $E$  is in one of the wells. Estimate the time required for the particle to tunnel to the other well.
7. Consider a particle trapped in the finite potential well whose potential is given by Equation ([e5.71]). Demonstrate that for a totally-symmetric state the ratio of the probability of finding the particle outside to the probability of finding the particle inside the well is

$$\frac{P_{\text{out}}}{P_{\text{in}}} = \frac{\cos^3 y}{\sin y (y + \sin y \cos y)}, \quad (4.E.5)$$

where  $(\lambda - y^2)^{1/2} = y \tan y$ , and  $\lambda = V/E_0$ . Hence, demonstrate that for a shallow well (i.e.,  $\lambda \ll 1$ )  $P_{\text{out}} \simeq 1 - 2\lambda$ , whereas for a deep well (i.e.,  $\lambda \gg 1$ )  $P_{\text{out}} \simeq (\pi^2/4)/\lambda^{3/2}$  (assuming that the particle is in the ground state).[ex12.3]

8. Consider the half-infinite potential well

$$V(x) = \begin{cases} \infty & x \leq 0 \\ -V_0 & 0 < x < L \\ 0 & x \geq L \end{cases}, \quad (4.E.6)$$

where  $V_0 > 0$ . Demonstrate that the bound-states of a particle of mass  $m$  and energy  $-V_0 < E < 0$  satisfy

$$\tan\left(\sqrt{2m(V_0 + E)} L/\hbar\right) = -\sqrt{(V_0 + E)/(-E)}. \quad (4.E.7)$$

9. Find the properly normalized first two excited energy eigenstates of the harmonic oscillator, as well as the expectation value of the potential energy in the  $n$ th energy eigenstate. Hint: Consider the raising and lowering operators  $a_{\pm}$ , defined in Equation ([e5.109]).

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [4.E: One-Dimensional Potentials \(Exercises\)](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## CHAPTER OVERVIEW

### 5: Multi-Particle Systems

In this chapter, we shall extend the single particle, one-dimensional formulation of non-relativistic quantum mechanics, introduced in the previous chapters, in order to investigate one-dimensional systems containing multiple particles.

[5.1: Fundamental Concepts of Multi-Particle Systems](#)

[5.2: Non-interacting Particles](#)

[5.3: Two-Particle Systems](#)

[5.4: Identical Particles](#)

[5.E: Multi-Particle Systems \(Exercises\)](#)

---

This page titled [5: Multi-Particle Systems](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 5.1: Fundamental Concepts of Multi-Particle Systems

We have already seen that the instantaneous state of a system consisting of a single non-relativistic particle, whose position coordinate is  $x$ , is fully specified by a complex wavefunction  $\psi(x, t)$ . This wavefunction is interpreted as follows. The probability of finding the particle between  $x$  and  $x + dx$  at time  $t$  is given by  $|\psi(x, t)|^2 dx$ . This interpretation only makes sense if the wavefunction is normalized such that

$$\int_{-\infty}^{\infty} |\psi(x, t)|^2 dx = 1 \quad (5.1.1)$$

at all times. The physical significance of this normalization requirement is that the probability of the particle being found anywhere on the  $x$ -axis must always be unity (which corresponds to certainty).

Consider a system containing  $N$  non-relativistic particles, labeled  $i = 1, N$ , moving in one dimension. Let  $x_i$  and  $m_i$  be the position coordinate and mass, respectively, of the  $i$ th particle. By analogy with the single-particle case, the instantaneous state of a multi-particle system is specified by a complex wavefunction  $\psi(x_1, x_2, \dots, x_N, t)$ . The probability of finding the first particle between  $x_1$  and  $x_1 + dx_1$ , the second particle between  $x_2$  and  $x_2 + dx_2$ , et cetera, at time  $t$  is given by  $|\psi(x_1, x_2, \dots, x_N, t)|^2 dx_1 dx_2 \dots dx_N$ . It follows that the wavefunction must satisfy the normalization condition

$$\int |\psi(x_1, x_2, \dots, x_N, t)|^2 dx_1 dx_2 \dots dx_N = 1 \quad (5.1.2)$$

at all times, where the integration is taken over all  $x_1 x_2 \dots x_N$  space.

In a single-particle system, position is represented by the algebraic operator  $x$ , whereas momentum is represented by the differential operator  $-i\hbar \partial/\partial x$ . (See Section [s4.6].) By analogy, in a multi-particle system, the position of the  $i$ th particle is represented by the algebraic operator  $x_i$ , whereas the corresponding momentum is represented by the differential operator

$$p_i = -i\hbar \frac{\partial}{\partial x_i}. \quad (5.1.3)$$

Because the  $x_i$  are independent variables (i.e.,  $\partial x_i / \partial x_j = \delta_{ij}$ ), we conclude that the various position and momentum operators satisfy the following commutation relations:

$$\begin{aligned} [x_i, x_j] &= 0, \\ [p_i, p_j] &= 0, \\ [x_i, p_j] &= i\hbar \delta_{ij}. \end{aligned}$$

Now, we know, from Section [smeas], that two dynamical variables can only be (exactly) measured simultaneously if the operators that represent them in quantum mechanics commute with one another. Thus, it is clear, from the previous commutation relations, that the only restriction on measurement in a one-dimensional multi-particle system is that it is impossible to simultaneously measure the position and momentum of the same particle. Note, in particular, that a knowledge of the position or momentum of a given particle does not in any way preclude a similar knowledge for a different particle. The commutation relations ([xe6.4])–([xe6.6]) illustrate an important point in quantum mechanics: namely, that operators corresponding to different degrees of freedom of a dynamical system tend to commute with one another. In this case, the different degrees of freedom correspond to the different motions of the various particles making up the system.

Finally, if  $H(x_1, x_2, \dots, x_N, t)$  is the Hamiltonian of the system then the multi-particle wavefunction  $\psi(x_1, x_2, \dots, x_N, t)$  satisfies the usual time-dependent Schrödinger equation [see Equation ([etimed])]

$$i\hbar \frac{\partial \psi}{\partial t} = H \psi. \quad (5.1.4)$$

Likewise, a multi-particle state of definite energy  $E$  (i.e., an eigenstate of the Hamiltonian with eigenvalue  $E$ ) is written (see Section [sstat])

$$\psi(x_1, x_2, \dots, x_N, t) = \psi_E(x_1, x_2, \dots, x_N) e^{-iEt/\hbar}, \quad (5.1.5)$$

where the stationary wavefunction  $\psi_E$  satisfies the time-independent Schrödinger equation [see Equation ([etimei])]

$$H \psi_E = E \psi_E. \quad (5.1.6)$$

Here,  $H$  is assumed not to be an explicit function of  $t$ .

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [5.1: Fundamental Concepts of Multi-Particle Systems](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 5.2: Non-interacting Particles

In general, we expect the Hamiltonian of a multi-particle system to take the form

$$H(x_1, x_2, \dots, x_N, t) = \sum_{i=1, N} \frac{p_i^2}{2m_i} + V(x_1, x_2, \dots, x_N, t). \quad (5.2.1)$$

Here, the first term on the right-hand side represents the total kinetic energy of the system, whereas the potential  $V$  specifies the nature of the interaction between the various particles making up the system, as well as the interaction of the particles with any external forces.

Suppose that the particles do not interact with one another. This implies that each particle moves in a common potential: that is,

$$V(x_1, x_2, \dots, x_N, t) = \sum_{i=1, N} V(x_i, t). \quad (5.2.2)$$

Hence, we can write

$$H(x_1, x_2, \dots, x_N, t) = \sum_{i=1, N} H_i(x_i, t), \quad (5.2.3)$$

where

$$H_i = \frac{p_i^2}{2m_i} + V(x_i, t). \quad (5.2.4)$$

In other words, for the case of non-interacting particles, the multi-particle Hamiltonian of the system can be written as the sum of  $N$  independent single-particle Hamiltonians. Here,  $H_i$  represents the energy of the  $i$ th particle, and is completely unaffected by the energies of the other particles. Furthermore, given that the various particles that make up the system are non-interacting, we expect their instantaneous positions to be completely uncorrelated with one another. This immediately implies that the multi-particle wavefunction  $\psi(x_1, x_2, \dots, x_N, t)$  can be written as the product of  $N$  independent single-particle wavefunctions: that is,

$$\psi(x_1, x_2, \dots, x_N, t) = \psi_1(x_1, t) \psi_2(x_2, t) \dots \psi_N(x_N, t). \quad (5.2.5)$$

Here,  $|\psi_i(x_i, t)|^2 dx_i$  is the probability of finding the  $i$ th particle between  $x_i$  and  $x_i + dx_i$  at time  $t$ . This probability is completely unaffected by the positions of the other particles. It is evident that  $\psi_i(x_i, t)$  must satisfy the normalization constraint

$$\int_{-\infty}^{\infty} |\psi_i(x_i, t)|^2 dx_i = 1. \quad (5.2.6)$$

If this is the case then the normalization constraint ([\[ex1\]](#)) for the multi-particle wavefunction is automatically satisfied. Equation ([\[ex13\]](#)) illustrates an important point in quantum mechanics: namely, that we can generally write the total wavefunction of a many degree of freedom system as a product of different wavefunctions corresponding to each degree of freedom.

According to Equations ([\[ex11\]](#)) and ([\[ex13\]](#)), the time-dependent Schrödinger equation ([\[ex7\]](#)) for a system of  $N$  non-interacting particles factorizes into  $N$  independent equations of the form

$$i\hbar \frac{\partial \psi_i}{\partial t} = H_i \psi_i. \quad (5.2.7)$$

Assuming that  $V(x, t) \equiv V(x)$ , the time-independent Schrödinger equation ([\[ex9\]](#)) also factorizes to give

$$H_i \psi_{E_i} = E_i \psi_{E_i}, \quad (5.2.8)$$

where  $\psi_i(x_i, t) = \psi_{E_i}(x_i) \exp(-i E_i t / \hbar)$ , and  $E_i$  is the energy of the  $i$ th particle. Hence, a multi-particle state of definite energy  $E$  has a wavefunction of the form

$$\psi(x_1, x_2, \dots, x_n, t) = \psi_E(x_1, x_2, \dots, x_N) e^{-i E t / \hbar}, \quad (5.2.9)$$

where

$$\psi_E(x_1, x_2, \dots, x_N) = \psi_{E_1}(x_1) \psi_{E_2}(x_2) \dots \psi_{E_N}(x_N), \quad (5.2.10)$$

and

$$E = \sum_{i=1, N} E_i. \quad (5.2.11)$$

Clearly, for the case of non-interacting particles, the energy of the whole system is simply the sum of the energies of the component particles.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [5.2: Non-interacting Particles](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).



## 5.3: Two-Particle Systems

Consider a system consisting of two particles, mass  $m_1$  and  $m_2$ , interacting via a potential  $V(x_1 - x_2)$  that only depends on the relative positions of the particles. According to Equations ([ex3]) and ([ex10]), the Hamiltonian of the system is written

$$H(x_1, x_2) = -\frac{\hbar^2}{2m_1} \frac{\partial^2}{\partial x_1^2} - \frac{\hbar^2}{2m_2} \frac{\partial^2}{\partial x_2^2} + V(x_1 - x_2). \quad (5.3.1)$$

Let

$$x' = x_1 - x_2 \quad (5.3.2)$$

be the particles' relative position coordinate, and

$$X = \frac{m_1 x_1 + m_2 x_2}{m_1 + m_2} \quad (5.3.3)$$

the coordinate of the center of mass. It is easily demonstrated that

$$\begin{aligned} \frac{\partial}{\partial x_1} &= \frac{m_1}{m_1 + m_2} \frac{\partial}{\partial X} + \frac{\partial}{\partial x'}, \\ \frac{\partial}{\partial x_2} &= \frac{m_2}{m_1 + m_2} \frac{\partial}{\partial X} - \frac{\partial}{\partial x'}. \end{aligned}$$

Hence, when expressed in terms of the new variables,  $x'$  and  $X$ , the Hamiltonian becomes

$$H(x', X) = -\frac{\hbar^2}{2M} \frac{\partial^2}{\partial X^2} - \frac{\hbar^2}{2\mu} \frac{\partial^2}{\partial x'^2} + V(x'), \quad (5.3.4)$$

where

$$M = m_1 + m_2 \quad (5.3.5)$$

is the total mass of the system, and

$$\mu = \frac{m_1 m_2}{m_1 + m_2} \quad (5.3.6)$$

the so-called *reduced mass*. Note that the total momentum of the system can be written

$$P = -i\hbar \left( \frac{\partial}{\partial x_1} + \frac{\partial}{\partial x_2} \right) = -i\hbar \frac{\partial}{\partial X}. \quad (5.3.7)$$

The fact that the Hamiltonian ([ex6.24]) is separable when expressed in terms of the new coordinates [i.e.,  $H(x', X) = H_{x'}(x') + H_X(X)$ ] suggests, by analogy with the analysis in the previous section, that the wavefunction can be factorized: that is,

$$\psi(x_1, x_2, t) = \psi_{x'}(x', t) \psi_X(X, t). \quad (5.3.8)$$

Hence, the time-dependent Schrödinger equation ([ex7]) also factorizes to give

$$i\hbar \frac{\partial \psi_{x'}}{\partial t} = -\frac{\hbar^2}{2\mu} \frac{\partial^2 \psi_{x'}}{\partial x'^2} + V(x') \psi_{x'}, \quad (5.3.9)$$

and

$$i\hbar \frac{\partial \psi_X}{\partial t} = -\frac{\hbar^2}{2M} \frac{\partial^2 \psi_X}{\partial X^2}. \quad (5.3.10)$$

The previous equation can be solved to give

$$\psi_X(X, t) = \psi_0 e^{i(P'X/\hbar - E't/\hbar)}, \quad (5.3.11)$$

where  $\psi_0$ ,  $P'$ , and  $E' = P'^2/2M$  are constants. It is clear, from Equations ([exa]), ([exb]), and ([ex33]), that the total momentum of the system takes the constant value  $P'$ . In other words, momentum is conserved.

Suppose that we work in the *centre of mass frame* of the system, which is characterized by  $P' = 0$ . It follows that  $\psi_X = \psi_0$ . In this case, we can write the wavefunction of the system in the form  $\psi(x_1, x_2, t) = \psi_{x'}(x', t) \psi_0 \equiv \psi(x_1 - x_2, t)$ , where

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2\mu} \frac{\partial^2 \psi}{\partial x^2} + V(x) \psi. \quad (5.3.12)$$

In other words, in the center of mass frame, two particles of mass  $m_1$  and  $m_2$ , moving in the potential  $V(x_1 - x_2)$ , are equivalent to a single particle of mass  $\mu$ , moving in the potential  $V(x)$ , where  $x = x_1 - x_2$ . This is a familiar result from classical dynamics.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [5.3: Two-Particle Systems](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 5.4: Identical Particles

Consider a system consisting of two identical particles of mass  $m$ . As before, the instantaneous state of the system is specified by the complex wavefunction  $\psi(x_1, x_2, t)$ . This wavefunction tells us that the probability of finding the first particle between  $x_1$  and  $x_1 + dx_1$ , and the second between  $x_2$  and  $x_2 + dx_2$ , at time  $t$  is  $|\psi(x_1, x_2, t)|^2 dx_1 dx_2$ . However, because the particles are identical, this must be the same as the probability of finding the first particle between  $x_2$  and  $x_2 + dx_2$ , and the second between  $x_1$  and  $x_1 + dx_1$ , at time  $t$  (because, in both cases, the result of the measurement is exactly the same). Hence, we conclude that

$$|\psi(x_1, x_2, t)|^2 = |\psi(x_2, x_1, t)|^2, \quad (5.4.1)$$

or

$$\psi(x_1, x_2, t) = e^{i\varphi} \psi(x_2, x_1, t), \quad (5.4.2)$$

where  $\varphi$  is a real constant. However, if we swap the labels on particles 1 and 2 (which are, after all, arbitrary for identical particles), and repeat the argument, we also conclude that

$$\psi(x_2, x_1, t) = e^{i\varphi} \psi(x_1, x_2, t). \quad (5.4.3)$$

Hence,

$$e^{2i\varphi} = 1. \quad (5.4.4)$$

The only solutions to the previous equation are  $\varphi = 0$  and  $\varphi = \pi$ . Thus, we infer that, for a system consisting of two identical particles, the wavefunction must be either symmetric or anti-symmetric under interchange of particle labels. That is, either

$$\psi(x_2, x_1, t) = \psi(x_1, x_2, t), \quad (5.4.5)$$

or

$$\psi(x_2, x_1, t) = -\psi(x_1, x_2, t). \quad (5.4.6)$$

The previous argument can easily be extended to systems containing more than two identical particles.

It turns out that the question of whether the wavefunction of a system containing many identical particles is symmetric or anti-symmetric under interchange of the labels on any two particles is determined by the nature of the particles themselves. Particles with wavefunctions that are symmetric under label interchange are said to obey *Bose-Einstein statistics*, and are called *bosons*. For instance, photons are bosons. Particles with wavefunctions that are anti-symmetric under label interchange are said to obey *Fermi-Dirac statistics*, and are called *fermions*. For instance, electrons, protons, and neutrons are fermions.

Consider a system containing two identical and non-interacting bosons. Let  $\psi(x, E)$  be a properly normalized, single-particle, stationary wavefunction corresponding to a state of definite energy  $E$ . The stationary wavefunction of the whole system is written

$$\psi_{E \text{ boson}}(x_1, x_2) = \frac{1}{\sqrt{2}} [\psi(x_1, E_a) \psi(x_2, E_b) + \psi(x_2, E_a) \psi(x_1, E_b)], \quad (5.4.7)$$

when the energies of the two particles are  $E_a$  and  $E_b$ . This expression automatically satisfies the symmetry requirement on the wavefunction. Incidentally, because the particles are identical, we cannot be sure which particle has energy  $E_a$ , and which has energy  $E_b$ —only that one particle has energy  $E_a$ , and the other  $E_b$ .

For a system consisting of two identical and non-interacting fermions, the stationary wavefunction of the whole system takes the form

$$\psi_{E \text{ fermion}}(x_1, x_2) = \frac{1}{\sqrt{2}} [\psi(x_1, E_a) \psi(x_2, E_b) - \psi(x_2, E_a) \psi(x_1, E_b)], \quad (5.4.8)$$

Again, this expression automatically satisfies the symmetry requirement on the wavefunction. Note that if  $E_a = E_b$  then the total wavefunction becomes zero everywhere. Now, in quantum mechanics, a null wavefunction corresponds to the absence of a state. We thus conclude that it is impossible for the two fermions in our system to occupy the same single-particle stationary state.

Finally, if the two particles are somehow distinguishable then the stationary wavefunction of the system is simply

$$\psi_{E \text{ dist}}(x_1, x_2) = \psi(x_1, E_a) \psi(x_2, E_b). \quad (5.4.9)$$

Let us evaluate the variance of the distance,  $x_1 - x_2$ , between the two particles, using the previous three wavefunctions. It is easily demonstrated that if the particles are distinguishable then

$$\langle (x_1 - x_2)^2 \rangle_{\text{dist}} = \langle x^2 \rangle_a + \langle x^2 \rangle_b - 2 \langle x \rangle_a \langle x \rangle_b, \quad (5.4.10)$$

where

$$\langle x^n \rangle_{a,b} = \int_{-\infty}^{\infty} \psi^*(x, E_{a,b}) x^n \psi(x, E_{a,b}) dx. \quad (5.4.11)$$

For the case of two identical bosons, we find

$$\langle (x_1 - x_2)^2 \rangle_{\text{boson}} = \langle (x_1 - x_2)^2 \rangle_{\text{dist}} - 2 |\langle x \rangle_{ab}|^2, \quad (5.4.12)$$

where

$$\langle x \rangle_{ab} = \int_{-\infty}^{\infty} \psi^*(x, E_a) x \psi(x, E_b) dx. \quad (5.4.13)$$

Here, we have assumed that  $E_a \neq E_b$ , so that

$$\int_{-\infty}^{\infty} \psi^*(x, E_a) \psi(x, E_b) dx = 0. \quad (5.4.14)$$

Finally, for the case of two identical fermions, we obtain

$$\langle (x_1 - x_2)^2 \rangle_{\text{fermion}} = \langle (x_1 - x_2)^2 \rangle_{\text{dist}} + 2 |\langle x \rangle_{ab}|^2, \quad (5.4.15)$$

Equation 5.4.12 indicates that the symmetry requirement on the total wavefunction of two identical bosons causes the particles to be, on average, closer together than two similar distinguishable particles. Conversely, Equation 5.4.15 indicates that the symmetry requirement on the total wavefunction of two identical fermions causes the particles to be, on average, further apart than two similar distinguishable particles. However, the strength of this effect depends on square of the magnitude of  $\langle x \rangle_{ab}$ , which measures the overlap between the wavefunctions  $\psi(x, E_a)$  and  $\psi(x, E_b)$ . It is evident, then, that if these two wavefunctions do not overlap to any great extent then identical bosons or fermions will act very much like distinguishable particles.

For a system containing  $N$  identical and non-interacting fermions, the anti-symmetric stationary wavefunction of the system is written

$$\psi_E(x_1, x_2, \dots, x_N) = \frac{1}{\sqrt{N!}} \begin{vmatrix} \psi(x_1, E_1) & \psi(x_2, E_1) & \dots & \psi(x_N, E_1) \\ \psi(x_1, E_2) & \psi(x_2, E_2) & \dots & \psi(x_N, E_2) \\ \vdots & \vdots & \ddots & \vdots \\ \psi(x_1, E_N) & \psi(x_2, E_N) & \dots & \psi(x_N, E_N) \end{vmatrix}. \quad (5.4.16)$$

This expression is known as the *Slater determinant*, and automatically satisfies the symmetry requirements on the wavefunction. Here, the energies of the particles are  $E_1, E_2, \dots, E_N$ . Note, again, that if any two particles in the system have the same energy (i.e., if  $E_i = E_j$  for some  $i \neq j$ ) then the total wavefunction is null. We conclude that it is impossible for any two identical fermions in a multi-particle system to occupy the same single-particle stationary state. This important result is known as the *Pauli exclusion principle*.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled 5.4: Identical Particles is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 5.E: Multi-Particle Systems (Exercises)

---

1. Consider a system consisting of two non-interacting particles, and three one-particle states,  $\psi_a(x)$ ,  $\psi_b(x)$ , and  $\psi_c(x)$ . How many different two-particle states can be constructed if the particles are (a) distinguishable, (b) indistinguishable bosons, or (c) indistinguishable fermions?
2. Consider two non-interacting particles, each of mass  $m$ , in a one-dimensional harmonic oscillator potential of classical oscillation frequency  $\omega$ . If one particle is in the ground-state, and the other in the first excited state, calculate  $\langle (x_1 - x_2)^2 \rangle$  assuming that the particles are (a) distinguishable, (b) indistinguishable bosons, or (c) indistinguishable fermions.
3. Two non-interacting particles, with the same mass  $m$ , are in a one-dimensional box of length  $a$ . What are the four lowest energies of the system? What are the degeneracies of these energies if the two particles are (a) distinguishable, (b) indistinguishable bosons, or (c) indistinguishable fermions?
4. Two particles in a one-dimensional box of length  $a$  occupy the  $n = 4$  and  $n' = 3$  states. Write the properly normalized wavefunctions if the particles are (a) distinguishable, (b) indistinguishable bosons, or (c) indistinguishable fermions.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [5.E: Multi-Particle Systems \(Exercises\)](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## CHAPTER OVERVIEW

### 6: Three-Dimensional Quantum Mechanics

In this chapter, we shall extend our previous one-dimensional formulation of non-relativistic quantum mechanics to produce a fully three-dimensional theory.

- [6.1: Fundamental Concepts](#)
- [6.2: Particle in Box](#)
- [6.3: Degenerate Electron Gases](#)
- [6.4: White Dwarf Stars](#)
- [6.5: Exercises](#)

#### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [6: Three-Dimensional Quantum Mechanics](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 6.1: Fundamental Concepts

We have seen that in one dimension the instantaneous state of a single non-relativistic particle is fully specified by a complex wavefunction,  $\psi(x, t)$ . The probability of finding the particle at time  $t$  between  $x$  and  $x + dx$  is  $P(x, t) dx$ , where

$$P(x, t) = |\psi(x, t)|^2. \quad (6.1.1)$$

Moreover, the wavefunction is normalized such that

$$\int_{-\infty}^{\infty} |\psi(x, t)|^2 dx = 1 \quad (6.1.2)$$

at all times.

In three dimensions, the instantaneous state of a single particle is also fully specified by a complex wavefunction,  $\psi(x, y, z, t)$ . By analogy with the one-dimensional case, the probability of finding the particle at time  $t$  between  $x$  and  $x + dx$ , between  $y$  and  $y + dy$ , and between  $z$  and  $z + dz$ , is  $P(x, y, z, t) dx dy dz$ , where

$$P(x, y, z, t) = |\psi(x, y, z, t)|^2. \quad (6.1.3)$$

As usual, this interpretation of the wavefunction only makes sense if the wavefunction is normalized such that

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\psi(x, y, z, t)|^2 dx dy dz = 1. \quad (6.1.4)$$

This normalization constraint ensures that the probability of finding the particle anywhere in space is always unity.

In one dimension, we can write the probability conservation equation (see Section [\[s4.5\]](#))

$$\frac{\partial |\psi|^2}{\partial t} + \frac{\partial j}{\partial x} = 0, \quad (6.1.5)$$

where

$$j = \frac{i\hbar}{2m} \left( \psi \frac{\partial \psi^*}{\partial x} - \psi^* \frac{\partial \psi}{\partial x} \right) \quad (6.1.6)$$

is the flux of probability along the  $x$ -axis. Integrating Equation ([\[e6.5\]](#)) over all space, and making use of the fact that  $\psi \rightarrow 0$  as  $|x| \rightarrow \infty$  if  $\psi$  is to be square-integrable, we obtain

$$\frac{d}{dt} \int_{-\infty}^{\infty} |\psi(x, t)|^2 dx = 0. \quad (6.1.7)$$

In other words, if the wavefunction is initially normalized then it stays normalized as time progresses. This is a necessary criterion for the viability of our basic interpretation of  $|\psi|^2$  as a probability density.

In three dimensions, by analogy with the one dimensional case, the probability conservation equation becomes

$$\frac{\partial |\psi|^2}{\partial t} + \frac{\partial j_x}{\partial x} + \frac{\partial j_y}{\partial y} + \frac{\partial j_z}{\partial z} = 0. \quad (6.1.8)$$

Here,

$$j_x = \frac{i\hbar}{2m} \left( \psi \frac{\partial \psi^*}{\partial x} - \psi^* \frac{\partial \psi}{\partial x} \right) \quad (6.1.9)$$

is the flux of probability along the  $x$ -axis, and

$$j_y = \frac{i\hbar}{2m} \left( \psi \frac{\partial \psi^*}{\partial y} - \psi^* \frac{\partial \psi}{\partial y} \right) \quad (6.1.10)$$

the flux of probability along the  $y$ -axis, et cetera. Integrating Equation ([\[e6.8\]](#)) over all space, and making use of the fact that  $\psi \rightarrow 0$  as  $|\mathbf{r}| \rightarrow \infty$  if  $\psi$  is to be square-integrable, we obtain

$$\frac{d}{dt} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\psi(x, y, z, t)|^2 dx dy dz = 0. \quad (6.1.11)$$

Thus, the normalization of the wavefunction is again preserved as time progresses, as must be the case if  $|\psi|^2$  is to be interpreted as a probability density.

In one dimension, position is represented by the algebraic operator  $x$ , whereas momentum is represented by the differential operator  $-i\hbar\partial/\partial x$ . (See Section [s4.6].) By analogy, in three dimensions, the Cartesian coordinates  $x$ ,  $y$ , and  $z$  are represented by the algebraic operators  $x$ ,  $y$ , and  $z$ , respectively, whereas the three Cartesian components of momentum,  $p_x$ ,  $p_y$ , and  $p_z$ , have the following representations:

$$\begin{aligned} p_x &\equiv -i\hbar \frac{\partial}{\partial x}, \\ p_y &\equiv -i\hbar \frac{\partial}{\partial y}, \\ p_z &\equiv -i\hbar \frac{\partial}{\partial z}. \end{aligned}$$

Let  $x_1 = x$ ,  $x_2 = y$ ,  $x_3 = z$ , and  $p_1 = p_x$ , et cetera. Because the  $x_i$  are independent variables (i.e.,  $\partial x_i / \partial x_j = \delta_{ij}$ ), we conclude that the various position and momentum operators satisfy the following commutation relations:

$$\begin{aligned} [x_i, x_j] &= 0, \\ [p_i, p_j] &= 0, \\ [x_i, p_j] &= i\hbar \delta_{ij}. \end{aligned}$$

Now, we know, from Section [smeas], that two dynamical variables can only be (exactly) measured simultaneously if the operators that represent them in quantum mechanics commute with one another. Thus, it is clear, from the previous commutation relations, that the only restriction on measurement in a system consisting of a single particle moving in three dimensions is that it is impossible to simultaneously measure a given position coordinate and the corresponding component of momentum. Note, however, that it is perfectly possible to simultaneously measure two different positions coordinates, or two different components of the momentum. The commutation relations ([commxx])–([commxp]) again illustrate the point that quantum mechanical operators corresponding to different degrees of freedom of a dynamical system (in this case, motion in different directions) tend to commute with one another. (See Section [sfuncon].)

In one dimension, the time evolution of the wavefunction is given by [see Equation ([etimed])]

$$i\hbar \frac{\partial \psi}{\partial t} = H \psi, \quad (6.1.12)$$

where  $H$  is the Hamiltonian. The same equation governs the time evolution of the wavefunction in three dimensions.

Now, in one dimension, the Hamiltonian of a non-relativistic particle of mass  $m$  takes the form

$$H = \frac{p_x^2}{2m} + V(x, t), \quad (6.1.13)$$

where  $V(x)$  is the potential energy. In three dimensions, this expression generalizes to

$$H = \frac{p_x^2 + p_y^2 + p_z^2}{2m} + V(x, y, z, t). \quad (6.1.14)$$

Hence, making use of Equations ([e6.12])–([e6.14]) and ([e6.15]), the three-dimensional version of the time-dependent Schrödinger equation becomes [see Equation ([e3.1])]

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \psi + V \psi. \quad (6.1.15)$$

Here, the differential operator

$$\nabla^2 \equiv \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \quad (6.1.16)$$



is known as the *Laplacian*. Incidentally, the probability conservation equation ([\[e6.8\]](#)) is easily derivable from Equation ([\[esh3d\]](#)). An eigenstate of the Hamiltonian corresponding to the eigenvalue  $E$  satisfies

$$H \psi = E \psi. \quad (6.1.17)$$

It follows from Equation ([\[e6.15\]](#)) that (see Section [\[sstat\]](#))

$$\psi(x, y, z, t) = \psi(x, y, z) e^{-i E t / \hbar}, \quad (6.1.18)$$

where the stationary wavefunction  $\psi(x, y, z)$  satisfies the three-dimensional version of the time-independent Schrödinger equation [see Equation ([\[etimeii\]](#))]:

$$\nabla^2 \psi = \frac{2m}{\hbar^2} (V - E) \psi, \quad (6.1.19)$$

where  $V$  is assumed not to depend explicitly on  $t$ .

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [6.1: Fundamental Concepts](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 6.2: Particle in Box

Consider a particle of mass  $m$  trapped inside a cubic box of dimension  $a$ . (See Section [\[s5.2\]](#).) The particle's stationary wavefunction,  $\psi(x, y, z)$ , satisfies

$$\left( \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) \psi = -\frac{2m}{\hbar^2} E \psi, \quad (6.2.1)$$

where  $E$  is the particle energy. The wavefunction satisfies the boundary condition that it must be zero at the edges of the box.

Let us search for a separable solution to the previous equation of the form

$$\psi(x, y, z) = X(x) Y(y) Z(z). \quad (6.2.2)$$

The factors of the wavefunction satisfy the boundary conditions  $X(0) = X(a) = 0$ ,  $Y(0) = Y(a) = 0$ , and  $Z(0) = Z(a) = 0$ . Substituting Equation ([\[e6.22\]](#)) into Equation ([\[e6.21\]](#)), and rearranging, we obtain

$$\frac{X''}{X} + \frac{Y''}{Y} + \frac{Z''}{Z} = -\frac{2m}{\hbar^2} E, \quad (6.2.3)$$

where  $'$  denotes a derivative with respect to argument. It is evident that the only way in which the previous equation can be satisfied at all points within the box is if

$$\begin{aligned} \frac{X''}{X} &= -k_x^2, \\ \frac{Y''}{Y} &= -k_y^2, \\ \frac{Z''}{Z} &= -k_z^2, \end{aligned}$$

where  $k_x^2$ ,  $k_y^2$ , and  $k_z^2$  are spatial constants. Note that the right-hand sides of the previous equations must contain negative, rather than positive, spatial constants, because it would not otherwise be possible to satisfy the boundary conditions. The solutions to the previous equations which are properly normalized, and satisfy the boundary conditions, are [see Equation ([\[e5.11\]](#))]

$$\begin{aligned} X(x) &= \sqrt{\frac{2}{a}} \sin(k_x x), \\ Y(y) &= \sqrt{\frac{2}{a}} \sin(k_y y), \\ Z(z) &= \sqrt{\frac{2}{a}} \sin(k_z z), \end{aligned}$$

where

$$\begin{aligned} k_x &= \frac{l_x \pi}{a}, \\ k_y &= \frac{l_y \pi}{a}, \\ k_z &= \frac{l_z \pi}{a}. \end{aligned}$$

Here,  $l_x$ ,  $l_y$ , and  $l_z$  are positive integers. Thus, from Equations ([\[e7.28\]](#))–([\[e7.31\]](#)), the energy of the system is written [see Equation ([\[eenergy\]](#))]

$$E = \frac{l^2 \pi^2 \hbar^2}{2 m a^2}. \quad (6.2.4)$$

where

$$l^2 = l_x^2 + l_y^2 + l_z^2. \quad (6.2.5)$$

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [6.2: Particle in Box](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 6.3: Degenerate Electron Gases

Consider  $N$  electrons trapped in a cubic box of dimension  $a$ . Let us treat the electrons as essentially non-interacting particles. According to Section [snon], the total energy of a system consisting of many non-interacting particles is simply the sum of the single-particle energies of the individual particles. Furthermore, electrons are subject to the *Pauli exclusion principle* (see Section [siden]), because they are indistinguishable fermions. The exclusion principle states that no two electrons in our system can occupy the same single-particle energy level. Now, from the previous section, the single-particle energy levels for a particle in a box are characterized by the three quantum numbers,  $l_x$ ,  $l_y$ , and  $l_z$ . Thus, we conclude that no two electrons in our system can have the same set of values of  $l_x$ ,  $l_y$ , and  $l_z$ . It turns out that this is not quite true, because electrons possess an intrinsic angular momentum called spin. The spin states of an electron are governed by an additional quantum number, which can take one of two different values. (See Chapter [sspin].) Hence, when spin is taken into account, we conclude that a maximum of two electrons (with different spin quantum numbers) can occupy a single-particle energy level corresponding to a particular set of values of  $l_x$ ,  $l_y$ , and  $l_z$ . Note, from Equations ([e7.38]) and ([e7.39]), that the associated particle energy is proportional to  $l^2 = l_x^2 + l_y^2 + l_z^2$ .

Suppose that our electrons are cold: that is, they have comparatively little thermal energy. In this case, we would expect them to fill the lowest single-particle energy levels available to them. We can imagine the single-particle energy levels as existing in a sort of three-dimensional quantum number space whose Cartesian coordinates are  $l_x$ ,  $l_y$ , and  $l_z$ . Thus, the energy levels are uniformly distributed in this space on a cubic lattice. Moreover, the distance between nearest neighbor energy levels is unity. This implies that the number of energy levels per unit volume is also unity. Finally, the energy of a given energy level is proportional to its distance,  $l^2 = l_x^2 + l_y^2 + l_z^2$ , from the origin.

Because we expect cold electrons to occupy the lowest energy levels available to them, but only two electrons can occupy a given energy level, it follows that if the number of electrons,  $N$ , is very large then the filled energy levels will be approximately distributed in a sphere centered on the origin of quantum number space. The number of energy levels contained in a sphere of radius  $l$  is approximately equal to the volume of the sphere—because the number of energy levels per unit volume is unity. It turns out that this is not quite correct, because we have forgotten that the quantum numbers  $l_x$ ,  $l_y$ , and  $l_z$  can only take positive values. Hence, the filled energy levels actually only occupy one octant of a sphere. The radius  $l_F$  of the octant of filled energy levels in quantum number space can be calculated by equating the number of energy levels it contains to the number of electrons,  $N$ . Thus, we can write

$$N = 2 \times \frac{1}{8} \times \frac{4\pi}{3} l_F^3. \quad (6.3.1)$$

Here, the factor 2 is to take into account the two spin states of an electron, and the factor  $1/8$  is to take account of the fact that  $l_x$ ,  $l_y$ , and  $l_z$  can only take positive values. Thus,

$$l_F = \left( \frac{3N}{\pi} \right)^{1/3}. \quad (6.3.2)$$

According to Equation ([e7.38]), the energy of the most energetic electrons—which is known as the *Fermi energy*—is given by

$$E_F = \frac{l_F^2 \pi^2 \hbar^2}{2 m_e a^2} = \frac{\pi^2 \hbar^2}{2 m a^2} \left( \frac{3N}{\pi} \right)^{2/3}, \quad (6.3.3)$$

where  $m_e$  is the electron mass. This can also be written as

$$E_F = \frac{\pi^2 \hbar^2}{2 m_e} \left( \frac{3n}{\pi} \right)^{2/3}, \quad (6.3.4)$$

where  $n = N/a^3$  is the number of electrons per unit volume (in real space). Note that the Fermi energy only depends on the number density of the confined electrons.

The mean energy of the electrons is given by

$$\bar{E} = E_F \int_0^{l_F} l^2 4\pi l^2 dl \bigg/ \frac{4}{3} \pi l_F^3 = \frac{3}{5} E_F, \quad (6.3.5)$$

because  $E \propto l^2$ , and the energy levels are uniformly distributed in quantum number space within an octant of radius  $l_F$ . Now, according to classical physics, the mean thermal energy of the electrons is  $(3/2) k_B T$ , where  $T$  is the electron temperature, and  $k_B$  the Boltzmann constant. Thus, if  $k_B T \ll E_F$  then our original assumption that the electrons are cold is valid. Note that, in this case, the electron energy is much larger than that predicted by classical physics—electrons in this state are termed *degenerate*. On the other hand, if  $k_B T \gg E_F$  then the electrons are hot, and are essentially governed by classical physics—electrons in this state are termed *non-degenerate*.

The total energy of a degenerate electron gas is

$$E_{\text{total}} = N \bar{E} = \frac{3}{5} N E_F. \quad (6.3.6)$$

Hence, the gas pressure takes the form

$$P = -\frac{\partial E_{\text{total}}}{\partial V} = \frac{2}{5} n E_F, \quad (6.3.7)$$

because  $E_F \propto a^{-2} = V^{-2/3}$ . [See Equation ([e7.42]).] Now, the pressure predicted by classical physics is  $P = n k_B T$ . Thus, a degenerate electron gas has a much higher pressure than that which would be predicted by classical physics. This is an entirely quantum mechanical effect, and is due to the fact that identical fermions cannot get significantly closer together than a de Broglie wavelength without violating the Pauli exclusion principle. Note that, according to Equation ([e7.43]), the mean spacing between degenerate electrons is

$$d \sim n^{-1/3} \sim \frac{h}{\sqrt{m_e E}} \sim \frac{h}{p} \sim \lambda, \quad (6.3.8)$$

where  $\lambda$  is the de Broglie wavelength. Thus, an electron gas is non-degenerate when the mean spacing between the electrons is much greater than the de Broglie wavelength, and becomes degenerate as the mean spacing approaches the de Broglie wavelength.

It turns out that the conduction (i.e., free) electrons inside metals are highly degenerate (because the number of electrons per unit volume is very large, and  $E_F \propto n^{2/3}$ ). Indeed, most metals are hard to compress as a direct consequence of the high degeneracy pressure of their conduction electrons. To be more exact, resistance to compression is usually measured in terms of a quantity known as the *bulk modulus*, which is defined

$$B = -V \frac{\partial P}{\partial V} \quad (6.3.9)$$

Now, for a fixed number of electrons,  $P \propto V^{-5/3}$ . [See Equations ([e7.42]) and ([e7.46]).] Hence,

$$B = \frac{5}{3} P = \frac{\pi^3 \hbar^2}{9 m} \left( \frac{3 n}{\pi} \right)^{5/3}. \quad (6.3.10)$$

For example, the number density of free electrons in magnesium is  $n \sim 8.6 \times 10^{28} \text{ m}^{-3}$ . This leads to the following estimate for the bulk modulus:  $B \sim 6.4 \times 10^{10} \text{ N m}^{-2}$ . The actual bulk modulus is  $B = 4.5 \times 10^{10} \text{ N m}^{-2}$ .

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [6.3: Degenerate Electron Gases](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 6.4: White Dwarf Stars

A main-sequence hydrogen-burning star, such as the Sun, is maintained in equilibrium via the balance of the gravitational attraction tending to make it collapse, and the thermal pressure tending to make it expand. Of course, the thermal energy of the star is generated by nuclear reactions occurring deep inside its core. Eventually, however, the star will run out of burnable fuel, and, therefore, start to collapse, as it radiates away its remaining thermal energy. What is the ultimate fate of such a star?

A burnt-out star is basically a gas of electrons and ions. As the star collapses, its density increases, and so the mean separation between its constituent particles decreases. Eventually, the mean separation becomes of order the de Broglie wavelength of the electrons, and the electron gas becomes degenerate. Note that the de Broglie wavelength of the ions is much smaller than that of the electrons, so the ion gas remains non-degenerate. Now, even at zero temperature, a degenerate electron gas exerts a substantial pressure, because the Pauli exclusion principle prevents the mean electron separation from becoming significantly smaller than the typical de Broglie wavelength. (See the previous section.) Thus, it is possible for a burnt-out star to maintain itself against complete collapse under gravity via the degeneracy pressure of its constituent electrons. Such stars are termed *white-dwarfs*. Let us investigate the physics of white-dwarfs in more detail.

The total energy of a white-dwarf star can be written

$$\mathcal{E} = K + U, \quad (6.4.1)$$

where  $K$  is the kinetic energy of the degenerate electrons (the kinetic energy of the ions is negligible), and  $U$  is the gravitational potential energy. Let us assume, for the sake of simplicity, that the density of the star is uniform. In this case, the gravitational potential energy takes the form

$$U = -\frac{3}{5} \frac{G M^2}{R}, \quad (6.4.2)$$

where  $G$  is the gravitational constant,  $M$  is the stellar mass, and  $R$  is the stellar radius.

From the previous section, the kinetic energy of a degenerate electron gas is simply

$$K = N \bar{E} = \frac{3}{5} N E_F = \frac{3}{5} N \frac{\pi^2 \hbar^2}{2 m_e} \left( \frac{3 N}{\pi V} \right)^{2/3}, \quad (6.4.3)$$

where  $N$  is the number of electrons,  $V$  the volume of the star, and  $m_e$  the electron mass.

The interior of a white-dwarf star is composed of atoms like  $C^{12}$  and  $O^{16}$  which contain equal numbers of protons, neutrons, and electrons. Thus,

$$M = 2 N m_p, \quad (6.4.4)$$

where  $m_p$  is the proton mass.

Equations (6.4.1)–(6.4.4) can be combined to give

$$\mathcal{E} = \frac{A}{R^2} - \frac{B}{R}, \quad (6.4.5)$$

where

$$A = \frac{3}{20} \left( \frac{9\pi}{8} \right)^{2/3} \frac{\hbar^2}{m_e} \left( \frac{M}{m_p} \right)^{5/3},$$

$$B = \frac{3}{5} G M^2.$$

The equilibrium radius of the star,  $R_*$ , is that which minimizes the total energy  $\mathcal{E}$ . In fact, it is easily demonstrated that

$$R_* = \frac{2 A}{B}, \quad (6.4.6)$$

which yields

$$R_* = \frac{(9\pi)^{2/3}}{8} \frac{\hbar^2}{G m_e m_p^{5/3} M^{1/3}}. \quad (6.4.7)$$

The previous formula can also be written

(6.4.8)

where  $R_\odot = 7 \times 10^5$  km is the solar radius, and  $M_\odot = 2 \times 10^{30}$  kg the solar mass. It follows that the radius of a typical solar mass white-dwarf is about 7000 km: that is, about the same as the radius of the Earth. The first white-dwarf to be discovered (in 1862) was the companion of Sirius. Nowadays, thousands of white-dwarfs have been observed, all with properties similar to those described previously.

Note from Equations ([e8wd4]), ([e8wd6]), and ([e8wd9]) that  $\bar{E} \propto M^{4/3}$ . In other words, the mean energy of the electrons inside a white dwarf increases as the stellar mass increases. Hence, for a sufficiently massive white dwarf, the electrons can become relativistic. It turns out that the degeneracy pressure for relativistic electrons only scales as  $R^{-1}$ , rather than  $R^{-2}$ , and thus is unable to balance the gravitational pressure [which also scales as  $R^{-1}$ —see Equation ([e7.52])]. It follows that electron degeneracy pressure is only able to halt the collapse of a burnt-out star provided that the stellar mass does not exceed some critical value, known as the *Chandrasekhar limit*, which turns out to be about 1.4 times the mass of the Sun. Stars whose mass exceeds the Chandrasekhar limit inevitably collapse to produce extremely compact objects, such as neutron stars (which are held up by the degeneracy pressure of their constituent neutrons), or black holes.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [6.4: White Dwarf Stars](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 6.5: Exercises

---

1. Consider a particle of mass  $m$  moving in a three-dimensional isotropic harmonic oscillator potential of force constant  $k$ . Solve the problem via the separation of variables, and obtain an expression for the allowed values of the total energy of the system (in a stationary state).
2. Repeat the calculation of the Fermi energy of a gas of fermions by assuming that the fermions are massless, so that the energy-momentum relation is  $E = pc$ .
3. Calculate the density of states of an electron gas in a cubic box of volume  $L^3$ , bearing in mind that there are two electrons per energy state. In other words, calculate the number of electron states in the interval  $E$  to  $E + dE$ . This number can be written  $dN = \rho(E) dE$ , where  $\rho$  is the density of states.
4. Repeat the previous calculation for a two-dimensional electron gas in a square box of area  $L^2$ .
5. Given that the number density of free electrons in copper is  $8.5 \times 10^{28} \text{ m}^{-3}$ , calculate the Fermi energy in electron volts, and the velocity of an electron whose kinetic energy is equal to the Fermi energy.
6. Obtain an expression for the Fermi energy (in eV) of an electron in a white dwarf star as a function of the stellar mass (in solar masses). At what mass does the Fermi energy equal the rest mass energy?

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)
- 

This page titled [6.5: Exercises](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).



## CHAPTER OVERVIEW

### 7: Orbital Angular Momentum

- [7.1: Angular Momentum Operators](#)
- [7.2: Representation of Angular Momentum](#)
- [7.3: Eigenstates of Angular Momentum](#)
- [7.4: Eigenvalues of  \$L\_z\$](#)
- [7.5: Eigenvalues of  \$L^2\$](#)
- [7.6: Spherical Harmonics](#)
- [7.E: Orbital Angular Momentum \(Exercises\)](#)

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [7: Orbital Angular Momentum](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 7.1: Angular Momentum Operators

In classical mechanics, the vector angular momentum,  $\mathbf{L}$ , of a particle of position vector  $\mathbf{r}$  and linear momentum  $\mathbf{p}$  is defined as

$$\mathbf{L} = \mathbf{r} \times \mathbf{p}. \quad (7.1.1)$$

It follows that

$$\begin{aligned} L_x &= y p_z - z p_y, \\ L_y &= z p_x - x p_z, \\ L_z &= x p_y - y p_x. \end{aligned}$$

Let us, first of all, consider whether it is possible to use the previous expressions as the definitions of the operators corresponding to the components of angular momentum in quantum mechanics, assuming that the  $x_i$  and  $p_i$  (where  $x_1 \equiv x$ ,  $p_1 \equiv p_x$ ,  $x_2 \equiv y$ , etc.) correspond to the appropriate quantum mechanical position and momentum operators. The first point to note is that expressions ([e8.1])–([e8.3]) are unambiguous with respect to the order of the terms in multiplicative factors, because the various position and momentum operators appearing in them all commute with one another. [See Equations ([commxp]).] Moreover, given that the  $x_i$  and the  $p_i$  are Hermitian operators, it is easily seen that the  $L_i$  are also Hermitian. This is important, because only Hermitian operators can represent physical variables in quantum mechanics. (See Section [s4.6].) We, thus, conclude that Equations ([e8.1])–([e8.3]) are plausible definitions for the quantum mechanical operators that represent the components of angular momentum.

Let us now derive the commutation relations for the  $L_i$ . For instance,

$$\begin{aligned} &= [y p_z - z p_y, z p_x - x p_z] = y p_x [p_z, z] + x p_y [z, p_z] \\ &= i \hbar (x p_y - y p_x) = i \hbar L_z, \end{aligned}$$

where use has been made of the definitions of the  $L_i$  [see Equations ([e8.1])–([e8.3])], and commutation relations ([commxx])–([commxp]) for the  $x_i$  and  $p_i$ . There are two similar commutation relations: one for  $L_y$  and  $L_z$ , and one for  $L_z$  and  $L_x$ . Collecting all of these commutation relations together, we obtain

$$\begin{aligned} [L_x, L_y] &= i \hbar L_z, \\ [L_y, L_z] &= i \hbar L_x, \\ [L_z, L_x] &= i \hbar L_y. \end{aligned}$$

By analogy with classical mechanics, the operator  $L^2$ , that represents the magnitude squared of the angular momentum vector, is defined

$$L^2 = L_x^2 + L_y^2 + L_z^2. \quad (7.1.2)$$

Now, it is easily demonstrated that if  $A$  and  $B$  are two general operators then

$$[A^2, B] = A [A, B] + [A, B] A. \quad (7.1.3)$$

Hence,

$$\begin{aligned} &= [L_y^2, L_x] + [L_z^2, L_x] \\ &= L_y [L_y, L_x] + [L_y, L_x] L_y + L_z [L_z, L_x] + [L_z, L_x] L_z \\ &= i \hbar (-L_y L_z - L_z L_y + L_z L_y + L_y L_z) = 0, \end{aligned}$$

where use has been made of Equations ([e8.6])–([e8.8]). In other words,  $L^2$  commutes with  $L_x$ . Likewise, it is easily demonstrated that  $L^2$  also commutes with  $L_y$ , and with  $L_z$ . Thus,

$$[L^2, L_x] = [L^2, L_y] = [L^2, L_z] = 0. \quad (7.1.4)$$

Recall, from Section [smeas], that in order for two physical quantities to be (exactly) measured simultaneously, the operators that represent them in quantum mechanics must commute with one another. Hence, the commutation relations ([e8.6])–([e8.8]) and ([e8.12]) imply that we can only simultaneously measure the magnitude squared of the angular momentum vector,  $L^2$ , together with, at most, one of its Cartesian components. By convention, we shall always choose to measure the  $z$ -component,  $L_z$ .

Finally, it is helpful to define the operators

$$L_{\pm} = L_x \pm i L_y. \quad (7.1.5)$$

Note that  $L_+$  and  $L_-$  are not Hermitian operators, but are the Hermitian conjugates of one another (see Section [s4.6]): that is,

$$(L_{\pm})^{\dagger} = L_{\mp}, \quad (7.1.6)$$

Moreover, it is easily seen that

$$\begin{aligned} L_+ L_- &= (L_x + i L_y)(L_x - i L_y) = L_x^2 + L_y^2 - i [L_x, L_y] = L_x^2 + L_y^2 + \hbar L_z \\ &= L^2 - L_z^2 + \hbar L_z. \end{aligned}$$

Likewise,

$$L_- L_+ = L^2 - L_z^2 - \hbar L_z, \quad (7.1.7)$$

giving

$$[L_+, L_-] = 2 \hbar L_z. \quad (7.1.8)$$

We also have

$$[L_+, L_z] = [L_x, L_z] + i [L_y, L_z] = -i \hbar L_y - \hbar L_x = -\hbar L_+, \quad (7.1.9)$$

and, similarly,

$$[L_-, L_z] = \hbar L_-. \quad (7.1.10)$$

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [7.1: Angular Momentum Operators](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 7.2: Representation of Angular Momentum

Now, we saw earlier, in Section [s7.2], that the operators,  $p_i$ , which represent the Cartesian components of linear momentum in quantum mechanics, can be represented as the spatial differential operators  $-i\hbar\partial/\partial x_i$ . Let us now investigate whether angular momentum operators can similarly be represented as spatial differential operators.

It is most convenient to perform our investigation using conventional spherical polar coordinates: that is,  $r$ ,  $\theta$ , and  $\phi$ . These are defined with respect to our usual Cartesian coordinates as follows:

$$\begin{aligned}x &= r \sin \theta \cos \phi, \\y &= r \sin \theta \sin \phi, \\z &= r \cos \theta.\end{aligned}$$

We deduce, after some tedious analysis, that

$$\begin{aligned}\frac{\partial}{\partial x} &= \sin \theta \cos \phi \frac{\partial}{\partial r} + \frac{\cos \theta \cos \phi}{r} \frac{\partial}{\partial \theta} - \frac{\sin \phi}{r \sin \theta} \frac{\partial}{\partial \phi}, \\ \frac{\partial}{\partial y} &= \sin \theta \sin \phi \frac{\partial}{\partial r} + \frac{\cos \theta \sin \phi}{r} \frac{\partial}{\partial \theta} + \frac{\cos \phi}{r \sin \theta} \frac{\partial}{\partial \phi}, \\ \frac{\partial}{\partial z} &= \cos \theta \frac{\partial}{\partial r} - \frac{\sin \theta}{r} \frac{\partial}{\partial \theta}.\end{aligned}$$

Making use of the definitions ([e8.1])–([e8.3]), ([e8.9]), and ([e8.13]), the fundamental representation ([e6.12])–([e6.14]) of the  $p_i$  operators as spatial differential operators, Equations ([e8.21])–([e8zz]), and a great deal of tedious analysis, we finally obtain

$$\begin{aligned}L_x &= -i\hbar \left( -\sin \phi \frac{\partial}{\partial \theta} - \cos \phi \cot \theta \frac{\partial}{\partial \phi} \right), \\ L_y &= -i\hbar \left( \cos \phi \frac{\partial}{\partial \theta} - \sin \phi \cot \theta \frac{\partial}{\partial \phi} \right), \\ L_z &= -i\hbar \frac{\partial}{\partial \phi},\end{aligned}$$

as well as

$$L^2 = -\hbar^2 \left[ \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2} \right], \quad (7.2.1)$$

and

$$L_{\pm} = \hbar e^{\pm i\phi} \left( \pm \frac{\partial}{\partial \theta} + i \cot \theta \frac{\partial}{\partial \phi} \right). \quad (7.2.2)$$

We, thus, conclude that all of our angular momentum operators can be represented as differential operators involving the angular spherical coordinates,  $\theta$  and  $\phi$ , but not involving the radial coordinate,  $r$ .

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled 7.2: Representation of Angular Momentum is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 7.3: Eigenstates of Angular Momentum

Let us find the simultaneous eigenstates of the angular momentum operators  $L_z$  and  $L^2$ . Because both of these operators can be represented as purely angular differential operators, it stands to reason that their eigenstates only depend on the angular coordinates  $\theta$  and  $\phi$ . Thus, we can write

$$\begin{aligned} L_z Y_{l,m}(\theta, \phi) &= m \hbar Y_{l,m}(\theta, \phi), \\ L^2 Y_{l,m}(\theta, \phi) &= l(l+1) \hbar^2 Y_{l,m}(\theta, \phi). \end{aligned}$$

Here, the  $Y_{l,m}(\theta, \phi)$  are the eigenstates in question, whereas the dimensionless quantities  $m$  and  $l$  parameterize the eigenvalues of  $L_z$  and  $L^2$ , which are  $m \hbar$  and  $l(l+1) \hbar^2$ , respectively. Of course, we expect the  $Y_{l,m}$  to be both mutually orthogonal and properly normalized (see Section [seig]), so that

$$\oint Y_{l',m'}^*(\theta, \phi) Y_{l,m}(\theta, \phi) d\Omega = \delta_{ll'} \delta_{mm'}, \quad (7.3.1)$$

where  $d\Omega = \sin \theta d\theta d\phi$  is an element of solid angle, and the integral is over all solid angle.

Now,

$$\begin{aligned} L_z (L_+ Y_{l,m}) &= (L_z L_+ + [L_z, L_+]) Y_{l,m} = (L_z L_+ + \hbar L_+) Y_{l,m} \\ &= (m+1) \hbar (L_+ Y_{l,m}), \end{aligned}$$

where use has been made of Equation ([e8.19]). We, thus, conclude that when the operator  $L_+$  operates on an eigenstate of  $L_z$  corresponding to the eigenvalue  $m \hbar$  it converts it to an eigenstate corresponding to the eigenvalue  $(m+1) \hbar$ . Hence,  $L_+$  is known as the *raising operator* (for  $L_z$ ). It is also easily demonstrated that

$$L_z (L_- Y_{l,m}) = (m-1) \hbar (L_- Y_{l,m}). \quad (7.3.2)$$

In other words, when  $L_-$  operates on an eigenstate of  $L_z$  corresponding to the eigenvalue  $m \hbar$  it converts it to an eigenstate corresponding to the eigenvalue  $(m-1) \hbar$ . Hence,  $L_-$  is known as the *lowering operator* (for  $L_z$ ).

Writing

$$\begin{aligned} L_+ Y_{l,m} &= c_{l,m}^+ Y_{l,m+1}, \\ L_- Y_{l,m} &= c_{l,m}^- Y_{l,m-1}, \end{aligned}$$

we obtain

$$L_- L_+ Y_{l,m} = c_{l,m}^+ c_{l,m+1}^- Y_{l,m} = [l(l+1) - m(m+1)] \hbar^2 Y_{l,m}, \quad (7.3.3)$$

where use has been made of Equation ([e8.17]). Likewise,

$$L_+ L_- Y_{l,m} = c_{l,m-1}^+ c_{l,m}^- Y_{l,m} = [l(l+1) - m(m-1)] \hbar^2 Y_{l,m}, \quad (7.3.4)$$

where use has been made of Equation ([e8.15]). It follows that

$$\begin{aligned} c_{l,m}^+ c_{l,m+1}^- &= [l(l+1) - m(m+1)] \hbar^2, \\ c_{l,m-1}^+ c_{l,m}^- &= [l(l+1) - m(m-1)] \hbar^2. \end{aligned}$$

These equations are satisfied when

$$c_{l,m}^\pm = [l(l+1) - m(m \pm 1)]^{1/2} \hbar. \quad (7.3.5)$$

Hence, we can write

$$\begin{aligned} L_+ Y_{l,m} &= [l(l+1) - m(m+1)]^{1/2} \hbar Y_{l,m+1}, \\ L_- Y_{l,m} &= [l(l+1) - m(m-1)]^{1/2} \hbar Y_{l,m-1}. \end{aligned}$$

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [7.3: Eigenstates of Angular Momentum](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 7.4: Eigenvalues of $L_z$

It seems reasonable to attempt to write the eigenstate  $Y_{l,m}(\theta, \phi)$  in the separable form

$$Y_{l,m}(\theta, \phi) = \Theta_{l,m}(\theta) \Phi_m(\phi). \quad (7.4.1)$$

We can satisfy the orthonormality constraint ([e8.31]) provided that

$$\int_0^\pi \Theta_{l',m'}^*(\theta) \Theta_{l,m}(\theta) \sin \theta d\theta = \delta_{ll'},$$

$$\int_0^{2\pi} \Phi_{m'}^*(\phi) \Phi_m(\phi) d\phi = \delta_{mm'}.$$

Note, from Equation ([e8.26]), that the differential operator which represents  $L_z$  only depends on the azimuthal angle  $\phi$ , and is independent of the polar angle  $\theta$ . It therefore follows from Equations ([e8.26]), ([e8.29]), and ([e8.34]) that

$$-i\hbar \frac{d\Phi_m}{d\phi} = m\hbar \Phi_m. \quad (7.4.2)$$

The solution of this equation is

$$\Phi_m(\phi) \sim e^{im\phi}. \quad (7.4.3)$$

Here, the symbol  $\sim$  just means that we are neglecting multiplicative constants.

Our basic interpretation of a wavefunction as a quantity whose modulus squared represents the probability density of finding a particle at a particular point in space suggests that a physical wavefunction must be single-valued in space. Otherwise, the probability density at a given point would not, in general, have a unique value, which does not make physical sense. Hence, we demand that the wavefunction ([e8.38]) be single-valued: that is,  $\Phi_m(\phi + 2\pi) = \Phi_m(\phi)$  for all  $\phi$ . This immediately implies that the quantity  $m$  is quantized. In fact,  $m$  can only take integer values. Thus, we conclude that the eigenvalues of  $L_z$  are also quantized, and take the values  $m\hbar$ , where  $m$  is an integer. [A more rigorous argument is that  $\Phi_m(\phi)$  must be continuous in order to ensure that  $L_z$  is an Hermitian operator, because the proof of hermiticity involves an integration by parts in  $\phi$  that has canceling contributions from  $\phi = 0$  and  $\phi = 2\pi$ .]

Finally, we can easily normalize the eigenstate ([e8.38]) by making use of the orthonormality constraint ([e8.36]). We obtain

$$\Phi_m(\phi) = \frac{e^{im\phi}}{\sqrt{2\pi}}. \quad (7.4.4)$$

This is the properly normalized eigenstate of  $L_z$  corresponding to the eigenvalue  $m\hbar$ .

### Contributors and Attributions

- { {template.ContribFitzpatrick()}}

This page titled 7.4: Eigenvalues of  $L_z$  is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 7.5: Eigenvalues of $L^2$

Consider the angular wavefunction  $\psi(\theta, \phi) = Y_{l,m}(\theta, \phi)$ . We know that

$$\oint \psi^*(\theta, \phi) \psi(\theta, \phi) d\Omega \geq 0, \quad (7.5.1)$$

because  $\psi^* \psi \equiv |\psi|^2$  is a positive-definite real quantity. Hence, making use of Equations ([e5.48]) and ([e8.14]), we find that

$$\oint (L_+ Y_{l,m})^* (L_+ Y_{l,m}) d\Omega = \oint Y_{l,m}^* (L_+)^{\dagger} (L_+ Y_{l,m}) d\Omega = \oint Y_{l,m}^* L_- L_+ Y_{l,m} d\Omega \geq 0.$$

It follows from Equations ([e8.17]), and ([e8.29])–([e8.31]) that

$$\begin{aligned} \oint Y_{l,m}^* (L^2 - L_z^2 - \hbar L_z) Y_{l,m} d\Omega &= \oint Y_{l,m}^* \hbar^2 [l(l+1) - m(m+1)] Y_{l,m} d\Omega \\ &= \hbar^2 [l(l+1) - m(m+1)] \oint Y_{l,m}^* Y_{l,m} d\Omega \\ &= \hbar^2 [l(l+1) - m(m+1)] \geq 0. \end{aligned}$$

We, thus, obtain the constraint

$$l(l+1) \geq m(m+1). \quad (7.5.2)$$

Likewise, the inequality

$$\oint (L_- Y_{l,m})^* (L_- Y_{l,m}) d\Omega = \oint Y_{l,m}^* L_+ L_- Y_{l,m} d\Omega \geq 0 \quad (7.5.3)$$

leads to a second constraint:

$$l(l+1) \geq m(m-1). \quad (7.5.4)$$

Without loss of generality, we can assume that  $l \geq 0$ . This is reasonable, from a physical standpoint, because  $l(l+1)\hbar^2$  is supposed to represent the magnitude squared of something, and should, therefore, only take non-negative values. If  $l$  is non-negative then the constraints ([e8.42]) and ([e8.44]) are equivalent to the following constraint:

$$-l \leq m \leq l. \quad (7.5.5)$$

We, thus, conclude that the quantum number  $m$  can only take a restricted range of integer values.

Now, if  $m$  can only take a restricted range of integer values then there must exist a lowest possible value that it can take. Let us call this special value  $m_-$ , and let  $Y_{l,m_-}$  be the corresponding eigenstate. Suppose we act on this eigenstate with the lowering operator  $L_-$ . According to Equation ([e8.32]), this will have the effect of converting the eigenstate into that of a state with a lower value of  $m$ . However, no such state exists. A non-existent state is represented in quantum mechanics by the null wavefunction,  $\psi = 0$ . Thus, we must have

$$L_- Y_{l,m_-} = 0. \quad (7.5.6)$$

From Equation ([e8.15]),

$$L^2 = L_+ L_- + L_z^2 - \hbar L_z \quad (7.5.7)$$

Hence,

$$L^2 Y_{l,m_-} = (L_+ L_- + L_z^2 - \hbar L_z) Y_{l,m_-}, \quad (7.5.8)$$

or

$$l(l+1) Y_{l,m_-} = m_- (m_- - 1) Y_{l,m_-}, \quad (7.5.9)$$

where use has been made of ([e8.29]), ([e8.30]), and ([e8.46]). It follows that

$$l(l+1) = m_- (m_- - 1). \quad (7.5.10)$$



Assuming that  $m_-$  is negative, the solution to the previous equation is

$$m_- = -l. \quad (7.5.11)$$

We can similarly show that the largest possible value of  $m$  is

$$m_+ = +l. \quad (7.5.12)$$

The previous two results imply that  $l$  is an integer, because  $m_-$  and  $m_+$  are both constrained to be integers.

We can now formulate the rules that determine the allowed values of the quantum numbers  $l$  and  $m$ . The quantum number  $l$  takes the non-negative integer values  $0, 1, 2, 3, \dots$ . Once  $l$  is given, the quantum number  $m$  can take any integer value in the range

$$-l, -l+1, \dots, 0, \dots, l-1, l. \quad (7.5.13)$$

Thus, if  $l = 0$  then  $m$  can only take the value 0, if  $l = 1$  then  $m$  can take the values  $-1, 0, +1$ , if  $l = 2$  then  $m$  can take the values  $-2, -1, 0, +1, +2$  and so on.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [7.5: Eigenvalues of  \$L^2\$](#)  is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 7.6: Spherical Harmonics

The simultaneous eigenstates,  $Y_{l,m}(\theta, \phi)$ , of  $L^2$  and  $L_z$  are known as the *spherical harmonics*. Let us investigate their functional form.

We know that

$$L_+ Y_{l,l}(\theta, \phi) = 0, \quad (7.6.1)$$

because there is no state for which  $m$  has a larger value than  $+l$ . Writing

$$Y_{l,l}(\theta, \phi) = \Theta_{l,l}(\theta) e^{i l \phi} \quad (7.6.2)$$

[see Equations ([e8.34]) and ([e8.38])], and making use of Equation ([e8.28]), we obtain

$$\hbar e^{i \phi} \left( \frac{\partial}{\partial \theta} + i \cot \theta \frac{\partial}{\partial \phi} \right) \Theta_{l,l}(\theta) e^{i l \phi} = 0. \quad (7.6.3)$$

This equation yields

$$\frac{d\Theta_{l,l}}{d\theta} - l \cot \theta \Theta_{l,l} = 0. \quad (7.6.4)$$

which can easily be solved to give

$$\Theta_{l,l} \sim (\sin \theta)^l. \quad (7.6.5)$$

Hence, we conclude that

$$Y_{l,l}(\theta, \phi) \sim (\sin \theta)^l e^{i l \phi}. \quad (7.6.6)$$

Likewise, it is easy to demonstrate that

$$Y_{l,-l}(\theta, \phi) \sim (\sin \theta)^l e^{-i l \phi}. \quad (7.6.7)$$

Once we know  $Y_{l,l}$ , we can obtain  $Y_{l,l-1}$  by operating on  $Y_{l,l}$  with the lowering operator  $L_-$ . Thus,

$$Y_{l,l-1} \sim L_- Y_{l,l} \sim e^{-i \phi} \left( -\frac{\partial}{\partial \theta} + i \cot \theta \frac{\partial}{\partial \phi} \right) (\sin \theta)^l e^{i l \phi}, \quad (7.6.8)$$

where use has been made of Equation ([e8.28]). The previous equation yields

$$Y_{l,l-1} \sim e^{i(l-1)\phi} \left( \frac{d}{d\theta} + l \cot \theta \right) (\sin \theta)^l. \quad (7.6.9)$$

Now,

$$\left( \frac{d}{d\theta} + l \cot \theta \right) f(\theta) \equiv \frac{1}{(\sin \theta)^l} \frac{d}{d\theta} [(\sin \theta)^l f(\theta)], \quad (7.6.10)$$

where  $f(\theta)$  is a general function. Hence, we can write

$$Y_{l,l-1}(\theta, \phi) \sim \frac{e^{i(l-1)\phi}}{(\sin \theta)^{l-1}} \left( \frac{1}{\sin \theta} \frac{d}{d\theta} \right) (\sin \theta)^{2l}. \quad (7.6.11)$$

ikewise, we can show that

$$Y_{l,-l+1}(\theta, \phi) \sim L_+ Y_{l,-l} \sim \frac{e^{-i(l-1)\phi}}{(\sin \theta)^{l-1}} \left( \frac{1}{\sin \theta} \frac{d}{d\theta} \right) (\sin \theta)^{2l}. \quad (7.6.12)$$

We can now obtain  $Y_{l,l-2}$  by operating on  $Y_{l,l-1}$  with the lowering operator. We get

$$Y_{l,l-2} \sim L_- Y_{l,l-1} \sim e^{-i \phi} \left( -\frac{\partial}{\partial \theta} + i \cot \theta \frac{\partial}{\partial \phi} \right) \frac{e^{i(l-1)\phi}}{(\sin \theta)^{l-1}} \left( \frac{1}{\sin \theta} \frac{d}{d\theta} \right) (\sin \theta)^{2l}, \quad (7.6.13)$$

which reduces to

$$Y_{l,l-2} \sim e^{-i(l-2)\phi} \left[ \frac{d}{d\theta} + (l-1) \cot \theta \right] \frac{1}{(\sin \theta)^{l-1}} \left( \frac{1}{\sin \theta} \frac{d}{d\theta} \right) (\sin \theta)^{2l}. \quad (7.6.14)$$

Finally, making use of Equation ([e8.64]), we obtain

$$Y_{l,l-2}(\theta, \phi) \sim \frac{e^{i(l-2)\phi}}{(\sin \theta)^{l-2}} \left( \frac{1}{\sin \theta} \frac{d}{d\theta} \right)^2 (\sin \theta)^{2l}. \quad (7.6.15)$$

Likewise, we can show that

$$Y_{l,-l+2}(\theta, \phi) \sim L_+ Y_{l,-l+1} \sim \frac{e^{-i(l-2)\phi}}{(\sin \theta)^{l-2}} \left( \frac{1}{\sin \theta} \frac{d}{d\theta} \right)^2 (\sin \theta)^{2l}. \quad (7.6.16)$$

A comparison of Equations ([e8.59]), ([e8.64a]), and ([e8.68]) reveals the general functional form of the spherical harmonics:

$$Y_{l,m}(\theta, \phi) \sim \frac{e^{im\phi}}{(\sin \theta)^m} \left( \frac{1}{\sin \theta} \frac{d}{d\theta} \right)^{l-m} (\sin \theta)^{2l}. \quad (7.6.17)$$

Here,  $m$  is assumed to be non-negative. Making the substitution  $u = \cos \theta$ , we can also write

$$Y_{l,m}(u, \phi) \sim e^{im\phi} (1-u^2)^{-m/2} \left( \frac{d}{du} \right)^{l-m} (1-u^2)^l. \quad (7.6.18)$$

Finally, it is clear from Equations ([e8.60]), ([e8.65]), and ([e8.69]) that

$$Y_{l,-m} \sim Y_{l,m}^*. \quad (7.6.19)$$




Figure 18: The  $|Y_{l,m}(\theta, \phi)|^2$  plotted as a functions of  $\theta$ . The solid, short-dashed, and long-dashed curves correspond to  $l, m = 0, 0$ , and  $1, 0$ , and  $1, \pm 1$ , respectively.

We now need to normalize our spherical harmonic functions so as to ensure that

$$\oint |Y_{l,m}(\theta, \phi)|^2 d\Omega = 1. \quad (7.6.20)$$

After a great deal of tedious analysis, the normalized spherical harmonic functions are found to take the form

$$Y_{l,m}(\theta, \phi) = (-1)^m \left[ \frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!} \right]^{1/2} P_{l,m}(\cos \theta) e^{im\phi} \quad (7.6.21)$$

for  $m \geq 0$ , where the  $P_{l,m}$  are known as *associated Legendre polynomials*, and are written

$$P_{l,m}(u) = (-1)^{l+m} \frac{(l+m)!}{(l-m)!} \frac{(1-u^2)^{-m/2}}{2^l l!} \left( \frac{d}{du} \right)^{l-m} (1-u^2)^l \quad (7.6.22)$$

for  $m \geq 0$ . Alternatively,

$$P_{l,m}(u) = (-1)^l \frac{(1-u^2)^{m/2}}{2^l l!} \left( \frac{d}{du} \right)^{l+m} (1-u^2)^l, \quad (7.6.23)$$

for  $m \geq 0$ . The spherical harmonics characterized by  $m < 0$  can be calculated from those characterized by  $m > 0$  via the identity

$$Y_{l,-m} = (-1)^m Y_{l,m}^*. \quad (7.6.24)$$

The spherical harmonics are orthonormal: that is,

$$\oint Y_{l',m'}^* Y_{l,m} d\Omega = \delta_{l'l} \delta_{m'm}, \quad (7.6.25)$$

and also form a complete set. In other words, any well-behaved function of  $\theta$  and  $\phi$  can be represented as a superposition of spherical harmonics. Finally, and most importantly, the spherical harmonics are the simultaneous eigenstates of  $L_z$  and  $L^2$  corresponding to the eigenvalues  $m \hbar$  and  $l(l+1) \hbar^2$ , respectively.




Figure 19: The  $|Y_{l,m}(\theta, \phi)|^2$  plotted as a functions of  $\theta$ . The solid, short-dashed, and long-dashed curves correspond to  $l, m = 2, 0$ , and  $2, \pm 1$ , and  $2, \pm 2$  respectively.

All of the  $l = 0$ ,  $l = 1$ , and  $l = 2$  spherical harmonics are listed below:

$$\begin{aligned}Y_{0,0} &= \frac{1}{\sqrt{4\pi}}, \\Y_{1,0} &= \sqrt{\frac{3}{4\pi}} \cos \theta, \\Y_{1,\pm 1} &= \mp \sqrt{\frac{3}{8\pi}} \sin \theta e^{\pm i \phi}, \\Y_{2,0} &= \sqrt{\frac{5}{16\pi}} (3 \cos^2 \theta - 1), \\Y_{2,\pm 1} &= \mp \sqrt{\frac{15}{8\pi}} \sin \theta \cos \theta e^{\pm i \phi}, \\Y_{2,\pm 2} &= \sqrt{\frac{15}{32\pi}} \sin^2 \theta e^{\pm 2 i \phi}.\end{aligned}$$

The  $\theta$  variation of these functions is illustrated in Figures [\[ylm1\]](#) and [\[ylm2\]](#).

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [7.6: Spherical Harmonics](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 7.E: Orbital Angular Momentum (Exercises)

---

1. A system is in the state  $\psi = Y_{l,m}(\theta, \phi)$ . Calculate  $\langle L_x \rangle$  and  $\langle L_x^2 \rangle$ .
2. Find the eigenvalues and eigenfunctions (in terms of the angles  $\theta$  and  $\phi$ ) of  $L_x$ .
3. Consider a beam of particles with  $l = 1$ . A measurement of  $L_x$  yields the result  $\hbar$ . What values will be obtained by a subsequent measurement of  $L_z$ , and with what probabilities? Repeat the calculation for the cases in which the measurement of  $L_x$  yields the results 0 and  $-\hbar$ .
4. The Hamiltonian for an axially symmetric rotator is given by

$$H = \frac{L_x^2 + L_y^2}{2 I_1} + \frac{L_z^2}{2 I_2}. \quad (7.E.1)$$

What are the eigenvalues of  $H$ ?

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [7.E: Orbital Angular Momentum \(Exercises\)](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## CHAPTER OVERVIEW

### 8: Central Potentials

In this chapter, we shall investigate the interaction of a non-relativistic particle of mass  $m$  and energy  $E$  with various so-called *central potentials*,  $V(r)$ , where  $r = (x^2 + y^2 + z^2)^{1/2}$  is the radial distance from the origin. It is, of course, most convenient to work in spherical coordinates— $r$ ,  $\theta$ ,  $\phi$ —during such an investigation. (See Section [\[s8.3\]](#).) Thus, we shall be searching for stationary wavefunctions,  $\psi(r, \theta, \phi)$ , that satisfy the time-independent Schrödinger equation (see Section [\[sstat\]](#))

$$H\psi = E\psi, \quad (8.1)$$

where the Hamiltonian takes the standard non-relativistic form

$$H = \frac{p^2}{2m} + V(r). \quad (8.2)$$

[8.1: Derivation of Radial Equation](#)

[8.2: Infinite Spherical Potential Well](#)

[8.3: Hydrogen Atom](#)

[8.4: Rydberg Formula](#)

[8.E: Central Potentials \(Exercises\)](#)

---

This page titled [8: Central Potentials](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 8.1: Derivation of Radial Equation

Now, we have seen that the Cartesian components of the momentum,  $\mathbf{p}$ , can be represented as (see Section [s7.2])

$$p_i = -i\hbar \frac{\partial}{\partial x_i} \quad (8.1.1)$$

for  $i = 1, 2, 3$ , where  $x_1 \equiv x$ ,  $x_2 \equiv y$ ,  $x_3 \equiv z$ , and  $\mathbf{r} \equiv (x_1, x_2, x_3)$ . Likewise, it is easily demonstrated, from the previous expressions, and the basic definitions of the spherical coordinates [see Equations ([e8.21])–([e8zz])], that the radial component of the momentum can be represented as

$$p_r \equiv \frac{\mathbf{p} \cdot \mathbf{r}}{r} = -i\hbar \frac{\partial}{\partial r} \quad (8.1.2)$$

Recall that the angular momentum vector,  $\mathbf{L}$ , is defined

$$\mathbf{L} = \mathbf{r} \times \mathbf{p} \quad (8.1.3)$$

[See Equation ([e8.0]).] This expression can also be written in the following form:

$$L_i = \epsilon_{ijk} x_j p_k. \quad (8.1.4)$$

Here, the  $\epsilon_{ijk}$  (where  $i, j, k$  all run from 1 to 3) are elements of the so-called *totally anti-symmetric tensor*. The values of the various elements of this tensor are determined via a simple rule:

$$\epsilon_{ijk} = \begin{cases} 0 & \text{if } i, j, k \text{ not all different} \\ 1 & \text{if } i, j, k \text{ are cyclic permutation of } 1, 2, 3 \\ -1 & \text{if } i, j, k \text{ are anti-cyclic permutation of } 1, 2, 3 \end{cases} \quad (8.1.5)$$

Thus,  $\epsilon_{123} = \epsilon_{231} = 1$ ,  $\epsilon_{321} = \epsilon_{132} = -1$ , and  $\epsilon_{112} = \epsilon_{131} = 0$ , et cetera. Equation ([e9.6]) also makes use of the *Einstein summation convention*, according to which repeated indices are summed (from 1 to 3). For instance,  $a_i b_i \equiv a_1 b_1 + a_2 b_2 + a_3 b_3$ . Making use of this convention, as well as Equation ([e9.7]), it is easily seen that Equations ([e9.5]) and ([e9.6]) are indeed equivalent.

Let us calculate the value of  $L^2$  using Equation ([e9.6]). According to our new notation,  $L^2$  is the same as  $L_i L_i$ . Thus, we obtain

$$L^2 = \epsilon_{ijk} x_j p_k \epsilon_{ilm} x_l p_m = \epsilon_{ijk} \epsilon_{ilm} x_j p_k x_l p_m. \quad (8.1.6)$$

Note that we are able to shift the position of  $\epsilon_{ilm}$  because its elements are just numbers, and, therefore, commute with all of the  $x_i$  and the  $p_i$ . Now, it is easily demonstrated that

$$\epsilon_{ijk} \epsilon_{ilm} \equiv \delta_{jl} \delta_{km} - \delta_{jm} \delta_{kl}. \quad (8.1.7)$$

Here  $\delta_{ij}$  is the usual *Kronecker delta*, whose elements are determined according to the rule

$$\delta_{ij} = \begin{cases} 1 & \text{if } i \text{ and } j \text{ the same} \\ 0 & \text{if } i \text{ and } j \text{ different} \end{cases} \quad (8.1.8)$$

It follows from Equations ([e9.8]) and ([e9.9]) that

$$L^2 = x_i p_j x_i p_j - x_i p_j x_j p_i. \quad (8.1.9)$$

Here, we have made use of the fairly self-evident result that  $\delta_{ij} a_i b_j \equiv a_i b_i$ . We have also been careful to preserve the order of the various terms on the right-hand side of the previous expression, because the  $x_i$  and the  $p_i$  do not necessarily commute with one another.

We now need to rearrange the order of the terms on the right-hand side of Equation ([e9.11]). We can achieve this goal by making use of the fundamental commutation relation for the  $x_i$  and the  $p_i$ :

$$[x_i, p_j] = i\hbar \delta_{ij}. \quad (8.1.10)$$

[See Equation ([commxp]).] Thus,



$$\begin{aligned}
 L^2 &= x_i (x_i p_j - [x_i, p_j]) p_j - x_i p_j (p_i x_j + [x_j, p_i]) \\
 &= x_i x_i p_j p_j - i \hbar \delta_{ij} x_i p_j - x_i p_j p_i x_j - i \hbar \delta_{ij} x_i p_j \\
 &= x_i x_i p_j p_j - x_i p_i p_j x_j - 2 i \hbar x_i p_i.
 \end{aligned}$$

Here, we have made use of the fact that  $p_j p_i = p_i p_j$ , because the  $p_i$  commute with one another. [See Equation ([\[commpp\]](#)).] Next,

$$L^2 = x_i x_i p_j p_j - x_i p_i (x_j p_j - [x_j, p_j]) - 2 i \hbar x_i p_i. \quad (8.1.11)$$

Now, according to Equation ([\[e9.12\]](#)),

$$[x_j, p_j] \equiv [x_1, p_1] + [x_2, p_2] + [x_3, p_3] = 3 i \hbar. \quad (8.1.12)$$

Hence, we obtain

$$L^2 = x_i x_i p_j p_j - x_i p_i x_j p_j + i \hbar x_i p_i. \quad (8.1.13)$$

When expressed in more conventional vector notation, the previous expression becomes

$$L^2 = r^2 p^2 - (\mathbf{r} \cdot \mathbf{p})^2 + i \hbar \mathbf{r} \cdot \mathbf{p}. \quad (8.1.14)$$

Note that if we had attempted to derive the previous expression directly from Equation ([\[e9.5\]](#)), using standard vector identities, then we would have missed the final term on the right-hand side. This term originates from the lack of commutation between the  $x_i$  and  $p_i$  operators in quantum mechanics. Of course, standard vector analysis assumes that all terms commute with one another.

Equation ([\[e9.17\]](#)) can be rearranged to give

$$p^2 = r^{-2} [(\mathbf{r} \cdot \mathbf{p})^2 - i \hbar \mathbf{r} \cdot \mathbf{p} + L^2]. \quad (8.1.15)$$

Now,

$$\mathbf{r} \cdot \mathbf{p} = r p_r = -i \hbar r \frac{\partial}{\partial r}, \quad (8.1.16)$$

where use has been made of Equation ([\[e9.4\]](#)). Hence, we obtain

$$p^2 = -\hbar^2 \left[ \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial}{\partial r} \right) + \frac{1}{r} \frac{\partial}{\partial r} - \frac{L^2}{\hbar^2 r^2} \right]. \quad (8.1.17)$$

Finally, the previous equation can be combined with Equation ([\[e9.2\]](#)) to give the following expression for the Hamiltonian:

$$H = -\frac{\hbar^2}{2m} \left( \frac{\partial^2}{\partial r^2} + \frac{2}{r} \frac{\partial}{\partial r} - \frac{L^2}{\hbar^2 r^2} \right) + V(r). \quad (8.1.18)$$

Let us now consider whether the previous Hamiltonian commutes with the angular momentum operators  $L_z$  and  $L^2$ . Recall, from Section [\[s8.3\]](#), that  $L_z$  and  $L^2$  are represented as differential operators that depend solely on the angular spherical coordinates,  $\theta$  and  $\phi$ , and do not contain the radial coordinate,  $r$ . Thus, any function of  $r$ , or any differential operator involving  $r$  (but not  $\theta$  and  $\phi$ ), will automatically commute with  $L^2$  and  $L_z$ . Moreover,  $L^2$  commutes both with itself, and with  $L_z$ . (See Section [\[s8.2\]](#).) It is, therefore, clear that the previous Hamiltonian commutes with both  $L_z$  and  $L^2$ .

According to Section [\[smeas\]](#), if two operators commute with one another then they possess simultaneous eigenstates. We thus conclude that for a particle moving in a central potential the eigenstates of the Hamiltonian are simultaneous eigenstates of  $L_z$  and  $L^2$ . Now, we have already found the simultaneous eigenstates of  $L_z$  and  $L^2$ —they are the spherical harmonics,  $Y_{l,m}(\theta, \phi)$ , discussed in Section [\[sharm\]](#). It follows that the spherical harmonics are also eigenstates of the Hamiltonian. This observation leads us to try the following separable form for the stationary wavefunction:

$$\psi(r, \theta, \phi) = R(r) Y_{l,m}(\theta, \phi). \quad (8.1.19)$$

It immediately follows, from Equation ([\[e8.29\]](#)) and ([\[e8.30\]](#)), and the fact that  $L_z$  and  $L^2$  both obviously commute with  $R(r)$ , that

$$\begin{aligned}
 L_z \psi &= m \hbar \psi, \\
 L^2 \psi &= l(l+1) \hbar^2 \psi.
 \end{aligned}$$

Recall that the quantum numbers  $m$  and  $l$  are restricted to take certain integer values, as explained in Section [\[slsq\]](#).

Finally, making use of Equations ([e9.1]), ([e9.21]), and ([e9.24]), we obtain the following differential equation which determines the radial variation of the stationary wavefunction:

$$-\frac{\hbar^2}{2m} \left[ \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} - \frac{l(l+1)}{r^2} \right] R_{n,l} + V R_{n,l} = E R_{n,l}. \quad (8.1.20)$$

Here, we have labeled the function  $R(r)$  by two quantum numbers,  $n$  and  $l$ . The second quantum number,  $l$ , is, of course, related to the eigenvalue of  $L^2$ . [Note that the azimuthal quantum number,  $m$ , does not appear in the previous equation, and, therefore, does not influence either the function  $R(r)$  or the energy,  $E$ .] As we shall see, the first quantum number,  $n$ , is determined by the constraint that the radial wavefunction be square-integrable.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [8.1: Derivation of Radial Equation](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 8.2: Infinite Spherical Potential Well

Consider a particle of mass  $m$  and energy  $E > 0$  moving in the following simple central potential:

$$V(r) = \begin{cases} 0 & \text{for } 0 \leq r \leq a \\ \infty & \text{otherwise} \end{cases}. \quad (8.2.1)$$

Clearly, the wavefunction  $\psi$  is only non-zero in the region  $0 \leq r \leq a$ . Within this region, it is subject to the physical boundary conditions that it be well behaved (i.e., square-integrable) at  $r = 0$ , and that it be zero at  $r = a$ . (See Section [\[s5.2\]](#).) Writing the wavefunction in the standard form

$$\psi(r, \theta, \phi) = R_{n,l}(r) Y_{l,m}(\theta, \phi), \quad (8.2.2)$$

we deduce (see the previous section) that the radial function  $R_{n,l}(r)$  satisfies

$$\frac{d^2 R_{n,l}}{dr^2} + \frac{2}{r} \frac{dR_{n,l}}{dr} + \left[ k^2 - \frac{l(l+1)}{r^2} \right] R_{n,l} = 0 \quad (8.2.3)$$

in the region  $0 \leq r \leq a$ , where

$$k^2 = \frac{2mE}{\hbar^2}. \quad (8.2.4)$$

Defining the scaled radial variable  $z = kr$ , the previous differential equation can

be transformed into the standard form

$$\left[ \frac{d^2}{dz^2} R_{n,l} \right] + \left[ 2 - \frac{l(l+1)}{z^2} \right] R_{n,l} = 0.$$

The two independent solutions to this well-known second-order differential equation are called *spherical Bessel functions*, and can be written

$$j_l(z) = z^l \left( -\frac{1}{z} \frac{d}{dz} \right)^l \left( \frac{\sin z}{z} \right),$$

$$y_l(z) = -z^l \left( -\frac{1}{z} \frac{d}{dz} \right)^l \left( \frac{\cos z}{z} \right).$$

Thus, the first few spherical Bessel functions take the form

$$j_0(z) = \frac{\sin z}{z},$$

$$j_1(z) = \frac{\sin z}{z^2} - \frac{\cos z}{z},$$

$$y_0(z) = -\frac{\cos z}{z},$$

$$y_1(z) = -\frac{\cos z}{z^2} - \frac{\sin z}{z}.$$

These functions are also plotted in Figure [\[sph\]](#). It can be seen that the spherical Bessel functions are oscillatory in nature, passing through zero many times. However, the  $y_l(z)$  functions are badly behaved (i.e., they are not square integrable) at  $z = 0$ , whereas the  $j_l(z)$  functions are well behaved everywhere. It follows from our boundary condition at  $r = 0$  that the  $y_l(z)$  are unphysical, and that the radial wavefunction  $R_{n,l}(r)$  is thus proportional to  $j_l(kr)$  only. In order to satisfy the boundary condition at  $r = a$  [i.e.,  $R_{n,l}(a) = 0$ ], the value of  $k$  must be chosen such that  $z = ka$  corresponds to one of the zeros of  $j_l(z)$ . Let us denote the  $n$ th zero of  $j_l(z)$  as  $z_{n,l}$ . It follows that

$$ka = z_{n,l}, \quad (8.2.5)$$

for  $n = 1, 2, 3, \dots$ . Hence, from Equation ([\[e9.29\]](#)), the allowed energy levels are

$$E_{n,l} = z_{n,l}^2 \frac{\hbar^2}{2 m a^2}. \quad (8.2.6)$$

The first few values of  $z_{n,l}$  are listed in Table [tsph]. It can be seen that  $z_{n,l}$  is an increasing function of both  $n$  and  $l$ .

The first few zeros of the spherical Bessel function  $j_l(z)$ .

	$n = 1$	$n = 2$	$n = 3$	$n = 4$
$l = 0$	3.142	6.283	9.425	12.566
[0.5ex] $l = 1$	4.493	7.725	10.904	14.066
[0.5ex] $l = 2$	5.763	9.095	12.323	15.515
[0.5ex] $l = 3$	6.988	10.417	13.698	16.924
[0.5ex] $l = 4$	8.183	11.705	15.040	18.301

We are now in a position to interpret the three quantum numbers— $n$ ,  $l$ , and  $m$ —which determine the form of the wavefunction specified in Equation ([e9.27]). As is clear from Chapter [sorb], the azimuthal quantum number  $m$  determines the number of nodes in the wavefunction as the azimuthal angle  $\phi$  varies between 0 and  $2\pi$ . Thus,  $m = 0$  corresponds to no nodes,  $m = 1$  to a single node,  $m = 2$  to two nodes, et cetera. Likewise, the polar quantum number  $l$  determines the number of nodes in the wavefunction as the polar angle  $\theta$  varies between 0 and  $\pi$ . Again,  $l = 0$  corresponds to no nodes,  $l = 1$  to a single node, et cetera. Finally, the radial quantum number  $n$  determines the number of nodes in the wavefunction as the radial variable  $r$  varies between 0 and  $a$  (not counting any nodes at  $r = 0$  or  $r = a$ ). Thus,  $n = 1$  corresponds to no nodes,  $n = 2$  to a single node,  $n = 3$  to two nodes, et cetera. Note that, for the case of an infinite potential well, the only restrictions on the values that the various quantum numbers can take are that  $n$  must be a positive integer,  $l$  must be a non-negative integer, and  $m$  must be an integer lying between  $-l$  and  $l$ . Note, further, that the allowed energy levels ([e9.39]) only depend on the values of the quantum numbers  $n$  and  $l$ . Finally, it is easily demonstrated that the spherical Bessel functions are mutually orthogonal: that is,

$$\int_0^a j_l(z_{n,l} r/a) j_l(z_{n',l} r/a) r^2 dr = 0 \quad (8.2.7)$$

when  $n \neq n'$ . Given that the  $Y_{l,m}(\theta, \phi)$  are mutually orthogonal (see Chapter [sorb]), this ensures that wavefunctions ([e9.27]) corresponding to distinct sets of values of the quantum numbers  $n$ ,  $l$ , and  $m$  are mutually orthogonal.

## Contributors and Attributions

- Richard Fitzpatrick (Professor of Physics, The University of Texas at Austin)

This page titled 8.2: Infinite Spherical Potential Well is shared under a not declared license and was authored, remixed, and/or curated by Richard Fitzpatrick.

## 8.3: Hydrogen Atom

A hydrogen atom consists of an electron, of charge  $-e$  and mass  $m_e$ , and a proton, of charge  $+e$  and mass  $m_p$ , moving in the Coulomb potential

$$V(\mathbf{r}) = -\frac{e^2}{4\pi\epsilon_0 |\mathbf{r}|}, \quad (8.3.1)$$

where  $\mathbf{r}$  is the position vector of the electron with respect to the proton. Now, according to the analysis in Section [stwo], this two-body problem can be converted into an equivalent one-body problem. In the latter problem, a particle of mass

$$\mu = \frac{m_e m_p}{m_e + m_p} \quad (8.3.2)$$

moves in the central potential

$$V(r) = -\frac{e^2}{4\pi\epsilon_0 r}. \quad (8.3.3)$$

Note, however, that because  $m_e/m_p \simeq 1/1836$  the difference between  $m_e$  and  $\mu$  is very small. Hence, in the following, we shall write neglect this difference entirely.

Writing the wavefunction in the usual form,

$$\psi(r, \theta, \phi) = R_{n,l}(r) Y_{l,m}(\theta, \phi), \quad (8.3.4)$$

it follows from Section 1.2 that the radial function  $R_{n,l}(r)$  satisfies

$$-\frac{\hbar^2}{2m_e} \left[ \frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} - \frac{l(l+1)}{r^2} \right] R_{n,l} - \left( \frac{e^2}{4\pi\epsilon_0 r} + E \right) R_{n,l} = 0. \quad (8.3.5)$$

Let  $r = az$ , with

$$a = \sqrt{\frac{\hbar^2}{2m_e(-E)}} = \sqrt{\frac{E_0}{E}} a_0, \quad (8.3.6)$$

where  $E_0$  and  $a_0$  are defined in Equations ([e9.56]) and ([e9.57]), respectively. Here, it is assumed that  $E < 0$ , because we are only interested in bound-states of the hydrogen atom. The previous differential equation transforms to

$$\left[ \frac{d^2}{dz^2} + \frac{2}{z} \frac{d}{dz} - \frac{l(l+1)}{z^2} + \frac{\zeta}{z} - 1 \right] R_{n,l} = 0, \quad (8.3.7)$$

where

$$\zeta = \frac{2m_e a e^2}{4\pi\epsilon_0 \hbar^2} = 2\sqrt{\frac{E_0}{E}}. \quad (8.3.8)$$

Suppose that  $R_{n,l}(r) = Z(r/a) \exp(-r/a)/(r/a)$ . It follows that

$$\left[ \frac{d^2}{dz^2} - 2 \frac{d}{dz} - \frac{l(l+1)}{z^2} + \frac{\zeta}{z} \right] Z = 0. \quad (8.3.9)$$

We now need to solve the previous differential equation in the domain  $z = 0$  to  $z = \infty$ , subject to the constraint that  $R_{n,l}(r)$  be square-integrable.

Let us look for a power-law solution of the form

$$Z(z) = \sum_k c_k z^k. \quad (8.3.10)$$

Substituting this solution into Equation ([e9.48]), we obtain

$$\sum_k c_k \{ k(k-1) z^{k-2} - 2k z^{k-1} - l(l+1) z^{k-2} + \zeta z^{k-1} \} = 0. \quad (8.3.11)$$

Equating the coefficients of  $z^{k-2}$  gives the recursion relation

$$c_k [k(k-1) - l(l+1)] = c_{k-1} [2(k-1) - \zeta]. \quad (8.3.12)$$

Now, the power series ([e9.49]) must terminate at small  $k$ , at some positive value of  $k$ , otherwise  $Z(z)$  behaves unphysically as  $z \rightarrow 0$  [i.e., it yields an  $R_{n,l}(r)$  that is not square integrable as  $r \rightarrow 0$ ]. From the previous recursion relation, this is only possible if  $[k_{\min}(k_{\min}-1) - l(l+1)] = 0$ , where the first term in the series is  $c_{k_{\min}} z^{k_{\min}}$ . There are two possibilities:  $k_{\min} = -l$  or  $k_{\min} = l+1$ . However, the former possibility predicts unphysical behavior of  $Z(z)$  at  $z=0$ . Thus, we conclude that  $k_{\min} = l+1$ . Note that, because  $R_{n,l}(r) \simeq Z(r/a)/(r/a) \simeq (r/a)^l$  at small  $r$ , there is a finite probability of finding the electron at the nucleus for an  $l=0$  state, whereas there is zero probability of finding the electron at the nucleus for an  $l>0$  state [i.e.,  $|\psi|^2 = 0$  at  $r=0$ , except when  $l=0$ ].

For large values of  $z$ , the ratio of successive coefficients in the power series ([e9.49]) is

$$\frac{c_k}{c_{k-1}} = \frac{2}{k}, \quad (8.3.13)$$

according to Equation ([e9.51]). This is the same as the ratio of successive coefficients in the power series

$$\sum_k \frac{(2z)^k}{k!}, \quad (8.3.14)$$

which converges to  $\exp(2z)$ . We conclude that  $Z(z) \rightarrow \exp(2z)$  as  $z \rightarrow \infty$ . It thus follows that  $R_{n,l}(r) \sim Z(r/a) \exp(-r/a)/(r/a) \rightarrow \exp(r/a)/(r/a)$  as  $r \rightarrow \infty$ . This does not correspond to physically acceptable behavior of the wavefunction, because  $\int |\psi|^2 dV$  must be finite. The only way in which we can avoid this unphysical behavior is if the power series ([e9.49]) terminates at some maximum value of  $k$ . According to the recursion relation ([e9.51]), this is only possible if

$$\frac{\zeta}{2} = n, \quad (8.3.15)$$

where  $n$  is an integer, and the last term in the series is  $c_n z^n$ . Because the first term in the series is  $c_{l+1} z^{l+1}$ , it follows that  $n$  must be greater than  $l$ , otherwise there are no terms in the series at all. Finally, it is clear from Equations ([e9.45]), ([e9.47]), and ([e9.54]) that

$$E = \frac{E_0}{n^2} \quad (8.3.16)$$

and

$$a = n a_0, \quad (8.3.17)$$

where

$$E_0 = -\frac{m_e e^4}{2(4\pi\epsilon_0)^2 \hbar^2} = -\frac{e^2}{8\pi\epsilon_0 a_0} = -13.6 \text{ eV}, \quad (8.3.18)$$

and

$$a_0 = \frac{4\pi\epsilon_0 \hbar^2}{m_e e^2} = 5.3 \times 10^{-11} \text{ m}. \quad (8.3.19)$$

Here,  $E_0$  is the energy of so-called *ground-state* (or lowest energy state) of the hydrogen atom, and the length  $a_0$  is known as the *Bohr radius*. Note that  $|E_0| \sim \alpha^2 m_e c^2$ , where  $\alpha = e^2/(4\pi\epsilon_0 \hbar c) \simeq 1/137$  is the dimensionless *fine-structure constant*. The fact that  $|E_0| \ll m_e c^2$  is the ultimate justification for our non-relativistic treatment of the hydrogen atom.

We conclude that the wavefunction of a hydrogen atom takes the form

$$\psi_{n,l,m}(r, \theta, \phi) = R_{n,l}(r) Y_{l,m}(\theta, \phi). \quad (8.3.20)$$

Here, the  $Y_{l,m}(\theta, \phi)$  are the spherical harmonics (see Section [sharm]), and  $R_{n,l}(z = r/a)$  is the solution of

$$\left[ \frac{1}{z^2} \frac{d}{dz} z^2 \frac{d}{dz} - \frac{l(l+1)}{z^2} + \frac{2n}{z} - 1 \right] R_{n,l} = 0 \quad (8.3.21)$$

which varies as  $z^l$  at small  $z$ . Furthermore, the quantum numbers  $n$ ,  $l$ , and  $m$  can only take values that satisfy the inequality

$$|m| \leq l < n, \quad (8.3.22)$$

where  $n$  is a positive integer,  $l$  a non-negative integer, and  $m$  an integer.

We expect the stationary states of the hydrogen atom to be orthonormal: that is,

$$\int \psi_{n',l',m'}^* \psi_{n,l,m} dV = \delta_{nn'} \delta_{ll'} \delta_{mm'}, \quad (8.3.23)$$

where  $dV$  is a volume element, and the integral is over all space. Of course,  $dV = r^2 dr d\Omega$ , where  $d\Omega$  is an element of solid angle. Moreover, we already know that the spherical harmonics are orthonormal [see Equation ([\[spho\]](#))]: that is,

$$\oint Y_{l',m'}^* Y_{l,m} d\Omega = \delta_{ll'} \delta_{mm'}. \quad (8.3.24)$$

It, thus, follows that the radial wavefunction satisfies the orthonormality constraint

$$\int_0^\infty R_{n',l}^* R_{n,l} r^2 dr = \delta_{nn'}. \quad (8.3.25)$$

The first few radial wavefunctions for the hydrogen atom are listed below:

$$\begin{aligned} R_{1,0}(r) &= \frac{2}{a_0^{3/2}} \exp\left(-\frac{r}{a_0}\right), \\ R_{2,0}(r) &= \frac{2}{(2a_0)^{3/2}} \left(1 - \frac{r}{2a_0}\right) \exp\left(-\frac{r}{2a_0}\right), \\ R_{2,1}(r) &= \frac{1}{\sqrt{3} (2a_0)^{3/2}} \frac{r}{a_0} \exp\left(-\frac{r}{2a_0}\right), \\ R_{3,0}(r) &= \frac{2}{(3a_0)^{3/2}} \left(1 - \frac{2r}{3a_0} + \frac{2r^2}{27a_0^2}\right) \exp\left(-\frac{r}{3a_0}\right), \\ R_{3,1}(r) &= \frac{4\sqrt{2}}{9 (3a_0)^{3/2}} \frac{r}{a_0} \left(1 - \frac{r}{6a_0}\right) \exp\left(-\frac{r}{3a_0}\right), \\ R_{3,2}(r) &= \frac{2\sqrt{2}}{27\sqrt{5} (3a_0)^{3/2}} \left(\frac{r}{a_0}\right)^2 \exp\left(-\frac{r}{3a_0}\right). \end{aligned}$$

These functions are illustrated in Figures [\[coul1\]](#) and [\[coul2\]](#).




Figure 21: The  $a_0 r^2 |R_{n,l}(r)|^2$  plotted as a functions of  $r/a_0$ . The solid, short-dashed, and long-dashed curves correspond to  $n, l = 1, 0$ , and  $2, 0$ , and  $2, 1$ , respectively.




Figure 22: The  $a_0 r^2 |R_{n,l}(r)|^2$  plotted as a functions of  $r/a_0$ . The solid, short-dashed, and long-dashed curves correspond to  $n, l = 3, 0$ , and  $3, 1$ , and  $3, 2$ , respectively.

Given the (properly normalized) hydrogen wavefunction ([\[e9.59\]](#)), plus our interpretation of  $|\psi|^2$  as a probability density, we can calculate



$$\langle r^k \rangle = \int_0^\infty r^{2+k} |R_{n,l}(r)|^2 dr, \quad (8.3.26)$$

where the angle-brackets denote an expectation value. For instance, it can be demonstrated (after much tedious algebra) that

$$\begin{aligned} \langle r^2 \rangle &= \frac{a_0^2 n^2}{2} [5n^2 + 1 - 3l(l+1)], \\ \langle r \rangle &= \frac{a_0}{2} [3n^2 - l(l+1)], \\ \left\langle \frac{1}{r} \right\rangle &= \frac{1}{n^2 a_0}, \\ \left\langle \frac{1}{r^2} \right\rangle &= \frac{1}{(l+1/2)n^3 a_0^2}, \\ \left\langle \frac{1}{r^3} \right\rangle &= \frac{1}{l(l+1/2)(l+1)n^3 a_0^3}. \end{aligned}$$

According to Equation ([e9.55]), the energy levels of the bound-states of a hydrogen atom only depend on the radial quantum number  $n$ . It turns out that this is a special property of a  $1/r$  potential. For a general central potential,  $V(r)$ , the quantized energy levels of a bound-state depend on both  $n$  and  $l$ . (See Section 1.3.)

The fact that the energy levels of a hydrogen atom only depend on  $n$ , and not on  $l$  and  $m$ , implies that the energy spectrum of a hydrogen atom is highly degenerate: that is, there are many different states which possess the same energy. According to the inequality ([e9.61]) (and the fact that  $n$ ,  $l$ , and  $m$  are integers), for a given value of  $l$ , there are  $2l+1$  different allowed values of  $m$  (i.e.,  $-l, -l+1, \dots, l-1, l$ ). Likewise, for a given value of  $n$ , there are  $n$  different allowed values of  $l$  (i.e.,  $0, 1, \dots, n-1$ ). Now, all states possessing the same value of  $n$  have the same energy (i.e., they are degenerate). Hence, the total number of degenerate states corresponding to a given value of  $n$  is

$$1 + 3 + 5 + \dots + 2(n-1) + 1 = n^2. \quad (8.3.27)$$

Thus, the ground-state ( $n=1$ ) is not degenerate, the first excited state ( $n=2$ ) is four-fold degenerate, the second excited state ( $n=3$ ) is nine-fold degenerate, et cetera (Actually, when we take into account the two spin states of an electron, the degeneracy of the  $n$ th energy level becomes  $2n^2$ .)

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [8.3: Hydrogen Atom](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 8.4: Rydberg Formula

An electron in a given stationary state of a hydrogen atom, characterized by the quantum numbers  $n$ ,  $l$ , and  $m$ , should, in principle, remain in that state indefinitely. In practice, if the state is slightly perturbed—for instance, via interaction with a photon—then the electron can make a transition to another stationary state with different quantum numbers. (See Chapter [\[s13\]](#).)

Suppose that an electron in a hydrogen atom makes a transition from an initial state whose radial quantum number is  $n_i$  to a final state whose radial quantum number is  $n_f$ . According to Equation ([\[e9.55\]](#)), the energy of the electron will change by

$$\Delta E = E_0 \left( \frac{1}{n_f^2} - \frac{1}{n_i^2} \right). \quad (8.4.1)$$

If  $\Delta E$  is negative then we would expect the electron to emit a photon of frequency  $\nu = -\Delta E/h$ . [See Equation ([\[ee3.15\]](#)).] Likewise, if  $\Delta E$  is positive then the electron must absorb a photon of energy  $\nu = \Delta E/h$ . Given that  $\lambda^{-1} = \nu/c$ , the possible wavelengths of the photons emitted by a hydrogen atom as its electron makes transitions between different energy levels are

$$\frac{1}{\lambda} = R \left( \frac{1}{n_f^2} - \frac{1}{n_i^2} \right), \quad (8.4.2)$$

where

$$R = \frac{-E_0}{h c} = \frac{m_e e^4}{(4\pi)^3 \epsilon_0^2 \hbar^3 c} = 1.097 \times 10^7 \text{ m}^{-1}. \quad (8.4.3)$$

Here, it is assumed that  $n_f < n_i$ . Note that the emission spectrum of hydrogen is quantized: that is, a hydrogen atom can only emit photons with certain fixed set of wavelengths. Likewise, a hydrogen atom can only absorb photons that have the same fixed set of wavelengths. This set of wavelengths constitutes the characteristic emission/absorption spectrum of the hydrogen atom, and can be observed as “spectral lines” using a spectroscope.

Equation ([\[e9.77\]](#)) is known as the *Rydberg formula*. Likewise,  $R$  is called the *Rydberg constant*. The Rydberg formula was actually discovered empirically in the nineteenth century by spectroscopists, and was first explained theoretically by Bohr in 1913 using a primitive version of quantum mechanics. Transitions to the ground-state ( $n_f = 1$ ) give rise to spectral lines in the ultraviolet band—this set of lines is called the *Lyman series*. Transitions to the first excited state ( $n_f = 2$ ) give rise to spectral lines in the visible band—this set of lines is called the *Balmer series*. Transitions to the second excited state ( $n_f = 3$ ) give rise to spectral lines in the infrared band—this set of lines is called the *Paschen series*, and so on.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [8.4: Rydberg Formula](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 8.E: Central Potentials (Exercises)

1. A particle of mass  $m$  is placed in a finite spherical well:

$$V(r) = \begin{cases} -V_0 & \text{for } r \leq a \\ 0 & \text{for } r > a \end{cases} \quad (8.E.1)$$

with  $V_0 > 0$  and  $a > 0$ . Find the ground-state by solving the radial equation with  $l = 0$ . Show that there is no ground-state if  $V_0 a^2 < \pi^2 \hbar^2 / (8m)$ .

2. Consider a particle of mass  $m$  in the three-dimensional harmonic oscillator potential  $V(r) = (1/2) m \omega^2 r^2$ . Solve the problem by separation of variables in spherical coordinates, and, hence, determine the energy eigenvalues of the system.
3. The normalized wavefunction for the ground-state of a hydrogen-like atom (neutral hydrogen,  $\text{He}^+$ ,  $\text{Li}^{++}$ , et cetera.) with nuclear charge  $Ze$  has the form

$$\psi = A \exp(-\beta r), \quad (8.E.2)$$

where  $A$  and  $\beta$  are constants, and  $r$  is the distance between the nucleus and the electron. Show the following:

1.  $A^2 = \beta^3 / \pi$ .
  2.  $\beta = Z/a_0$ , where  $a_0 = (\hbar^2 / m_e) (4\pi \epsilon_0 / e^2)$ .
  3. The energy is  $E = -Z^2 E_0$  where  $E_0 = (m_e / 2 \hbar^2) (e^2 / 4\pi \epsilon_0)^2$ .
  4. The expectation values of the potential and kinetic energies are  $2E$  and  $-E$ , respectively.
  5. The expectation value of  $r$  is  $(3/2) (a_0 / Z)$ .
  6. The most probable value of  $r$  is  $a_0 / Z$ .
4. An atom of tritium is in its ground-state. Suddenly the nucleus decays into a helium nucleus, via the emission of a fast electron that leaves the atom without perturbing the extranuclear electron, Find the probability that the resulting  $\text{He}^+$  ion will be left in an  $n = 1, l = 0$  state. Find the probability that it will be left in a  $n = 2, l = 0$  state. What is the probability that the ion will be left in an  $l > 0$  state?
  5. Calculate the wavelengths of the photons emitted from the  $n = 2, l = 1$  to  $n = 1, l = 0$  transition in hydrogen, deuterium, and positronium.
  6. To conserve linear momentum, an atom emitting a photon must recoil, which means that not all of the energy made available in the downward jump goes to the photon. Find a hydrogen atom's recoil energy when it emits a photon in an  $n = 2$  to  $n = 1$  transition. What fraction of the transition energy is the recoil energy?

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [8.E: Central Potentials \(Exercises\)](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## CHAPTER OVERVIEW

### 9: Spin Angular Momentum

Broadly speaking, a classical extended object (e.g., the Earth) can possess two different types of angular momentum. The first type is due to the rotation of the object's center of mass about some fixed external point (e.g., the Sun)—this is generally known as *orbital angular momentum*. The second type is due to the object's internal motion—this is generally known as *spin angular momentum* (because, for a rigid object, the internal motion consists of spinning about an axis passing through the center of mass). By analogy, quantum particles can possess both orbital angular momentum due to their motion through space (see Chapter [\[sorb\]](#)), and spin angular momentum due to their internal motion. Actually, the analogy with classical extended objects is not entirely accurate, because electrons, for instance, are structureless point particles. In fact, in quantum mechanics, it is best to think of spin angular momentum as a kind of intrinsic angular momentum possessed by particles. It turns out that each type of elementary particle has a characteristic spin angular momentum, just as each type has a characteristic charge and mass.

[9.1: Spin Operators](#)

[9.2: Spin Space](#)

[9.3: Eigenstates of  \$S\_z\$  and  \$S^2\$](#)

[9.4: Pauli Representation](#)

[9.5: Spin Precession](#)

[9.E: Spin Angular Momentum \(Exercises\)](#)

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [9: Spin Angular Momentum](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 9.1: Spin Operators

Because spin is a type of angular momentum, it is reasonable to suppose that it possesses similar properties to orbital angular momentum. Thus, by analogy with Section [s8.2], we would expect to be able to define three operators— $S_x$ ,  $S_y$ , and  $S_z$ —that represent the three Cartesian components of spin angular momentum. Moreover, it is plausible that these operators possess analogous commutation relations to the three corresponding orbital angular momentum operators,  $L_x$ ,  $L_y$ , and  $L_z$ . [See Equations ([e8.6])–([e8.8]).] In other words,

$$\begin{aligned}[S_x, S_y] &= i\hbar S_z, \\ [S_y, S_z] &= i\hbar S_x, \\ [S_z, S_x] &= i\hbar S_y.\end{aligned}$$

We can represent the magnitude squared of the spin angular momentum vector by the operator

$$S^2 = S_x^2 + S_y^2 + S_z^2. \quad (9.1.1)$$

By analogy with the analysis in Section [s8.2], it is easily demonstrated that

$$[S^2, S_x] = [S^2, S_y] = [S^2, S_z] = 0. \quad (9.1.2)$$

We thus conclude (see Section [smeas]) that we can simultaneously measure the magnitude squared of the spin angular momentum vector, together with, at most, one Cartesian component. By convention, we shall always choose to measure the  $z$ -component,  $S_z$ .

By analogy with Equation ([e8.13]), we can define raising and lowering operators for spin angular momentum:

$$S_{\pm} = S_x \pm i S_y. \quad (9.1.3)$$

If  $S_x$ ,  $S_y$ , and  $S_z$  are Hermitian operators, as must be the case if they are to represent physical quantities, then  $S_{\pm}$  are the Hermitian conjugates of one another: that is,

$$(S_{\pm})^{\dagger} = S_{\mp}. \quad (9.1.4)$$

Finally, by analogy with Section [s8.2], it is easily demonstrated that

$$\begin{aligned}S_+ S_- &= S^2 - S_z^2 + \hbar S_z, \\ S_- S_+ &= S^2 - S_z^2 - \hbar S_z, \\ [S_+, S_z] &= -\hbar S_+, \\ [S_-, S_z] &= +\hbar S_-.\end{aligned}$$

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled 9.1: Spin Operators is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 9.2: Spin Space

We now have to discuss the wavefunctions upon which the previously introduced spin operators act. Unlike regular wavefunctions, spin wavefunctions do not exist in real space. Likewise, the spin angular momentum operators cannot be represented as differential operators in real space. Instead, we need to think of spin wavefunctions as existing in an abstract (complex) vector space. The different members of this space correspond to the different internal configurations of the particle under investigation. Note that only the directions of our vectors have any physical significance (just as only the shape of a regular wavefunction has any physical significance). Thus, if the vector  $\chi$  corresponds to a particular internal state then  $c\chi$  corresponds to the same state, where  $c$  is a complex number. Now, we expect the internal states of our particle to be superposable, because the superposability of states is one of the fundamental assumptions of quantum mechanics. It follows that the vectors making up our vector space must also be superposable. Thus, if  $\chi_1$  and  $\chi_2$  are two vectors corresponding to two different internal states then  $c_1\chi_1 + c_2\chi_2$  is another vector corresponding to the state obtained by superposing  $c_1$  times state 1 with  $c_2$  times state 2 (where  $c_1$  and  $c_2$  are complex numbers). Finally, the dimensionality of our vector space is simply the number of linearly independent vectors required to span it (i.e., the number of linearly independent internal states of the particle under investigation).

We now need to define the length of our vectors. We can do this by introducing a second, or *dual*, vector space whose elements are in one to one correspondence with the elements of our first space. Let the element of the second space that corresponds to the element  $\chi$  of the first space be called  $\chi^\dagger$ . Moreover, the element of the second space that corresponds to  $c\chi$  is  $c^*\chi^\dagger$ . We shall assume that it is possible to combine  $\chi$  and  $\chi^\dagger$  in a multiplicative fashion to generate a real positive-definite number that we shall interpret as the length, or *norm*, of  $\chi$ . Let us denote this number  $\chi^\dagger\chi$ . Thus, we have

$$\chi^\dagger\chi \geq 0 \quad (9.2.1)$$

for all  $\chi$ . We shall also assume that it is possible to combine unlike states in an analogous multiplicative fashion to produce complex numbers. The product of two unlike states  $\chi$  and  $\chi'$  is denoted  $\chi^\dagger\chi'$ . Two states  $\chi$  and  $\chi'$  are said to be mutually orthogonal, or independent, if  $\chi^\dagger\chi' = 0$ .

Now, when a general spin operator,  $A$ , operates on a general spin-state,  $\chi$ , it converts it into a different spin-state that we shall denote  $A\chi$ . The dual of this state is  $(A\chi)^\dagger \equiv \chi^\dagger A^\dagger$ , where  $A^\dagger$  is the Hermitian conjugate of  $A$  (this is the definition of an Hermitian conjugate in spin space). An eigenstate of  $A$  corresponding to the eigenvalue  $a$  satisfies

$$A\chi_a = a\chi_a. \quad (9.2.2)$$

As before, if  $A$  corresponds to a physical variable then a measurement of  $A$  will result in one of its eigenvalues. (See Section [smeas].) In order to ensure that these eigenvalues are all real,  $A$  must be Hermitian; that is,  $A^\dagger = A$ . (See Section [seig].) We expect the  $\chi_a$  to be mutually orthogonal. We can also normalize them such that they all have unit length. In other words,

$$\chi_a^\dagger\chi_{a'} = \delta_{aa'}. \quad (9.2.3)$$

Finally, a general spin state can be written as a superposition of the normalized eigenstates of  $A$ : that is,

$$\chi = \sum_a c_a \chi_a. \quad (9.2.4)$$

A measurement of  $\chi$  will then yield the result  $a$  with probability  $|c_a|^2$ .

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled 9.2: Spin Space is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 9.3: Eigenstates of $S_z$ and $S^2$

Because the operators  $S_z$  and  $S^2$  commute, they must possess simultaneous eigenstates. (See Section [smeas].) Let these eigenstates take the form [see Equations ([e8.29]) and ([e8.30])]:

$$\begin{aligned} S_z \chi_{s,m_s} &= m_s \hbar \chi_{s,m_s}, \\ S^2 \chi_{s,m_s} &= s(s+1) \hbar^2 \chi_{s,m_s}. \end{aligned}$$

Now, it is easily demonstrated, from the commutation relations ([e10.9]) and ([e10.10]), that

$$S_z (S_+ \chi_{s,m_s}) = (m_s + 1) \hbar (S_+ \chi_{s,m_s}), \quad (9.3.1)$$

and

$$S_z (S_- \chi_{s,m_s}) = (m_s - 1) \hbar (S_- \chi_{s,m_s}). \quad (9.3.2)$$

Thus,  $S_+$  and  $S_-$  are indeed the raising and lowering operators, respectively, for spin angular momentum. (See Section [seian].) The eigenstates of  $S_z$  and  $S^2$  are assumed to be orthonormal: that is,

$$\chi_{s,m_s}^\dagger \chi_{s',m'_s} = \delta_{ss'} \delta_{m_s m'_s}. \quad (9.3.3)$$

Consider the wavefunction  $\chi = S_+ \chi_{s,m_s}$ . Because we know, from Equation ([e10.11]), that  $\chi^\dagger \chi \geq 0$ , it follows that

$$(S_+ \chi_{s,m_s})^\dagger (S_+ \chi_{s,m_s}) = \chi_{s,m_s}^\dagger S_+^\dagger S_+ \chi_{s,m_s} = \chi_{s,m_s}^\dagger S_- S_+ \chi_{s,m_s} \geq 0, \quad (9.3.4)$$

where use has been made of Equation ([e10.7]). Equations ([e10.8]), ([e10.16]), ([e10.17]), and ([e10.20]) yield

$$s(s+1) \geq m_s(m_s+1). \quad (9.3.5)$$

Likewise, if  $\chi = S_- \chi_{s,m_s}$  then we obtain

$$s(s+1) \geq m_s(m_s-1). \quad (9.3.6)$$

Assuming that  $s \geq 0$ , the previous two inequalities imply that

$$-s \leq m_s \leq s. \quad (9.3.7)$$

Hence, at fixed  $s$ , there is both a maximum and a minimum possible value that  $m_s$  can take.

Let  $m_{s \min}$  be the minimum possible value of  $m_s$ . It follows that (see Section [slsq])

$$S_- \chi_{s,m_{s \min}} = 0. \quad (9.3.8)$$

Now, from Equation ([e10.7a]),

$$S^2 = S_+ S_- + S_z^2 - \hbar S_z. \quad (9.3.9)$$

Hence,

$$S^2 \chi_{s,m_{s \min}} = (S_+ S_- + S_z^2 - \hbar S_z) \chi_{s,m_{s \min}}, \quad (9.3.10)$$

giving

$$s(s+1) = m_{s \min}(m_{s \min} - 1). \quad (9.3.11)$$

Assuming that  $m_{s \min} < 0$ , this equation yields

$$m_{s \min} = -s. \quad (9.3.12)$$

Likewise, it is easily demonstrated that

$$m_{s \max} = +s. \quad (9.3.13)$$

Moreover,

$$S_- \chi_{s,-s} = S_+ \chi_{s,s} = 0. \quad (9.3.14)$$

Now, the raising operator  $S_+$ , acting upon  $\chi_{s,-s}$ , converts it into some multiple of  $\chi_{s,-s+1}$ . Employing the raising operator a second time, we obtain a multiple of  $\chi_{s,-s+2}$ . However, this process cannot continue indefinitely, because there is a maximum possible value of  $m_s$ . Indeed, after acting upon  $\chi_{s,-s}$  a sufficient number of times with the raising operator  $S_+$ , we must obtain a multiple of  $\chi_{s,s}$ , so that employing the raising operator one more time leads to the null state. [See Equation ([e10.31]).] If this is not the case then we will inevitably obtain eigenstates of  $S_z$  corresponding to  $m_s > s$ , which we have already demonstrated is impossible.

It follows, from the previous argument, that

$$m_{s \max} - m_{s \min} = 2s = k, \quad (9.3.15)$$

where  $k$  is a positive integer. Hence, the quantum number  $s$  can either take positive integer or positive half-integer values. Up to now, our analysis has been very similar to that which we used earlier to investigate orbital angular momentum. (See Section [sorb].) Recall, that for orbital angular momentum the quantum number  $m$ , which is analogous to  $m_s$ , is restricted to take integer values. (See Section [slz].) This implies that the quantum number  $l$ , which is analogous to  $s$ , is also restricted to take integer values. However, the origin of these restrictions is the representation of the orbital angular momentum operators as differential operators in real space. (See Section [s8.3].) There is no equivalent representation of the corresponding spin angular momentum operators. Hence, we conclude that there is no reason why the quantum number  $s$  cannot take half-integer, as well as integer, values.

In 1940, Wolfgang Pauli proved the so-called *spin-statistics theorem* using relativistic quantum mechanics. According to this theorem, all fermions possess half-integer spin (i.e., a half-integer value of  $s$ ), whereas all bosons possess integer spin (i.e., an integer value of  $s$ ). In fact, all presently known fermions, including electrons and protons, possess *spin one-half*. In other words, electrons and protons are characterized by  $s = 1/2$  and  $m_s = \pm 1/2$ .

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [9.3: Eigenstates of  \$S\_z\$  and  \$S^2\$](#)  is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).



## 9.4: Pauli Representation

Let us denote the two independent spin eigenstates of an electron as

$$\chi_{\pm} \equiv \chi_{1/2, \pm 1/2}. \quad (9.4.1)$$

It thus follows, from Equations ([e10.16]) and ([e10.17]), that

$$\begin{aligned} S_z \chi_{\pm} &= \pm \frac{1}{2} \hbar \chi_{\pm}, \\ S^2 \chi_{\pm} &= \frac{3}{4} \hbar^2 \chi_{\pm}. \end{aligned}$$

Note that  $\chi_+$  corresponds to an electron whose spin angular momentum vector has a positive component along the  $z$ -axis. Loosely speaking, we could say that the spin vector points in the  $+z$ -direction (or its spin is “up”). Likewise,  $\chi_-$  corresponds to an electron whose spin points in the  $-z$ -direction (or whose spin is “down”). These two eigenstates satisfy the orthonormality requirements

$$\chi_+^\dagger \chi_+ = \chi_-^\dagger \chi_- = 1, \quad (9.4.2)$$

and

$$\chi_+^\dagger \chi_- = 0. \quad (9.4.3)$$

A general spin state can be represented as a linear combination of  $\chi_+$  and  $\chi_-$ : that is,

$$\chi = c_+ \chi_+ + c_- \chi_-. \quad (9.4.4)$$

It is thus evident that electron spin space is two-dimensional.

Up to now, we have discussed spin space in rather abstract terms. In the following, we shall describe a particular representation of electron spin space due to Pauli. This so-called *Pauli representation* allows us to visualize spin space, and also facilitates calculations involving spin.

Let us attempt to represent a general spin state as a complex column vector in some two-dimensional space: that is,

$$\chi \equiv \begin{pmatrix} c_+ \\ c_- \end{pmatrix}. \quad (9.4.5)$$

The corresponding dual vector is represented as a row vector: that is,

$$\chi^\dagger \equiv (c_+^*, c_-^*). \quad (9.4.6)$$

Furthermore, the product  $\chi^\dagger \chi$  is obtained according to the ordinary rules of matrix multiplication: that is,

$$\chi^\dagger \chi = (c_+^*, c_-^*) \begin{pmatrix} c_+ \\ c_- \end{pmatrix} = c_+^* c_+ + c_-^* c_- = |c_+|^2 + |c_-|^2 \geq 0. \quad (9.4.7)$$

Likewise, the product  $\chi^\dagger \chi'$  of two different spin states is also obtained from the rules of matrix multiplication: that is,

$$\chi^\dagger \chi' = (c_+^*, c_-^*) \begin{pmatrix} c'_+ \\ c'_- \end{pmatrix} = c_+^* c'_+ + c_-^* c'_-. \quad (9.4.8)$$

Note that this particular representation of spin space is in complete accordance with the discussion in Section 1.3. For obvious reasons, a vector used to represent a spin state is generally known as *spinor*.

A general spin operator  $A$  is represented as a  $2 \times 2$  matrix which operates on a spinor: that is,

$$A \chi \equiv \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \begin{pmatrix} c_+ \\ c_- \end{pmatrix}. \quad (9.4.9)$$

As is easily demonstrated, the Hermitian conjugate of  $A$  is represented by the transposed complex conjugate of the matrix used to represent  $A$ : that is,

$$A^\dagger \equiv \begin{pmatrix} A_{11}^* & A_{21}^* \\ A_{12}^* & A_{22}^* \end{pmatrix}. \quad (9.4.10)$$

Let us represent the spin eigenstates  $\chi_+$  and  $\chi_-$  as

$$\chi_+ \equiv \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad (9.4.11)$$

and

$$\chi_- \equiv \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad (9.4.12)$$

respectively. Note that these forms automatically satisfy the orthonormality constraints ([e10.35]) and ([e10.36]). It is convenient to write the spin operators  $S_i$  (where  $i = 1, 2, 3$  corresponds to  $x, y, z$ ) as

$$S_i = \frac{\hbar}{2} \sigma_i. \quad (9.4.13)$$

Here, the  $\sigma_i$  are dimensionless  $2 \times 2$  matrices. According to Equations ([e10.1x])–([e10.2x]), the  $\sigma_i$  satisfy the commutation relations

$$\begin{aligned} [\sigma_y, \sigma_z] &= 2i\sigma_x, \\ [\sigma_z, \sigma_x] &= 2i\sigma_y, \\ [\sigma_x, \sigma_y] &= 2i\sigma_z. \end{aligned}$$

Furthermore, Equation ([e10.34]) yields

$$\sigma_z \chi_\pm = \pm \chi_\pm. \quad (9.4.14)$$

It is easily demonstrated, from the previous expressions, that the  $\sigma_i$  are represented by the following matrices:

$$\begin{aligned} \sigma_x &\equiv \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \\ \sigma_y &\equiv \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \\ \sigma_z &\equiv \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \end{aligned}$$

Incidentally, these matrices are generally known as the *Pauli matrices*.

Finally, a general spinor takes the form

$$\chi = c_+ \chi_+ + c_- \chi_- = \begin{pmatrix} c_+ \\ c_- \end{pmatrix}. \quad (9.4.15)$$

If the spinor is properly normalized then

$$\chi^\dagger \chi = |c_+|^2 + |c_-|^2 = 1. \quad (9.4.16)$$

In this case, we can interpret  $|c_+|^2$  as the probability that an observation of  $S_z$  will yield the result  $+\hbar/2$ , and  $|c_-|^2$  as the probability that an observation of  $S_z$  will yield the result  $-\hbar/2$ .

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [9.4: Pauli Representation](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 9.5: Spin Precession

According to classical physics, a small current loop possesses a *magnetic moment* of magnitude  $\mu = I A$ , where  $I$  is the current circulating around the loop, and  $A$  the area of the loop. The direction of the magnetic moment is conventionally taken to be normal to the plane of the loop, in the sense given by a standard right-hand circulation rule. Consider a small current loop consisting of an electron in uniform circular motion. It is easily demonstrated that the electron's orbital angular momentum  $\mathbf{L}$  is related to the magnetic moment  $\mu$  of the loop via

$$\mu = -\frac{e}{2m_e} \mathbf{L}, \quad (9.5.1)$$

where  $e$  is the magnitude of the electron charge, and  $m_e$  the electron mass.

The previous expression suggests that there may be a similar relationship between magnetic moment and spin angular momentum. We can write

$$\mu = -\frac{ge}{2m_e} \mathbf{S}, \quad (9.5.2)$$

where  $g$  is called the *gyromagnetic ratio*. Classically, we would expect  $g = 1$ . In fact,

$$g = 2 \left( 1 + \frac{\alpha}{2\pi} + \dots \right) = 2.0023192, \quad (9.5.3)$$

here  $\alpha = e^2 / (2 \epsilon_0 \hbar c) \simeq 1/137$  is the so-called *fine-structure constant*. The fact that the gyromagnetic ratio is (almost) twice that expected from classical physics is only explicable using relativistic quantum mechanics. Furthermore, the small corrections to the relativistic result  $g = 2$  come from quantum field theory.

The energy of a classical magnetic moment  $\mu$  in a uniform magnetic field  $\mathbf{B}$  is

$$H = -\mu \cdot \mathbf{B}. \quad (9.5.4)$$

Assuming that the previous expression also holds good in quantum mechanics, the Hamiltonian of an electron in a  $z$ -directed magnetic field of magnitude  $B$  takes the form

$$H = \Omega S_z, \quad (9.5.5)$$

where

$$\Omega = \frac{geB}{2m_e}. \quad (9.5.6)$$

Here, for the sake of simplicity, we are neglecting the electron's translational degrees of freedom.

Schrödinger's equation can be written [see Equation ([\[etimed\]](#))]

$$i\hbar \frac{\partial \chi}{\partial t} = H \chi, \quad (9.5.7)$$

where the spin state of the electron is characterized by the spinor  $\chi$ . Adopting the Pauli representation, we obtain

$$\chi = \begin{pmatrix} c_+(t) \\ c_-(t) \end{pmatrix}, \quad (9.5.8)$$

where  $|c_+|^2 + |c_-|^2 = 1$ . Here,  $|c_+|^2$  is the probability of observing the spin-up state, and  $|c_-|^2$  the probability of observing the spin-down state. It follows from Equations ([\[e10.46\]](#)), ([\[e10.53\]](#)), ([\[e10.60\]](#)), ([\[e10.62\]](#)), and ([\[e10.63\]](#)) that

$$i\hbar \begin{pmatrix} \dot{c}_+ \\ \dot{c}_- \end{pmatrix} = \frac{\Omega \hbar}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} c_+ \\ c_- \end{pmatrix}, \quad (9.5.9)$$

where  $\dot{\phantom{x}} \equiv d/dt$ . Hence,

$$\dot{c}_{\pm} = \mp i \frac{\Omega}{2} c_{\pm} \quad (9.5.10)$$

Let

$$\begin{aligned}c_+(0) &= \cos(\alpha/2), \\c_-(0) &= \sin(\alpha/2).\end{aligned}$$

The significance of the angle  $\alpha$  will become apparent presently. Solving Equation ([e10.65]), subject to the initial conditions ([e10.66]) and ([e10.67]), we obtain

$$\begin{aligned}c_+(t) &= \cos(\alpha/2) \exp(-i \Omega t/2), \\c_-(t) &= \sin(\alpha/2) \exp(+i \Omega t/2).\end{aligned}$$

We can most easily visualize the effect of the time dependence in the previous expressions for  $c_{\pm}$  by calculating the expectation values of the three Cartesian components of the electron's spin angular momentum. By analogy with Equation ([e3.55]), the expectation value of a general spin operator  $A$  is simply

$$\langle A \rangle = \chi^\dagger A \chi. \quad (9.5.11)$$

Hence, the expectation value of  $S_z$  is

$$\langle S_z \rangle = \frac{\hbar}{2} (c_+^*, c_-^*) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} c_+ \\ c_- \end{pmatrix}, \quad (9.5.12)$$

which reduces to

$$\langle S_z \rangle = \frac{\hbar}{2} \cos \alpha \quad (9.5.13)$$

with the help of Equations ([e10.68]) and ([e10.69]). Likewise, the expectation value of  $S_x$  is

$$\langle S_x \rangle = \frac{\hbar}{2} (c_+^*, c_-^*) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} c_+ \\ c_- \end{pmatrix}, \quad (9.5.14)$$

which reduces to

$$\langle S_x \rangle = \frac{\hbar}{2} \sin \alpha \cos(\Omega t). \quad (9.5.15)$$

Finally, the expectation value of  $S_y$  is

$$\langle S_y \rangle = \frac{\hbar}{2} \sin \alpha \sin(\Omega t). \quad (9.5.16)$$

According to Equations ([e10.72]), ([e10.74]), and ([e10.75]), the expectation value of the spin angular momentum vector subtends a constant angle  $\alpha$  with the  $z$ -axis, and precesses about this axis at the frequency

$$\Omega \simeq \frac{e B}{m_e}. \quad (9.5.17)$$

This behavior is actually equivalent to that predicted by classical physics. Note, however, that a measurement of  $S_x$ ,  $S_y$ , or  $S_z$  will always yield either  $+\hbar/2$  or  $-\hbar/2$ . It is the relative probabilities of obtaining these two results that varies as the expectation value of a given component of the spin varies.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled 9.5: Spin Precession is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 9.E: Spin Angular Momentum (Exercises)

1. Find the Pauli representations of  $S_x$ ,  $S_y$ , and  $S_z$  for a spin-1 particle.
2. Find the Pauli representations of the normalized eigenstates of  $S_x$  and  $S_y$  for a spin-1/2 particle.
3. Suppose that a spin-1/2 particle has a spin vector that lies in the  $x$ - $z$  plane, making an angle  $\theta$  with the  $z$ -axis. Demonstrate that a measurement of  $S_z$  yields  $\hbar/2$  with probability  $\cos^2(\theta/2)$ , and  $-\hbar/2$  with probability  $\sin^2(\theta/2)$ .
4. An electron is in the spin-state

$$\chi = A \begin{pmatrix} 1 - 2i \\ 2 \end{pmatrix} \quad (9.E.1)$$

in the Pauli representation. Determine the constant  $A$  by normalizing  $\chi$ . If a measurement of  $S_z$  is made, what values will be obtained, and with what probabilities? What is the expectation value of  $S_z$ ? Repeat the previous calculations for  $S_x$  and  $S_y$ .

5. Consider a spin-1/2 system represented by the normalized spinor

$$\chi = \begin{pmatrix} \cos \alpha \\ \sin \alpha \exp(i\beta) \end{pmatrix} \quad (9.E.2)$$

in the Pauli representation, where  $\alpha$  and  $\beta$  are real. What is the probability that a measurement of  $S_y$  yields  $-\hbar/2$ ?

6. An electron is at rest in an oscillating magnetic field

$$\mathbf{B} = B_0 \cos(\omega t) \mathbf{e}_z, \quad (9.E.3)$$

where  $B_0$  and  $\omega$  are real positive constants.

1. Find the Hamiltonian of the system.
2. If the electron starts in the spin-up state with respect to the  $x$ -axis, determine the spinor  $\chi(t)$  which represents the state of the system in the Pauli representation at all subsequent times.
3. Find the probability that a measurement of  $S_x$  yields the result  $-\hbar/2$  as a function of time.
4. What is the minimum value of  $B_0$  required to force a complete flip in  $S_x$ ?

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [9.E: Spin Angular Momentum \(Exercises\)](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## CHAPTER OVERVIEW

### 10: Addition of Angular Momentum

Consider an electron in a hydrogen atom. As we have already seen, the electron's motion through space is parameterized by the three quantum numbers  $n$ ,  $l$ , and  $m$ . (See Section [\[s10.4\]](#).) To these we must now add the two quantum numbers  $s$  and  $m_s$  that parameterize the electron's internal motion. (See the previous chapter.) Now, the quantum numbers  $l$  and  $m$  specify the electron's orbital angular momentum vector,  $\mathbf{L}$ , (as much as it can be specified) whereas the quantum numbers  $s$  and  $m_s$  specify its spin angular momentum vector,  $\mathbf{S}$ . But, if the electron possesses both orbital and spin angular momentum then what is its total angular momentum?

[10.1: General Principles of Angular Momentum](#)

[10.2: Angular Momentum in Hydrogen Atom](#)

[10.3: Two Spin One-Half Particles](#)

[10.E: Addition of Angular Momentum \(Exercises\)](#)

---

This page titled [10: Addition of Angular Momentum](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 10.1: General Principles of Angular Momentum

The three fundamental orbital angular momentum operators,  $L_x$ ,  $L_y$ , and  $L_z$ , obey the commutation relations ([e8.6])–([e8.8]), which can be written in the convenient vector form:

$$\mathbf{L} \times \mathbf{L} = i \hbar \mathbf{L}. \quad (10.1.1)$$

Likewise, the three fundamental spin angular momentum operators,  $S_x$ ,  $S_y$ , and  $S_z$ , obey the commutation relations ([e10.1x])–([e10.2x]), which can also be written in vector form: that is,

$$\mathbf{S} \times \mathbf{S} = i \hbar \mathbf{S}. \quad (10.1.2)$$

Because the orbital angular momentum operators are associated with the electron's motion through space, whereas the spin angular momentum operators are associated with its internal motion, and these two types of motion are completely unrelated (i.e., they correspond to different degrees of freedom—see Section [sfuncon]), it is reasonable to suppose that the two sets of operators commute with one another: that is,

$$[L_i, S_j] = 0, \quad (10.1.3)$$

where  $i, j = 1, 2, 3$  corresponds to  $x, y, z$ , respectively.

Let us now consider the electron's total angular momentum vector

$$\mathbf{J} = \mathbf{L} + \mathbf{S}. \quad (10.1.4)$$

We have

$$\begin{aligned} \mathbf{J} \times \mathbf{J} &= (\mathbf{L} + \mathbf{S}) \times (\mathbf{L} + \mathbf{S}) \\ &= \mathbf{L} \times \mathbf{L} + \mathbf{S} \times \mathbf{S} + \mathbf{L} \times \mathbf{S} + \mathbf{S} \times \mathbf{L} = \mathbf{L} \times \mathbf{L} + \mathbf{S} \times \mathbf{S} \\ &= i \hbar \mathbf{L} + i \hbar \mathbf{S} = i \hbar \mathbf{J}. \end{aligned}$$

In other words,

$$\mathbf{J} \times \mathbf{J} = i \hbar \mathbf{J}. \quad (10.1.5)$$

It is thus evident that the three fundamental total angular momentum operators,  $J_x$ ,  $J_y$ , and  $J_z$ , obey analogous commutation relations to the corresponding orbital and spin angular momentum operators. It therefore follows that the total angular momentum has similar properties to the orbital and spin angular momenta. For instance, it is only possible to simultaneously measure the magnitude squared of the total angular momentum vector,

$$J^2 = J_x^2 + J_y^2 + J_z^2, \quad (10.1.6)$$

together with a single Cartesian component. By convention, we shall always choose to measure  $J_z$ . A simultaneous eigenstate of  $J_z$  and  $J^2$  satisfies

$$\begin{aligned} J_z \psi_{j,m_j} &= m_j \hbar \psi_{j,m_j}, \\ J^2 \psi_{j,m_j} &= j(j+1) \hbar^2 \psi_{j,m_j}, \end{aligned}$$

where the quantum number  $j$  can take positive integer, or half-integer, values, and the quantum number  $m_j$  is restricted to the following range of values:

$$-j, -j+1, \dots, j-1, j. \quad (10.1.7)$$

Now,

$$J^2 = (\mathbf{L} + \mathbf{S}) \cdot (\mathbf{L} + \mathbf{S}) = L^2 + S^2 + 2 \mathbf{L} \cdot \mathbf{S}, \quad (10.1.8)$$

which can also be written as

$$J^2 = L^2 + S^2 + 2 L_z S_z + L_+ S_- + L_- S_+. \quad (10.1.9)$$

We know that the operator  $L^2$  commutes with itself, with all of the Cartesian components of  $\mathbf{L}$  (and, hence, with the raising and lowering operators  $L_{\pm}$ ), and with all of the spin angular momentum operators. (See Section [s8.2].) It is therefore clear that

$$[J^2, L^2] = 0. \quad (10.1.10)$$

A similar argument allows us to also conclude that

$$[J^2, S^2] = 0. \quad (10.1.11)$$

Now, the operator  $L_z$  commutes with itself, with  $L^2$ , with all of the spin angular momentum operators, but not with the raising and lowering operators  $L_{\pm}$ . (See Section [s8.2].) It follows that

$$[J^2, L_z] \neq 0. \quad (10.1.12)$$

Likewise, we can also show that

$$[J^2, S_z] \neq 0. \quad (10.1.13)$$

Finally, we have

$$J_z = L_z + S_z, \quad (10.1.14)$$

where  $[J_z, L_z] = [J_z, S_z] = 0$ .

Recalling that only commuting operators correspond to physical quantities that can be simultaneously measured (see Section [smeas]), it follows, from the previous discussion, that there are two alternative sets of physical variables associated with angular momentum that we can measure simultaneously. The first set correspond to the operators  $L^2$ ,  $S^2$ ,  $L_z$ ,  $S_z$ , and  $J_z$ . The second set correspond to the operators  $L^2$ ,  $S^2$ ,  $J^2$ , and  $J_z$ . In other words, we can always measure the magnitude squared of the orbital and spin angular momentum vectors, together with the  $z$ -component of the total angular momentum vector. In addition, we can either choose to measure the  $z$ -components of the orbital and spin angular momentum vectors, or the magnitude squared of the total angular momentum vector.

Let  $\psi_{l,s;m,m_s}^{(1)}$  represent a simultaneous eigenstate of  $L^2$ ,  $S^2$ ,  $L_z$ , and  $S_z$  corresponding to the following eigenvalues:

$$\begin{aligned} L^2 \psi_{l,s;m,m_s}^{(1)} &= l(l+1) \hbar^2 \psi_{l,s;m,m_s}^{(1)}, \\ S^2 \psi_{l,s;m,m_s}^{(1)} &= s(s+1) \hbar^2 \psi_{l,s;m,m_s}^{(1)}, \\ L_z \psi_{l,s;m,m_s}^{(1)} &= m \hbar \psi_{l,s;m,m_s}^{(1)}, \\ S_z \psi_{l,s;m,m_s}^{(1)} &= m_s \hbar \psi_{l,s;m,m_s}^{(1)}. \end{aligned}$$

It is easily seen that

$$\begin{aligned} J_z \psi_{l,s;m,m_s}^{(1)} &= (L_z + S_z) \psi_{l,s;m,m_s}^{(1)} = (m + m_s) \hbar \psi_{l,s;m,m_s}^{(1)} \\ &= m_j \hbar \psi_{l,s;m,m_s}^{(1)}. \end{aligned}$$

Hence,

$$m_j = m + m_s. \quad (10.1.15)$$

In other words, the quantum numbers controlling the  $z$ -components of the various angular momentum vectors can simply be added algebraically.

Finally, let  $\psi_{l,s;j,m_j}^{(2)}$  represent a simultaneous eigenstate of  $L^2$ ,  $S^2$ ,  $J^2$ , and  $J_z$  corresponding to the following eigenvalues:

$$\begin{aligned} L^2 \psi_{l,s;j,m_j}^{(2)} &= l(l+1) \hbar^2 \psi_{l,s;j,m_j}^{(2)}, \\ S^2 \psi_{l,s;j,m_j}^{(2)} &= s(s+1) \hbar^2 \psi_{l,s;j,m_j}^{(2)}, \\ J^2 \psi_{l,s;j,m_j}^{(2)} &= j(j+1) \hbar^2 \psi_{l,s;j,m_j}^{(2)}, \\ J_z \psi_{l,s;j,m_j}^{(2)} &= m_j \hbar \psi_{l,s;j,m_j}^{(2)}. \end{aligned}$$



## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [10.1: General Principles of Angular Momentum](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 10.2: Angular Momentum in Hydrogen Atom

In a hydrogen atom, the wavefunction of an electron in a simultaneous eigenstate of  $L^2$  and  $L_z$  has an angular dependence specified by the spherical harmonic  $Y_{l,m}(\theta, \phi)$ . (See Section [sharm].) If the electron is also in an eigenstate of  $S^2$  and  $S_z$  then the quantum numbers  $s$  and  $m_s$  take the values  $1/2$  and  $\pm 1/2$ , respectively, and the internal state of the electron is specified by the spinors  $\chi_{\pm}$ . (See Section [spauli].) Hence, the simultaneous eigenstates of  $L^2$ ,  $S^2$ ,  $L_z$ , and  $S_z$  can be written in the separable form

$$\psi_{l,1/2;m,\pm 1/2}^{(1)} = Y_{l,m} \chi_{\pm}. \quad (10.2.1)$$

Here, it is understood that orbital angular momentum operators act on the spherical harmonic functions,  $Y_{l,m}$ , whereas spin angular momentum operators act on the spinors,  $\chi_{\pm}$ .

Because the eigenstates  $\psi_{l,1/2;m,\pm 1/2}^{(1)}$  are (presumably) orthonormal, and form a complete set, we can express the eigenstates  $\psi_{l,1/2;j,m_j}^{(2)}$  as linear combinations of them. For instance,

$$\psi_{l,1/2;j,m+1/2}^{(2)} = \alpha \psi_{l,1/2;m,1/2}^{(1)} + \beta \psi_{l,1/2;m+1,-1/2}^{(1)} \quad (10.2.2)$$

where  $\alpha$  and  $\beta$  are, as yet, unknown coefficients. Note that the number of  $\psi^{(1)}$  states that can appear on the right-hand side of the previous expression is limited to two by the constraint that  $m_j = m + m_s$  [see Equation ([e11.23])], and the fact that  $m_s$  can only take the values  $\pm 1/2$ . Assuming that the  $\psi^{(2)}$  eigenstates are properly normalized, we have

$$\alpha^2 + \beta^2 = 1. \quad (10.2.3)$$

Now, it follows from Equation ([e11.26]) that

$$J^2 \psi_{l,1/2;j,m+1/2}^{(2)} = j(j+1) \hbar^2 \psi_{l,1/2;j,m+1/2}^{(2)} \quad (10.2.4)$$

where [see Equation ([e11.12])]

$$J^2 = L^2 + S^2 + 2L_z S_z + L_+ S_- + L_- S_+. \quad (10.2.5)$$

Moreover, according to Equations ([e11.28]) and ([e11.29]), we can write

$$\psi_{l,1/2;j,m+1/2}^{(2)} = \alpha Y_{l,m} \chi_+ + \beta Y_{l,m+1} \chi_-. \quad (10.2.6)$$

Recall, from Equations ([e11.28]) and ([e11.29]), that

$$\begin{aligned} L_+ Y_{l,m} &= [l(l+1) - m(m+1)]^{1/2} \hbar Y_{l,m+1}, \\ L_- Y_{l,m} &= [l(l+1) - m(m-1)]^{1/2} \hbar Y_{l,m-1}. \end{aligned}$$

By analogy, when the spin raising and lowering operators,  $S_{\pm}$ , act on a general spinor,  $\chi_{s,m_s}$ , we obtain

$$\begin{aligned} S_+ \chi_{s,m_s} &= [s(s+1) - m_s(m_s+1)]^{1/2} \hbar \chi_{s,m_s+1}, \\ S_- \chi_{s,m_s} &= [s(s+1) - m_s(m_s-1)]^{1/2} \hbar \chi_{s,m_s-1}. \end{aligned}$$

For the special case of spin one-half spinors (i.e.,  $s = 1/2$ ,  $m_s = \pm 1/2$ ), the previous expressions reduce to

$$S_+ \chi_+ = S_- \chi_- = 0, \quad (10.2.7)$$

and

$$S_{\pm} \chi_{\mp} = \hbar \chi_{\pm}. \quad (10.2.8)$$

It follows from Equations ([e11.32]) and ([e11.34])–([e11.39]) that

$$\begin{aligned} J^2 Y_{l,m} \chi_+ &= [l(l+1) + 3/4 + m] \hbar^2 Y_{l,m} \chi_+ \\ &\quad + [l(l+1) - m(m+1)]^{1/2} \hbar^2 Y_{l,m+1} \chi_-, \end{aligned}$$

and

$$J^2 Y_{l,m+1} \chi_- = [l(l+1) + 3/4 - m - 1] \hbar^2 Y_{l,m+1} \chi_- \\ + [l(l+1) - m(m+1)]^{1/2} \hbar^2 Y_{l,m} \chi_+.$$

Hence, Equations ([e11.31]) and ([e11.33]) yield

$$(x - m) \alpha - [l(l+1) - m(m+1)]^{1/2} \beta = 0, \\ -[l(l+1) - m(m+1)]^{1/2} \alpha + (x + m + 1) \beta = 0,$$

where

$$x = j(j+1) - l(l+1) - 3/4. \quad (10.2.9)$$

Equations ([e11.42]) and ([e11.43]) can be solved to give

$$x(x+1) = l(l+1), \quad (10.2.10)$$

and

$$\frac{\alpha}{\beta} = \frac{[(l-m)(l+m+1)]^{1/2}}{x-m}. \quad (10.2.11)$$

It follows that  $x = l$  or  $x = -l - 1$ , which corresponds to  $j = l + 1/2$  or  $j = l - 1/2$ , respectively. Once  $x$  is specified, Equations ([e11.30]) and ([e11.45]) can be solved for  $\alpha$  and  $\beta$ . We obtain

$$\psi_{l+1/2, m+1/2}^{(2)} = \left( \frac{l+m+1}{2l+1} \right)^{1/2} \psi_{m, 1/2}^{(1)} + \left( \frac{l-m}{2l+1} \right)^{1/2} \psi_{m+1, -1/2}^{(1)}, \quad (10.2.12)$$

and

$$\psi_{l-1/2, m+1/2}^{(2)} = \left( \frac{l-m}{2l+1} \right)^{1/2} \psi_{m, 1/2}^{(1)} - \left( \frac{l+m+1}{2l+1} \right)^{1/2} \psi_{m+1, -1/2}^{(1)}. \quad (10.2.13)$$

Here, we have neglected the common subscripts  $l, 1/2$  for the sake of clarity: that is,  $\psi_{l+1/2, m+1/2}^{(2)} \equiv \psi_{l, 1/2; l+1/2, m+1/2}^{(2)}$ , et cetera. The previous equations can easily be inverted to give the  $\psi^{(1)}$  eigenstates in terms of the  $\psi^{(2)}$  eigenstates:

$$\psi_{m, 1/2}^{(1)} = \left( \frac{l+m+1}{2l+1} \right)^{1/2} \psi_{l+1/2, m+1/2}^{(2)} + \left( \frac{l-m}{2l+1} \right)^{1/2} \psi_{l-1/2, m+1/2}^{(2)}, \\ \psi_{m+1, -1/2}^{(1)} = \left( \frac{l-m}{2l+1} \right)^{1/2} \psi_{l+1/2, m+1/2}^{(2)} - \left( \frac{l+m+1}{2l+1} \right)^{1/2} \psi_{l-1/2, m+1/2}^{(2)}.$$

The information contained in Equations ([e11.47])–([e11.50]) is neatly summarized in Table [t2]. For instance, Equation ([e11.47]) is obtained by reading the first row of this table, whereas Equation ([e11.50]) is obtained by reading the second column. The coefficients in this type of table are generally known as *Clebsch-Gordon coefficients*.

Clebsch-Gordon coefficients for adding spin one-half to spin  $l$ .

	$m, 1/2$	$m+1, -1/2$	$m, m_s$
[0.5ex] $l+1/2, m+1/2$	$\sqrt{(l+m+1)/(2l+1)}$	$\sqrt{(l-m)/(2l+1)}$	
[0.5ex] $l-1/2, m+1/2$	$\sqrt{(l-m)/(2l+1)}$	$-\sqrt{(l+m+1)/(2l+1)}$	
[0.5ex] $j, m_j$			

As an example, let us consider the  $l = 1$  states of a hydrogen atom. The eigenstates of  $L^2$ ,  $S^2$ ,  $L_z$ , and  $S_z$ , are denoted  $\psi_{m, m_s}^{(1)}$ . Because  $m$  can take the values  $-1, 0, 1$ , whereas  $m_s$  can take the values  $\pm 1/2$ , there are clearly six such states: that is,  $\psi_{1, \pm 1/2}^{(1)}$ ,  $\psi_{0, \pm 1/2}^{(1)}$ , and  $\psi_{-1, \pm 1/2}^{(1)}$ . The eigenstates of  $L^2$ ,  $S^2$ ,  $J^2$ , and  $J_z$ , are denoted  $\psi_{j, m_j}^{(2)}$ . Because  $l = 1$  and  $s = 1/2$  can be combined together to form either  $j = 3/2$  or  $j = 1/2$  (see previously), there are also six such states: that is,  $\psi_{3/2, \pm 3/2}^{(2)}$ ,  $\psi_{3/2, \pm 1/2}^{(2)}$ , and  $\psi_{1/2, \pm 1/2}^{(2)}$ . According to Table [t2], the various different eigenstates are interrelated as follows:

$$\begin{aligned}\psi_{3/2,\pm 3/2}^{(2)} &= \psi_{\pm 1,\pm 1/2}^{(1)}, \\ \psi_{3/2,1/2}^{(2)} &= \sqrt{\frac{2}{3}} \psi_{0,1/2}^{(1)} + \sqrt{\frac{1}{3}} \psi_{1,-1/2}^{(1)}, \\ \psi_{1/2,1/2}^{(2)} &= \sqrt{\frac{1}{3}} \psi_{0,1/2}^{(1)} - \sqrt{\frac{2}{3}} \psi_{1,-1/2}^{(1)}, \\ \psi_{1/2,-1/2}^{(2)} &= \sqrt{\frac{2}{3}} \psi_{-1,1/2}^{(1)} - \sqrt{\frac{1}{3}} \psi_{0,-1/2}^{(1)}, \\ \psi_{3/2,-1/2}^{(2)} &= \sqrt{\frac{1}{3}} \psi_{-1,1/2}^{(1)} + \sqrt{\frac{2}{3}} \psi_{0,-1/2}^{(1)},\end{aligned}$$

and

$$\begin{aligned}\psi_{\pm 1,\pm 1/2}^{(1)} &= \psi_{3/2,\pm 3/2}^{(2)}, \\ \psi_{1,-1/2}^{(1)} &= \sqrt{\frac{1}{3}} \psi_{3/2,1/2}^{(2)} - \sqrt{\frac{2}{3}} \psi_{1/2,1/2}^{(2)}, \\ \psi_{0,1/2}^{(1)} &= \sqrt{\frac{2}{3}} \psi_{3/2,1/2}^{(2)} + \sqrt{\frac{1}{3}} \psi_{1/2,1/2}^{(2)}, \\ \psi_{0,-1/2}^{(1)} &= \sqrt{\frac{2}{3}} \psi_{3/2,-1/2}^{(2)} - \sqrt{\frac{1}{3}} \psi_{1/2,-1/2}^{(2)}, \\ \psi_{-1,1/2}^{(1)} &= \sqrt{\frac{1}{3}} \psi_{3/2,-1/2}^{(2)} + \sqrt{\frac{2}{3}} \psi_{1/2,-1/2}^{(2)}.\end{aligned}$$

Thus, if we know that an electron in a hydrogen atom is in an  $l = 1$  state characterized by  $m = 0$  and  $m_s = 1/2$  [i.e., the state represented by  $\psi_{0,1/2}^{(1)}$ ] then, according to Equation ([e11.57]), a measurement of the total angular momentum will yield  $j = 3/2$ ,  $m_j = 1/2$  with probability  $2/3$ , and  $j = 1/2$ ,  $m_j = 1/2$  with probability  $1/3$ . Suppose that we make such a measurement, and obtain the result  $j = 3/2$ ,  $m_j = 1/2$ . As a result of the measurement, the electron is thrown into the corresponding eigenstate,  $\psi_{3/2,1/2}^{(2)}$ . It thus follows from Equation ([e11.52]) that a subsequent measurement of  $L_z$  and  $S_z$  will yield  $m = 0$ ,  $m_s = 1/2$  with probability  $2/3$ , and  $m = 1$ ,  $m_s = -1/2$  with probability  $1/3$ .

Clebsch-Gordon coefficients for adding spin one-half to spin one. Only non-zero coefficients are shown.

	$-1, -1/2$	$-1, 1/2$	$0, -1/2$	$0, 1/2$	$1, -1/2$	$1, 1/2$	$m, m_s$
[0.5ex] $3/2, -3/2$	1						
[0.5ex] $3/2, -1/2$		$\sqrt{1/3}$	$\sqrt{2/3}$				
[0.5ex] $1/2, -1/2$		$\sqrt{2/3}$	$-\sqrt{1/3}$				
[0.5ex] $3/2, 1/2$				$\sqrt{2/3}$	$\sqrt{1/3}$		
[0.5ex] $1/2, 1/2$				$\sqrt{1/3}$	$-\sqrt{2/3}$		
[0.5ex] $3/2, 3/2$						1	
$j, m_j$							

The information contained in Equations ([ecgs])–([ecge]) is neatly summed up in Table [t3]. Note that each row and column of this table has unit norm, and also that the different rows and different columns are mutually orthogonal. Of course, this is because the

$\psi^{(1)}$  and  $\psi^{(2)}$  eigenstates are orthonormal.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [10.2: Angular Momentum in Hydrogen Atom](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 10.3: Two Spin One-Half Particles

Consider a system consisting of two spin one-half particles. Suppose that the system does not possess any orbital angular momentum. Let  $\mathbf{S}_1$  and  $\mathbf{S}_2$  be the spin angular momentum operators of the first and second particles, respectively, and let

$$\mathbf{S} = \mathbf{S}_1 + \mathbf{S}_2 \quad (10.3.1)$$

be the total spin angular momentum operator. By analogy with the previous analysis, we conclude that it is possible to simultaneously measure either  $S_1^2$ ,  $S_2^2$ ,  $S^2$ , and  $S_z$ , or  $S_1^2$ ,  $S_2^2$ ,  $S_{1z}$ ,  $S_{2z}$ , and  $S_z$ . Let the quantum numbers associated with measurements of  $S_1^2$ ,  $S_{1z}$ ,  $S_2^2$ ,  $S_{2z}$ ,  $S^2$ , and  $S_z$  be  $s_1$ ,  $m_{s_1}$ ,  $s_2$ ,  $m_{s_2}$ ,  $s$ , and  $m_s$ , respectively. In other words, if the spinor  $\chi_{s_1, s_2; m_{s_1}, m_{s_2}}^{(1)}$  is a simultaneous eigenstate of  $S_1^2$ ,  $S_2^2$ ,  $S_{1z}$ , and  $S_{2z}$ , then

$$\begin{aligned} S_1^2 \chi_{s_1, s_2; m_{s_1}, m_{s_2}}^{(1)} &= s_1(s_1 + 1) \hbar^2 \chi_{s_1, s_2; m_{s_1}, m_{s_2}}^{(1)}, \\ S_2^2 \chi_{s_1, s_2; m_{s_1}, m_{s_2}}^{(1)} &= s_2(s_2 + 1) \hbar^2 \chi_{s_1, s_2; m_{s_1}, m_{s_2}}^{(1)}, \\ S_{1z} \chi_{s_1, s_2; m_{s_1}, m_{s_2}}^{(1)} &= m_{s_1} \hbar \chi_{s_1, s_2; m_{s_1}, m_{s_2}}^{(1)}, \\ S_{2z} \chi_{s_1, s_2; m_{s_1}, m_{s_2}}^{(1)} &= m_{s_2} \hbar \chi_{s_1, s_2; m_{s_1}, m_{s_2}}^{(1)}, \\ S_z \chi_{s_1, s_2; m_{s_1}, m_{s_2}}^{(1)} &= m_s \hbar \chi_{s_1, s_2; m_{s_1}, m_{s_2}}^{(1)}. \end{aligned}$$

Likewise, if the spinor  $\chi_{s_1, s_2; s, m_s}^{(2)}$  is a simultaneous eigenstate of  $S_1^2$ ,  $S_2^2$ ,  $S^2$ , and  $S_z$ , then

$$\begin{aligned} S_1^2 \chi_{s_1, s_2; s, m_s}^{(2)} &= s_1(s_1 + 1) \hbar^2 \chi_{s_1, s_2; s, m_s}^{(2)}, \\ S_2^2 \chi_{s_1, s_2; s, m_s}^{(2)} &= s_2(s_2 + 1) \hbar^2 \chi_{s_1, s_2; s, m_s}^{(2)}, \\ S^2 \chi_{s_1, s_2; s, m_s}^{(2)} &= s(s + 1) \hbar^2 \chi_{s_1, s_2; s, m_s}^{(2)}, \\ S_z \chi_{s_1, s_2; s, m_s}^{(2)} &= m_s \hbar \chi_{s_1, s_2; s, m_s}^{(2)}. \end{aligned}$$

Of course, because both particles have spin one-half,  $s_1 = s_2 = 1/2$ , and  $s_{1z}, s_{2z} = \pm 1/2$ . Furthermore, by analogy with previous analysis,

$$m_s = m_{s_1} + m_{s_2}. \quad (10.3.2)$$

Now, we saw, in the previous section, that when spin  $l$  is added to spin one-half then the possible values of the total angular momentum quantum number are  $j = l \pm 1/2$ . By analogy, when spin one-half is added to spin one-half then the possible values of the total spin quantum number are  $s = 1/2 \pm 1/2$ . In other words, when two spin one-half particles are combined, we either obtain a state with overall spin  $s = 1$ , or a state with overall spin  $s = 0$ . To be more exact, there are three possible  $s = 1$  states (corresponding to  $m_s = -1, 0, 1$ ), and one possible  $s = 0$  state (corresponding to  $m_s = 0$ ). The three  $s = 1$  states are generally known as the *triplet* states, whereas the  $s = 0$  state is known as the *singlet* state.

Clebsch-Gordon coefficients for adding spin one-half to spin one-half. Only non-zero coefficients are shown.

	$-1/2, -1/2$	$-1/2, 1/2$	$1/2, -1/2$	$1/2, 1/2$	$m_{s_1}, m_{s_2}$
[0.5ex] 1, -1	1				
[0.5ex] 1, 0		$1/\sqrt{2}$	$1/\sqrt{2}$		
[0.5ex] 0, 0		$1/\sqrt{2}$	$-1/\sqrt{2}$		
[0.5ex] 1, 1				1	
$s, m_s$					

The Clebsch-Gordon coefficients for adding spin one-half to spin one-half can easily be inferred from Table [t2] (with  $l = 1/2$ ), and are listed in Table [t4]. It follows from this table that the three triplet states are:

$$\begin{aligned}\chi_{1,-1}^{(2)} &= \chi_{-1/2,-1.2}^{(1)}, \\ \chi_{1,0}^{(2)} &= \frac{1}{\sqrt{2}} \left( \chi_{-1/2,1/2}^{(1)} + \chi_{1/2,-1/2}^{(1)} \right), \\ \chi_{1,1}^{(2)} &= \chi_{1/2,1/2}^{(1)},\end{aligned}$$

where  $\chi_{s,m_s}^{(2)}$  is shorthand for  $\chi_{s_1,s_2;s,m_s}^{(2)}$ , et cetera. Likewise, the singlet state is written:

$$\chi_{0,0}^{(2)} = \frac{1}{\sqrt{2}} \left( \chi_{-1/2,1/2}^{(1)} - \chi_{1/2,-1/2}^{(1)} \right). \quad (10.3.3)$$

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [10.3: Two Spin One-Half Particles](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 10.E: Addition of Angular Momentum (Exercises)

1. An electron in a hydrogen atom occupies the combined spin and position state

$$R_{2,1}(r) \left[ \sqrt{1/3} Y_{1,0}(\theta, \phi) \chi_+ + \sqrt{2/3} Y_{1,1}(\theta, \phi) \chi_- \right]. \quad (10.E.1)$$

1. What values would a measurement of  $L^2$  yield, and with what probabilities?
  2. Same for  $L_z$ .
  3. Same for  $S^2$ .
  4. Same for  $S_z$ .
  5. Same for  $J^2$ .
  6. Same for  $J_z$ .
  7. What is the probability density for finding the electron at  $r, \theta, \phi$ ?
  8. What is the probability density for finding the electron in the spin up state (with respect to the  $z$ -axis) at radius  $r$ ?
2. In a low energy neutron-proton system (with zero orbital angular momentum), the potential energy is given by

$$V(r) = V_1(r) + V_2(r) \left[ 3 \frac{(\sigma_1 \cdot \mathbf{r})(\sigma_2 \cdot \mathbf{r})}{r^2} - \sigma_1 \cdot \sigma_2 \right] + V_3(r) \sigma_1 \cdot \sigma_2, \quad (10.E.2)$$

where  $\sigma_1$  denotes the vector of the Pauli matrices of the neutron, and  $\sigma_2$  denotes the vector of the Pauli matrices of the proton. Calculate the potential energy for the neutron-proton system:

1. In the spin singlet state.
  2. In the spin triplet state.
3. Consider two electrons in a spin singlet state.
    1. If a measurement of the spin of one of the electrons shows that it is in the state with  $S_z = \hbar/2$ , what is the probability that a measurement of the  $z$ -component of the spin of the other electron yields  $S_z = \hbar/2$ ?
    2. If a measurement of the spin of one of the electrons shows that it is in the state with  $S_y = \hbar/2$ , what is the probability that a measurement of the  $x$ -component of the spin of the other electron yields  $S_x = -\hbar/2$ ?

Finally, if electron 1 is in a spin state described by  $\cos \alpha_1 \chi_+ + \sin \alpha_1 e^{i\beta_1} \chi_-$ , and electron 2 is in a spin state described by  $\cos \alpha_2 \chi_+ + \sin \alpha_2 e^{i\beta_2} \chi_-$ , what is the probability that the two-electron spin state is a triplet state?

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [10.E: Addition of Angular Momentum \(Exercises\)](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).



## CHAPTER OVERVIEW

### 11: Time-Independent Perturbation Theory

Consider the following very commonly occurring problem. The Hamiltonian of a quantum mechanical system is written

$$H = H_0 + H_1. \quad (11.1)$$

Here,  $H_0$  is a simple Hamiltonian whose eigenvalues and eigenstates are known exactly.  $H_1$  introduces some interesting additional physics into the problem, but is sufficiently complicated that when we add it to  $H_0$  we can no longer find the exact energy eigenvalues and eigenstates. However,  $H_1$  can, in some sense (which we shall specify more precisely later on), be regarded as small compared to  $H_0$ . Can we find approximate eigenvalues and eigenstates of the modified Hamiltonian,  $H_0 + H_1$ , by performing some sort of perturbation expansion about the eigenvalues and eigenstates of the original Hamiltonian,  $H_0$ ? Let us investigate.

Incidentally, in this chapter, we shall only discuss so-called *time-independent perturbation theory*, in which the modification to the Hamiltonian,  $H_1$ , has no explicit dependence on time. It is also assumed that the unperturbed Hamiltonian,  $H_0$ , is time independent.

[11.1: Exercises](#)

[11.2: Improved Notation](#)

[11.3: Two-State System](#)

[11.4: Non-Degenerate Perturbation Theory](#)

[11.5: Quadratic Stark Effect](#)

[11.6: Degenerate Perturbation Theory](#)

[11.7: Linear Stark Effect](#)

[11.8: Fine Structure of Hydrogen](#)

[11.9: Zeeman Effect](#)

[11.10: Hyperfine Structure](#)

---

This page titled [11: Time-Independent Perturbation Theory](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 11.1: Exercises

1. Consider the two-state system investigated in Section 1.3. Show that the most general expressions for the perturbed energy eigenvalues and eigenstates are

$$E'_1 = E_1 + e_{11} + \frac{|e_{12}|^2}{E_1 - E_2} + \mathcal{O}(\epsilon^3),$$

$$E'_2 = E_2 + e_{22} - \frac{|e_{12}|^2}{E_1 - E_2} + \mathcal{O}(\epsilon^3),$$

and

$$\psi'_1 = \psi_1 + \frac{e_{12}^*}{E_1 - E_2} \psi_2 + \mathcal{O}(\epsilon^2),$$

$$\psi'_2 = \psi_2 - \frac{e_{12}}{E_1 - E_2} \psi_1 + \mathcal{O}(\epsilon^2),$$

respectively. Here,  $\epsilon = |e_{12}|/(E_1 - E_2) \ll 1$ . You may assume that  $|e_{11}|/(E_1 - E_2), |e_{22}|/(E_1 - E_2) \sim \mathcal{O}(\epsilon)$ .

2. Consider the two-state system investigated in Section 1.3. Show that if the unperturbed energy eigenstates are also eigenstates of the perturbing Hamiltonian then

$$E'_1 = E_1 + e_{11},$$

$$E'_2 = E_2 + e_{22},$$

and

$$\psi'_1 = \psi_1$$

$$\psi'_2 = \psi_2$$

to all orders in the perturbation expansion.

3. Consider the two-state system investigated in Section 1.3. Show that if the unperturbed energy eigenstates are degenerate, so that  $E_1 = E_2 = E_{12}$ , then the most general expressions for the perturbed energy eigenvalues and eigenstates are

$$E^\pm = E_{12} + e^\pm, \quad (11.1.1)$$

and

$$\psi^\pm = \langle 1 | \psi^\pm \rangle \psi_1 + \langle 2 | \psi^\pm \rangle \psi_2, \quad (11.1.2)$$

respectively, where

$$e^\pm = \frac{1}{2} (e_{11} + e_{22}) \pm \frac{1}{2} \left[ (e_{11} - e_{22})^2 + 4 |e_{12}|^2 \right]^{1/2}, \quad (11.1.3)$$

and

$$\frac{\langle 1 | \psi^\pm \rangle}{\langle 2 | \psi^\pm \rangle} = - \left( \frac{e_{12}}{e_{11} - e^\pm} \right) = - \left( \frac{e_{22} - e^\pm}{e_{12}^*} \right). \quad (11.1.4)$$

Demonstrate that the  $\psi^\pm$  are the simultaneous eigenstates of the unperturbed Hamiltonian,  $H_0$ , and the perturbed Hamiltonian,  $H_1$ , and that the  $e^\pm$  are the corresponding eigenvalues of  $H_1$ .

4. Calculate the lowest-order energy-shift in the ground state of the one-dimensional harmonic oscillator when the perturbation

$$V = \lambda x^4 \quad (11.1.5)$$

is added to

$$H = \frac{p_x^2}{2m} + \frac{1}{2} m \omega^2 x^2. \quad (11.1.6)$$

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [11.1: Exercises](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 11.2: Improved Notation

Before commencing our investigation, it is helpful to introduce some improved notation. Let the  $\psi_i$  be a complete set of eigenstates of the Hamiltonian,  $H$ , corresponding to the eigenvalues  $E_i$ : that is,

$$H \psi_i = E_i \psi_i. \quad (11.2.1)$$

Now, we expect the  $\psi_i$  to be orthonormal. (See Section [\[seig\]](#).) In one dimension, this implies that

$$\int_{-\infty}^{\infty} \psi_i^* \psi_j dx = \delta_{ij}. \quad (11.2.2)$$

In three dimensions (see Chapter [\[sthree\]](#)), the previous expression generalizes to

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \psi_i^* \psi_j dx dy dz = \delta_{ij}. \quad (11.2.3)$$

Finally, if the  $\psi_i$  are spinors (see Chapter [\[sspin\]](#)) then we have

$$\psi_i^\dagger \psi_j = \delta_{ij}. \quad (11.2.4)$$

The generalization to the case where  $\psi$  is a product of a regular wavefunction and a spinor is fairly obvious. We can represent all of the previous possibilities by writing

$$\langle \psi_i | \psi_j \rangle \equiv \langle i | j \rangle = \delta_{ij}. \quad (11.2.5)$$

Here, the term in angle brackets represents the integrals appearing in Equations ([\[e12.1\]](#)) and ([\[e12.2\]](#)) in one- and three-dimensional regular space, respectively, and the spinor product appearing in Equation ([\[e12.3\]](#)) in spin-space. The advantage of our new notation is its great generality: that is, it can deal with one-dimensional wavefunctions, three-dimensional wavefunctions, spinors, et cetera.

Expanding a general wavefunction,  $\psi_a$ , in terms of the energy eigenstates,  $\psi_i$ , we obtain

$$\psi_a = \sum_i c_i \psi_i. \quad (11.2.6)$$

In one dimension, the expansion coefficients take the form (see Section [\[seig\]](#))

$$c_i = \int_{-\infty}^{\infty} \psi_i^* \psi_a dx, \quad (11.2.7)$$

whereas in three dimensions we get

$$c_i = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \psi_i^* \psi_a dx dy dz. \quad (11.2.8)$$

Finally, if  $\psi$  is a spinor then we have

$$c_i = \psi_i^\dagger \psi_a. \quad (11.2.9)$$

We can represent all of the previous possibilities by writing

$$c_i = \langle \psi_i | \psi_a \rangle \equiv \langle i | a \rangle. \quad (11.2.10)$$

The expansion ([\[e12.7\]](#)) thus becomes

$$\psi_a = \sum_i \langle \psi_i | \psi_a \rangle \psi_i \equiv \sum_i \langle i | a \rangle \psi_i. \quad (11.2.11)$$

Incidentally, it follows that

$$\langle i | a \rangle^* = \langle a | i \rangle. \quad (11.2.12)$$

Finally, if  $A$  is a general operator, and the wavefunction  $\psi_a$  is expanded in the manner shown in Equation ([\[e12.7\]](#)), then the expectation value of  $A$  is written (see Section [\[seig\]](#))

$$\langle A \rangle = \sum_{i,j} c_i^* c_j A_{ij}. \quad (11.2.13)$$

Here, the  $A_{ij}$  are unsurprisingly known as the *matrix elements* of  $A$ . In one dimension, the matrix elements take the form

$$A_{ij} = \int_{-\infty}^{\infty} \psi_i^* A \psi_j dx, \quad (11.2.14)$$

whereas in three dimensions we get

$$A_{ij} = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \psi_i^* A \psi_j dx dy dz. \quad (11.2.15)$$

Finally, if  $\psi$  is a spinor then we have

$$A_{ij} = \psi_i^\dagger A \psi_j. \quad (11.2.16)$$

We can represent all of the previous possibilities by writing

$$A_{ij} = \langle \psi_i | A | \psi_j \rangle \equiv \langle i | A | j \rangle. \quad (11.2.17)$$

The expansion ([e12.14]) thus becomes

$$\langle A \rangle \equiv \langle a | A | a \rangle = \sum_{i,j} \langle a | i \rangle \langle i | A | j \rangle \langle j | a \rangle. \quad (11.2.18)$$

Incidentally, it follows that [see Equation ([e5.48])]

$$\langle i | A | j \rangle^* = \langle j | A^\dagger | i \rangle. \quad (11.2.19)$$

Finally, it is clear from Equation ([e12.20a]) that

$$\sum_i |i\rangle \langle i| \equiv 1, \quad (11.2.20)$$

where the  $\psi_i$  are a complete set of eigenstates, and 1 is the identity operator.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [11.2: Improved Notation](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 11.3: Two-State System

Consider the simplest possible non-trivial quantum mechanical system. In such a system, there are only two independent eigenstates of the unperturbed Hamiltonian: that is,

$$\begin{aligned} H_0 \psi_1 &= E_1 \psi_1, \\ H_0 \psi_2 &= E_2 \psi_2. \end{aligned}$$

It is assumed that these states, and their associated eigenvalues, are known. We also expect the states to be orthonormal, and to form a complete set.

Let us now try to solve the modified energy eigenvalue problem

$$(H_0 + H_1) \psi_E = E \psi_E. \quad (11.3.1)$$

We can, in fact, solve this problem exactly. Because the eigenstates of  $H_0$  form a complete set, we can write [see Equation (e12.13a)]

$$\psi_E = \langle 1|E \rangle \psi_1 + \langle 2|E \rangle \psi_2. \quad (11.3.2)$$

It follows from Equation (e12.23) that

$$\langle i|H_0 + H_1|E \rangle = E \langle i|E \rangle, \quad (11.3.3)$$

where  $i = 1$  or  $2$ . Equations (e12.21), (e12.22), (e12.24), (e12.25), and the orthonormality condition

$$\langle i|j \rangle = \delta_{ij}, \quad (11.3.4)$$

yield two coupled equations that can be written in matrix form:

$$\begin{pmatrix} E_1 - E + e_{11}, & e_{12} \\ e_{12}^*, & E_2 - E + e_{22} \end{pmatrix} \begin{pmatrix} \langle 1|E \rangle \\ \langle 2|E \rangle \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix},$$

where

$$\begin{aligned} e_{11} &= \langle 1|H_1|1 \rangle, \\ e_{22} &= \langle 2|H_1|2 \rangle, \\ e_{12} &= \langle 1|H_1|2 \rangle = \langle 2|H_1|1 \rangle^*. \end{aligned}$$

Here, use has been made of the fact that  $H_1$  is an Hermitian operator.

Consider the special (but not uncommon) case of a perturbing Hamiltonian whose diagonal matrix elements are zero, so that

$$e_{11} = e_{22} = 0. \quad (11.3.5)$$

The solution of Equation (e12.27) (obtained by setting the determinant of the matrix to zero) is

$$E = \frac{(E_1 + E_2) \pm \sqrt{(E_1 - E_2)^2 + 4|e_{12}|^2}}{2}. \quad (11.3.6)$$

Let us expand in the supposedly small parameter

$$\epsilon = \frac{|e_{12}|}{|E_1 - E_2|}. \quad (11.3.7)$$

We obtain

$$E \simeq \frac{1}{2} (E_1 + E_2) \pm \frac{1}{2} (E_1 - E_2) (1 + 2\epsilon^2 + \dots). \quad (11.3.8)$$

The previous expression yields the modification of the energy eigenvalues due to the perturbing Hamiltonian:

$$E'_1 = E_1 + \frac{|e_{12}|^2}{E_1 - E_2} + \dots,$$
$$E'_2 = E_2 - \frac{|e_{12}|^2}{E_1 - E_2} + \dots.$$

Note that  $H_1$  causes the upper eigenvalue to rise, and the lower to fall. It is easily demonstrated that the modified eigenstates take the form

$$\psi'_1 = \psi_1 + \frac{e_{12}^*}{E_1 - E_2} \psi_2 + \dots,$$
$$\psi'_2 = \psi_2 - \frac{e_{12}}{E_1 - E_2} \psi_1 + \dots.$$

Thus, the modified energy eigenstates consist of one of the unperturbed eigenstates, plus a slight admixture of the other. Now, our expansion procedure is only valid when  $\epsilon \ll 1$ . This suggests that the condition for the validity of the perturbation method as a whole is

$$|e_{12}| \ll |E_1 - E_2|. \quad (11.3.9)$$

In other words, when we say that  $H_1$  needs to be small compared to  $H_0$ , what we are really saying is that the previous inequality must be satisfied.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [11.3: Two-State System](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 11.4: Non-Degenerate Perturbation Theory

Let us now generalize our perturbation analysis to deal with systems possessing more than two energy eigenstates. Consider a system in which the energy eigenstates of the unperturbed Hamiltonian,  $H_0$ , are denoted

$$H_0 \psi_n = E_n \psi_n, \quad (11.4.1)$$

where  $n$  runs from 1 to  $N$ . The eigenstates are assumed to be orthonormal, so that

$$\langle m | n \rangle = \delta_{nm}, \quad (11.4.2)$$

and to form a complete set. Let us now try to solve the energy eigenvalue problem for the perturbed Hamiltonian:

$$(H_0 + H_1) \psi_E = E \psi_E. \quad (11.4.3)$$

It follows that

$$\langle m | H_0 + H_1 | E \rangle = E \langle m | E \rangle, \quad (11.4.4)$$

where  $m$  can take any value from 1 to  $N$ . Now, we can express  $\psi_E$  as a linear superposition of the unperturbed energy eigenstates:

$$\psi_E = \sum_k \langle k | E \rangle \psi_k, \quad (11.4.5)$$

where  $k$  runs from 1 to  $N$ . We can combine the previous equations to give

$$(E_m - E + e_{mm}) \langle m | E \rangle + \sum_{k \neq m} e_{mk} \langle k | E \rangle = 0, \quad (11.4.6)$$

where

$$e_{mk} = \langle m | H_1 | k \rangle. \quad (11.4.7)$$

Let us now develop our perturbation expansion. We assume that

$$\frac{e_{mk}}{E_m - E_k} \sim \mathcal{O}(\epsilon) \quad (11.4.8)$$

for all  $m \neq k$ , where  $\epsilon \ll 1$  is our expansion parameter. We also assume that

$$\frac{e_{mm}}{E_m} \sim \mathcal{O}(\epsilon) \quad (11.4.9)$$

for all  $m$ . Let us search for a modified version of the  $n$ th unperturbed energy eigenstate for which

$$E = E_n + \mathcal{O}(\epsilon), \quad (11.4.10)$$

and

$$\begin{aligned} \langle n | E \rangle &= 1, \\ \langle m | E \rangle &= \mathcal{O}(\epsilon) \end{aligned}$$

for  $m \neq n$ . Suppose that we write out Equation (11.4.6) for  $m \neq n$ , neglecting terms that are  $\mathcal{O}(\epsilon^2)$  according to our expansion scheme. We find that

$$(E_m - E_n) \langle m | E \rangle + e_{mn} \simeq 0, \quad (11.4.11)$$

giving

$$\langle m | E \rangle \simeq -\frac{e_{mn}}{E_m - E_n}. \quad (11.4.12)$$

Substituting the previous expression into Equation (11.4.6), evaluated for  $m = n$ , and neglecting  $\mathcal{O}(\epsilon^3)$  terms, we obtain

$$(E_n - E + e_{nn}) - \sum_{k \neq n} \frac{|e_{nk}|^2}{E_k - E_n} \simeq 0. \quad (11.4.13)$$



Thus, the modified  $n$ th energy eigenstate possesses an eigenvalue

$$E'_n = E_n + e_{nn} + \sum_{k \neq n} \frac{|e_{nk}|^2}{E_n - E_k} + \mathcal{O}(\epsilon^3) \quad (11.4.14)$$

and a wavefunction

$$\psi'_n = \psi_n + \sum_{k \neq n} \frac{e_{kn}}{E_n - E_k} \psi_k + \mathcal{O}(\epsilon^2). \quad (11.4.15)$$

Incidentally, it is easily demonstrated that the modified eigenstates remain orthonormal to  $\mathcal{O}(\epsilon^2)$ .

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [11.4: Non-Degenerate Perturbation Theory](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 11.5: Quadratic Stark Effect

Suppose that a hydrogen atom is subject to a uniform external electric field, of magnitude  $|\mathbf{E}|$ , directed along the  $z$ -axis. The Hamiltonian of the system can be split into two parts. Namely, the unperturbed Hamiltonian,

$$H_0 = \frac{p^2}{2m_e} - \frac{e^2}{4\pi\epsilon_0 r}, \quad (11.5.1)$$

and the perturbing Hamiltonian

$$H_1 = e |\mathbf{E}| z. \quad (11.5.2)$$

Note that the electron spin is irrelevant to this problem (because the spin operators all commute with  $H_1$ ), so we can ignore the spin degrees of freedom of the system. Hence, the energy eigenstates of the unperturbed Hamiltonian are characterized by three quantum numbers—the radial quantum number  $n$ , and the two angular quantum numbers  $l$  and  $m$ . (See Chapter [\[scent\]](#).) Let us denote these states as the  $\psi_{nlm}$ , and let their corresponding energy eigenvalues be the  $E_{nlm}$ . According to the analysis in the previous section, the change in energy of the eigenstate characterized by the quantum numbers  $n, l, m$  in the presence of a small electric field is given by

$$\begin{aligned} \Delta E_{nlm} = & e |\mathbf{E}| \langle n, l, m | z | n, l, m \rangle \\ & + e^2 |\mathbf{E}|^2 \sum_{n', l', m' \neq n, l, m} \frac{|\langle n, l, m | z | n', l', m' \rangle|^2}{E_{nlm} - E_{n'l'm'}}. \end{aligned}$$

This energy-shift is known as the *Stark effect*.

The sum on the right-hand side of the previous equation seems very complicated. However, it turns out that most of the terms in this sum are zero. This follows because the matrix elements  $\langle n, l, m | z | n', l', m' \rangle$  are zero for virtually all choices of the two sets of quantum number,  $n, l, m$  and  $n', l', m'$ . Let us try to find a set of rules that determine when these matrix elements are non-zero. These rules are usually referred to as the *selection rules* for the problem in hand.

Now, because [see Equation ([\[e8.3\]](#))]

$$L_z = x p_y - y p_x, \quad (11.5.3)$$

it follows that [see Equations ([\[commxx\]](#))–([\[commxp\]](#))]

$$[L_z, z] = 0. \quad (11.5.4)$$

Thus,

$$\begin{aligned} \langle n, l, m | [L_z, z] | n', l', m' \rangle &= \langle n, l, m | L_z z - z L_z | n', l', m' \rangle \\ &= \hbar (m - m') \langle n, l, m | z | n', l', m' \rangle = 0, \end{aligned}$$

because  $\psi_{nlm}$  is, by definition, an eigenstate of  $L_z$  corresponding to the eigenvalue  $m\hbar$ . Hence, it is clear, from the previous equation, that one of the selection rules is that the matrix element  $\langle n, l, m | z | n', l', m' \rangle$  is zero unless

$$m' = m. \quad (11.5.5)$$

Let us now determine the selection rule for  $l$ . We have

$$\begin{aligned} &= [L_x^2, z] + [L_y^2, z] \\ &= L_x [L_x, z] + [L_x, z] L_x + L_y [L_y, z] + [L_y, z] L_y \\ &= i\hbar (-L_x y - y L_x + L_y x + x L_y) \\ &= 2i\hbar (L_y x - L_x y + i\hbar z) \\ &= 2i\hbar (L_y x - y L_x) = 2i\hbar (x L_y - L_x y), \end{aligned}$$

where use has been made of Equations ([\[commxx\]](#))–([\[commxp\]](#)), ([\[e8.1\]](#))–([\[e8.3\]](#)), and ([\[e8.10\]](#)). Thus,

$$\begin{aligned}
 ] &= 2i\hbar (L^2, L_y x - L_x y + i\hbar z) \\
 &= 2i\hbar (L_y [L^2, x] - L_x [L^2, y] + i\hbar [L^2, z]) \\
 &= -4\hbar^2 L_y (y L_z - L_y z) + 4\hbar^2 L_x (L_x z - x L_z) \\
 &\quad - 2\hbar^2 (L^2 z - z L^2),
 \end{aligned}$$

which reduces to

$$\begin{aligned}
 ] &= -\hbar^2 \{4(L_x x + L_y y + L_z z) L_z - 4(L_x^2 + L_y^2 + L_z^2) z \\
 &\quad + 2(L^2 z - z L^2)\} \\
 &= -\hbar^2 \{4(L_x x + L_y y + L_z z) L_z - 2(L^2 z + z L^2)\}.
 \end{aligned}$$

However, it is clear from Equations ([e8.1])–([e8.3]) that

$$L_x x + L_y y + L_z z = 0. \quad (11.5.6)$$

Hence, we obtain

$$[L^2, [L^2, z]] = 2\hbar^2 (L^2 z + z L^2). \quad (11.5.7)$$

Finally, the previous expression expands to give

$$L^4 z - 2L^2 z L^2 + z L^4 - 2\hbar^2 (L^2 z + z L^2) = 0. \quad (11.5.8)$$

Equation ([e12.69]) implies that

$$\langle n, l, m | L^4 z - 2L^2 z L^2 + z L^4 - 2\hbar^2 (L^2 z + z L^2) | n', l', m \rangle = 0. \quad (11.5.9)$$

Because, by definition,  $\psi_{nlm}$  is an eigenstate of  $L^2$  corresponding to the eigenvalue  $l(l+1)\hbar^2$ , this expression yields

$$\begin{aligned}
 \{l^2(l+1)^2 - 2l(l+1)l'(l'+1) + l'^2(l'+1)^2 \\
 - 2l(l+1) - 2l'(l'+1)\} \langle n, l, m | z | n', l', m \rangle = 0,
 \end{aligned}$$

which reduces to

$$(l+l'+2)(l+l')(l-l'+1)(l-l'-1) \langle n, l, m | z | n', l', m \rangle = 0. \quad (11.5.10)$$

According to the previous formula, the matrix element  $\langle n, l, m | z | n', l', m \rangle$  vanishes unless  $l = l' = 0$  or  $l' = l \pm 1$ . [Of course, the factor  $l+l'+2$ , in the previous equation, can never be zero, because  $l$  and  $l'$  can never be negative.] Recall, however, from Chapter [scent], that an  $l = 0$  wavefunction is spherically symmetric. It, therefore, follows, from symmetry, that the matrix element  $\langle n, l, m | z | n', l', m \rangle$  is zero when  $l = l' = 0$ . In conclusion, the selection rule for  $l$  is that the matrix element  $\langle n, l, m | z | n', l', m \rangle$  is zero unless

$$l' = l \pm 1. \quad (11.5.11)$$

Application of the selection rules ([e12.63]) and ([e12.73]) to Equation ([e12.59]) yields

$$\Delta E_{nlm} = e^2 |\mathbf{E}|^2 \sum_{n', l' = l \pm 1} \frac{|\langle n, l, m | z | n', l', m \rangle|^2}{E_{nlm} - E_{n'l'm}}. \quad (11.5.12)$$

Note that, according to the selection rules, all of the terms in Equation ([e12.59]) that vary linearly with the electric field-strength vanish. Only those terms which vary quadratically with the field-strength survive. Hence, this type of energy-shift of an atomic state in the presence of a small electric field is known as the *quadratic Stark effect*. Now, the *electric polarizability* of an atom is defined in terms of the energy-shift of the atomic state as follows :

$$\Delta E = -\frac{1}{2} \alpha |\mathbf{E}|^2. \quad (11.5.13)$$

Hence, we can write

$$\alpha_{nlm} = 2e^2 \sum_{n', l' = l \pm 1} \frac{|\langle n, l, m | z | n', l', m \rangle|^2}{E_{n'l'm} - E_{nlm}}. \quad (11.5.14)$$

Unfortunately, there is one fairly obvious problem with Equation (11.5.14). Namely, it predicts an infinite energy-shift if there exists some non-zero matrix element  $\langle n, l, m | z | n', l', m \rangle$  that couples two degenerate unperturbed energy eigenstates: that is, if  $\langle n, l, m | z | n', l', m \rangle \neq 0$  and  $E_{nlm} = E_{n'l'm}$ . Clearly, our perturbation method breaks down completely in this situation. Hence, we conclude that Equations (11.5.14) and (11.5.16) are only applicable to cases where the coupled eigenstates are non-degenerate. For this reason, the type of perturbation theory employed here is known as *non-degenerate perturbation theory*. The unperturbed eigenstates of a hydrogen atom have energies that only depend on the radial quantum number  $n$ . (See Chapter 9.) It follows that we can only apply the previous results to the  $n = 1$  eigenstate (because for  $n > 1$  there will be coupling to degenerate eigenstates with the same value of  $n$  but different values of  $l$ ).

Thus, according to non-degenerate perturbation theory, the polarizability of the ground-state (i.e.,  $n = 1$ ) of a hydrogen atom is given by

$$\alpha = 2e^2 \sum_{n>1} \frac{|\langle 1, 0, 0 | z | n, 1, 0 \rangle|^2}{E_{n00} - E_{100}}. \quad (11.5.15)$$

Here, we have made use of the fact that  $E_{n10} = E_{n00}$ . The sum in the previous expression can be evaluated approximately by noting that (see Section 10.4)

$$E_{n00} = -\frac{e^2}{8\pi\epsilon_0 a_0 n^2}, \quad (11.5.16)$$

where

$$a_0 = \frac{4\pi\epsilon_0 \hbar^2}{m_e e^2} \quad (11.5.17)$$

is the Bohr radius. Hence, we can write

$$E_{n00} - E_{100} \geq E_{200} - E_{100} = \frac{3}{4} \frac{e^2}{8\pi\epsilon_0 a_0}, \quad (11.5.18)$$

which implies that

$$\alpha < \frac{16}{3} 4\pi\epsilon_0 a_0 \sum_{n>1} |\langle 1, 0, 0 | z | n, 1, 0 \rangle|^2. \quad (11.5.19)$$

However, [see Equation (11.5.20)]

$$\begin{aligned} \sum_{n>1} |\langle 1, 0, 0 | z | n, 1, 0 \rangle|^2 &= \sum_{n>1} \langle 1, 0, 0 | z | n, 1, 0 \rangle \langle n, 1, 0 | z | 1, 0, 0 \rangle \\ &= \sum_{n', l', m'} \langle 1, 0, 0 | z | n', l', m' \rangle \langle n', l', m' | z | 1, 0, 0 \rangle \\ &= \langle 1, 0, 0 | z^2 | 1, 0, 0 \rangle = \frac{1}{3} \langle 1, 0, 0 | r^2 | 1, 0, 0 \rangle, \end{aligned}$$

where we have made use of the selection rules, the fact that the  $\psi_{n', l', m'}$  form a complete set, and the fact the the ground-state of hydrogen is spherically symmetric. Finally, it follows from Equation (9.73) that

$$\langle 1, 0, 0 | r^2 | 1, 0, 0 \rangle = 3a_0^2. \quad (11.5.20)$$

Hence, we conclude that

$$\alpha < \frac{16}{3} 4\pi\epsilon_0 a_0^3 \simeq 5.3 \cdot 4\pi\epsilon_0 a_0^3. \quad (11.5.21)$$

The exact result (which can be obtained by solving Schrödinger's equation in parabolic coordinates) is

$$\alpha = \frac{9}{2} 4\pi \epsilon_0 a_0^3 = 4.5 4\pi \epsilon_0 a_0^3. \quad (11.5.22)$$

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [11.5: Quadratic Stark Effect](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 11.6: Degenerate Perturbation Theory

Let us, rather naively, investigate the Stark effect in an excited (i.e.,  $n > 1$ ) state of the hydrogen atom using standard non-degenerate perturbation theory. We can write

$$H_0 \psi_{nlm} = E_n \psi_{nlm}, \quad (11.6.1)$$

because the energy eigenstates of the unperturbed Hamiltonian only depend on the quantum number  $n$ . Making use of the selection rules ([e12.63]) and ([e12.73]), non-degenerate perturbation theory yields the following expressions for the perturbed energy levels and eigenstates [see Equations ([e12.56]) and ([e12.57])]:

$$E'_{nl} = E_n + e_{nl} + \sum_{n', l' = l \pm 1} \frac{|e_{n'l'nl}|^2}{E_n - E_{n'}}, \quad (11.6.2)$$

and

$$\psi'_{nlm} = \psi_{nlm} + \sum_{n', l' = l \pm 1} \frac{e_{n'l'nl}}{E_n - E_{n'}} \psi_{n'l'm}, \quad (11.6.3)$$

where

$$e_{n'l'nl} = \langle n', l', m | H_1 | n, l, m \rangle. \quad (11.6.4)$$

Unfortunately, if  $n > 1$  then the summations in the previous expressions are not well defined, because there exist non-zero matrix elements,  $e_{n'l'nl}$ , that couple degenerate eigenstates: that is, there exist non-zero matrix elements that couple states with the same value of  $n$ , but different values of  $l$ . These particular matrix elements give rise to singular factors  $1/(E_n - E_n)$  in the summations. This does not occur if  $n = 1$  because, in this case, the selection rule  $l' = l \pm 1$ , and the fact that  $l = 0$  (because  $0 \leq l < n$ ), only allow  $l'$  to take the single value 1. Of course, there is no  $n = 1$  state with  $l' = 1$ . Hence, there is only one coupled state corresponding to the eigenvalue  $E_1$ . Unfortunately, if  $n > 1$  then there are multiple coupled states corresponding to the eigenvalue  $E_n$ .

Note that our problem would disappear if the matrix elements of the perturbed Hamiltonian corresponding to the same value of  $n$ , but different values of  $l$ , were all zero: that is, if

$$\langle n, l', m | H_1 | n, l, m \rangle = \lambda_{nl} \delta_{ll'}. \quad (11.6.5)$$

In this case, all of the singular terms in Equations ([e12.88]) and ([e12.89]) would reduce to zero. Unfortunately, the previous equation is not satisfied in general. Fortunately, we can always redefine the unperturbed eigenstates corresponding to the eigenvalue  $E_n$  in such a manner that Equation ([e12.91]) is satisfied. Suppose that there are  $N_n$  coupled eigenstates belonging to the eigenvalue  $E_n$ . Let us define  $N_n$  new states which are linear combinations of our  $N_n$  original degenerate eigenstates:

$$\psi_{nlm}^{(1)} = \sum_{k=1, N_n} \langle n, k, m | n, l^{(1)}, m \rangle \psi_{nkm}. \quad (11.6.6)$$

Note that these new states are also degenerate energy eigenstates of the unperturbed Hamiltonian,  $H_0$ , corresponding to the eigenvalue  $E_n$ . The  $\psi_{nlm}^{(1)}$  are chosen in such a manner that they are also eigenstates of the perturbing Hamiltonian,  $H_1$ : that is, they are simultaneous eigenstates of  $H_0$  and  $H_1$ . Thus,

$$H_1 \psi_{nlm}^{(1)} = \lambda_{nl} \psi_{nlm}^{(1)}. \quad (11.6.7)$$

The  $\psi_{nlm}^{(1)}$  are also chosen so as to be orthonormal: that is,

$$\langle n, l'^{(1)}, m | n, l^{(1)}, m \rangle = \delta_{ll'}. \quad (11.6.8)$$

It follows that

$$\langle n, l'^{(1)}, m | H_1 | n, l^{(1)}, m \rangle = \lambda_{nl} \delta_{ll'}. \quad (11.6.9)$$

Thus, if we use the new eigenstates, instead of the old ones, then we can employ Equations ([e12.88]) and ([e12.89]) directly, because all of the singular terms vanish. The only remaining difficulty is to determine the new eigenstates in terms of the original

ones.

Now [see Equation ([\[e12.20\]](#))]

$$\sum_{l=1, N_n} |n, l, m\rangle \langle n, l, m| \equiv 1, \quad (11.6.10)$$

where 1 denotes the identity operator in the sub-space of all coupled unperturbed eigenstates corresponding to the eigenvalue  $E_n$ . Using this completeness relation, the eigenvalue equation ([\[e12.93\]](#)) can be transformed into a straightforward matrix equation:

$$\sum_{l''=1, N_n} \langle n, l', m | H_1 | n, l'', m \rangle \langle n, l'', m | n, l^{(1)}, m \rangle = \lambda_{nl} \langle n, l', m | n, l^{(1)}, m \rangle. \quad (11.6.11)$$

This can be written more transparently as

$$\mathbf{U} \mathbf{x} = \lambda \mathbf{x}, \quad (11.6.12)$$

where the elements of the  $N_n \times N_n$  Hermitian matrix  $\mathbf{U}$  are

$$U_{jk} = \langle n, j, m | H_1 | n, k, m \rangle. \quad (11.6.13)$$

Provided that the determinant of  $\mathbf{U}$  is non-zero, Equation ([\[e12.100\]](#)) can always be solved to give  $N_n$  eigenvalues  $\lambda_{nl}$  (for  $l = 1$  to  $N_n$ ), with  $N_n$  corresponding eigenvectors  $\mathbf{x}_{nl}$ . The normalized eigenvectors specify the weights of the new eigenstates in terms of the original eigenstates: that is,

$$(\mathbf{x}_{nl})_k = \langle n, k, m | n, l^{(1)}, m \rangle, \quad (11.6.14)$$

for  $k = 1$  to  $N_n$ . In our new scheme, Equations ([\[e12.88\]](#)) and ([\[e12.89\]](#)) yield

$$E'_{nl} = E_n + \lambda_{nl} + \sum_{n' \neq n, l' = l \pm 1} \frac{|e_{n'l'nl}|^2}{E_n - E_{n'}}, \quad (11.6.15)$$

and

$$\psi_{nlm}^{(1)'} = \psi_{nlm}^{(1)} + \sum_{n' \neq n, l' = l \pm 1} \frac{e_{n'l'nl}}{E_n - E_{n'}} \psi_{n'l'm}. \quad (11.6.16)$$

There are no singular terms in these expressions, because the summations are over  $n' \neq n$ : that is, they specifically exclude the problematic, degenerate, unperturbed energy eigenstates corresponding to the eigenvalue  $E_n$ . Note that the first-order energy shifts are equivalent to the eigenvalues of the matrix equation ([\[e12.100\]](#)).

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [11.6: Degenerate Perturbation Theory](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 11.7: Linear Stark Effect

Returning to the Stark effect, let us examine the effect of an external electric field on the energy levels of the  $n = 2$  states of a hydrogen atom. There are four such states: an  $l = 0$  state, usually referred to as  $2S$ , and three  $l = 1$  states (with  $m = -1, 0, 1$ ), usually referred to as  $2P$ . All of these states possess the same unperturbed energy,  $E_{200} = -e^2 / (32\pi \epsilon_0 a_0)$ . As before, the perturbing Hamiltonian is

$$H_1 = e |\mathbf{E}| z. \quad (11.7.1)$$

According to the previously determined selection rules (i.e.,  $m' = m$ , and  $l' = l \pm 1$ ), this Hamiltonian couples  $\psi_{200}$  and  $\psi_{210}$ . Hence, non-degenerate perturbation theory breaks down when applied to these two states. On the other hand, non-degenerate perturbation theory works fine for the  $\psi_{211}$  and  $\psi_{21-1}$  states, because these are not coupled to any other  $n = 2$  states by the perturbing Hamiltonian.

In order to apply perturbation theory to the  $\psi_{200}$  and  $\psi_{210}$  states, we have to solve the matrix eigenvalue equation

$$\mathbf{U} \mathbf{x} = \lambda \mathbf{x}, \quad (11.7.2)$$

where  $\mathbf{U}$  is the matrix of the matrix elements of  $H_1$  between these states. Thus,

$$\mathbf{U} = e |\mathbf{E}| \begin{pmatrix} 0, & \langle 2, 0, 0 | z | 2, 1, 0 \rangle \\ \langle 2, 1, 0 | z | 2, 0, 0 \rangle, & 0 \end{pmatrix}, \quad (11.7.3)$$

where the rows and columns correspond to  $\psi_{200}$  and  $\psi_{210}$ , respectively. Here, we have again made use of the selection rules, which tell us that the matrix element of  $z$  between two hydrogen atom states is zero unless the states possess  $l$  quantum numbers that differ by unity. It is easily demonstrated, from the exact forms of the  $2S$  and  $2P$  wavefunctions, that

$$\langle 2, 0, 0 | z | 2, 1, 0 \rangle = \langle 2, 1, 0 | z | 2, 0, 0 \rangle = 3 a_0. \quad (11.7.4)$$

It can be seen, by inspection, that the eigenvalues of  $\mathbf{U}$  are  $\lambda_1 = 3 e a_0 |\mathbf{E}|$  and  $\lambda_2 = -3 e a_0 |\mathbf{E}|$ . The corresponding normalized eigenvectors are

$$\begin{aligned} \mathbf{x}_1 &= \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}, \\ \mathbf{x}_2 &= \begin{pmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \end{pmatrix}. \end{aligned}$$

It follows that the simultaneous eigenstates of  $H_0$  and  $H_1$  take the form

$$\begin{aligned} \psi_1 &= \frac{\psi_{200} + \psi_{210}}{\sqrt{2}}, \\ \psi_2 &= \frac{\psi_{200} - \psi_{210}}{\sqrt{2}}. \end{aligned}$$

In the absence of an external electric field, both of these states possess the same energy,  $E_{200}$ . The first-order energy shifts induced by an external electric field are given by

$$\begin{aligned} \Delta E_1 &= +3 e a_0 |\mathbf{E}|, \\ \Delta E_2 &= -3 e a_0 |\mathbf{E}|. \end{aligned}$$

Thus, in the presence of an electric field, the energies of states 1 and 2 are shifted upwards and downwards, respectively, by an amount  $3 e a_0 |\mathbf{E}|$ . These states are orthogonal linear combinations of the original  $\psi_{200}$  and  $\psi_{210}$  states. Note that the energy shifts are linear in the electric field-strength, so this effect—which is known as the *linear Stark effect*—is much larger than the quadratic effect described in Section 1.5. Note, also, that the energies of the  $\psi_{211}$  and  $\psi_{21-1}$  states are not affected by the electric field to first-order. Of course, to second-order the energies of these states are shifted by an amount which depends on the square of the electric field-strength. (See Section 1.5.)



## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [11.7: Linear Stark Effect](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 11.8: Fine Structure of Hydrogen

According to special relativity, the kinetic energy (i.e., the difference between the total energy and the rest mass energy) of a particle of rest mass  $m$  and momentum  $p$  is

$$T = \sqrt{p^2 c^2 + m^2 c^4} - m c^2. \quad (11.8.1)$$

In the non-relativistic limit  $p \ll m c$ , we can expand the square-root in the previous expression to give

$$T = \frac{p^2}{2m} \left[ 1 - \frac{1}{4} \left( \frac{p}{m c} \right)^2 + \mathcal{O} \left( \frac{p}{m c} \right)^4 \right]. \quad (11.8.2)$$

Hence,

$$T \simeq \frac{p^2}{2m} - \frac{p^4}{8m^3 c^2}. \quad (11.8.3)$$

Of course, we recognize the first term on the right-hand side of this equation as the standard non-relativistic expression for the kinetic energy. The second term is the lowest-order relativistic correction to this energy. Let us consider the effect of this type of correction on the energy levels of a hydrogen atom. So, the unperturbed Hamiltonian is given by Equation ([e12.58]), and the perturbing Hamiltonian takes the form

$$H_1 = -\frac{p^4}{8m_e^3 c^2}. \quad (11.8.4)$$

Now, according to standard first-order perturbation theory (see Section 1.4), the lowest-order relativistic correction to the energy of a hydrogen atom state characterized by the standard quantum numbers  $n$ ,  $l$ , and  $m$  is given by

$$\begin{aligned} \Delta E_{nlm} &= \langle n, l, m | H_1 | n, l, m \rangle = -\frac{1}{8m_e^3 c^2} \langle n, l, m | p^4 | n, l, m \rangle \\ &= -\frac{1}{8m_e^3 c^2} \langle n, l, m | p^2 p^2 | n, l, m \rangle. \end{aligned}$$

However, Schrödinger's equation for a unperturbed hydrogen atom can be written

$$p^2 \psi_{n,l,m} = 2m_e (E_n - V) \psi_{n,l,m}, \quad (11.8.5)$$

where  $V = -e^2 / (4\pi \epsilon_0 r)$ . Because  $p^2$  is an Hermitian operator, it follows that

$$\begin{aligned} \Delta E_{nlm} &= -\frac{1}{2m_e c^2} \langle n, l, m | (E_n - V)^2 | n, l, m \rangle \\ &= -\frac{1}{2m_e c^2} (E_n^2 - 2E_n \langle n, l, m | V | n, l, m \rangle + \langle n, l, m | V^2 | n, l, m \rangle) \\ &= -\frac{1}{2m_e c^2} \left[ E_n^2 + 2E_n \left( \frac{e^2}{4\pi \epsilon_0} \right) \left\langle \frac{1}{r} \right\rangle + \left( \frac{e^2}{4\pi \epsilon_0} \right)^2 \left\langle \frac{1}{r^2} \right\rangle \right]. \end{aligned}$$

It follows from Equations ([e9.74]) and ([e9.75]) that

$$\Delta E_{nlm} = -\frac{1}{2m_e c^2} \left[ E_n^2 + 2E_n \left( \frac{e^2}{4\pi \epsilon_0} \right) \frac{1}{n^2 a_0} + \left( \frac{e^2}{4\pi \epsilon_0} \right)^2 \frac{1}{(l+1/2) n^3 a_0^2} \right].$$

Finally, making use of Equations ([e9.55]), ([e9.56]), and ([e9.57]), the previous expression reduces to

$$\Delta E_{nlm} = E_n \frac{\alpha^2}{n^2} \left( \frac{n}{l+1/2} - \frac{3}{4} \right), \quad (11.8.6)$$

where

$$\alpha = \frac{e^2}{4\pi\epsilon_0\hbar c} \simeq \frac{1}{137} \quad (11.8.7)$$

is the dimensionless *fine structure constant*.

Note that the previous derivation implicitly assumes that  $p^4$  is an Hermitian operator. It turns out that this is not the case for  $l = 0$  states. However, somewhat fortuitously, our calculation still gives the correct answer when  $l = 0$ . Note, also, that we are able to employ non-degenerate perturbation theory in the previous calculation, using the  $\psi_{nlm}$  eigenstates, because the perturbing Hamiltonian commutes with both  $L^2$  and  $L_z$ . It follows that there is no coupling between states with different  $l$  and  $m$  quantum numbers. Hence, all coupled states have different  $n$  quantum numbers, and therefore have different energies.

Now, an electron in a hydrogen atom experiences an electric field

$$\mathbf{E} = \frac{e\mathbf{r}}{4\pi\epsilon_0 r^3} \quad (11.8.8)$$

due to the charge on the nucleus. However, according to electromagnetic theory, a non-relativistic particle moving in a electric field  $\mathbf{E}$  with velocity  $\mathbf{v}$  also experiences an effective magnetic field

$$\mathbf{B} = -\frac{\mathbf{v} \times \mathbf{E}}{c^2} \quad (11.8.9)$$

Recall, that an electron possesses a magnetic moment [see Equations ([e10.58]) and ([e10.59])]

$$\mu = -\frac{e}{m_e} \mathbf{S} \quad (11.8.10)$$

due to its spin angular momentum,  $\mathbf{S}$ . We, therefore, expect an additional contribution to the Hamiltonian of a hydrogen atom of the form [see Equation ([e10.60a])]

$$H_1 = -\mu \cdot \mathbf{B} = -\frac{e^2}{4\pi\epsilon_0 m_e c^2 r^3} \mathbf{v} \times \mathbf{r} \cdot \mathbf{S} = \frac{e^2}{4\pi\epsilon_0 m_e^2 c^2 r^3} \mathbf{L} \cdot \mathbf{S},$$

where  $\mathbf{L} = m_e \mathbf{r} \times \mathbf{v}$  is the electron's orbital angular momentum. This effect is known as *spin-orbit coupling*. It turns out that the previous expression is too large, by a factor 2, due to an obscure relativistic effect known as *Thomas precession*. Hence, the true spin-orbit correction to the Hamiltonian is

$$H_1 = \frac{e^2}{8\pi\epsilon_0 m_e^2 c^2 r^3} \mathbf{L} \cdot \mathbf{S}. \quad (11.8.11)$$

Let us now apply perturbation theory to the hydrogen atom, using the previous expression as the perturbing Hamiltonian.

Now,

$$\mathbf{J} = \mathbf{L} + \mathbf{S} \quad (11.8.12)$$

is the total angular momentum of the system. Hence,

$$J^2 = L^2 + S^2 + 2\mathbf{L} \cdot \mathbf{S}, \quad (11.8.13)$$

giving

$$\mathbf{L} \cdot \mathbf{S} = \frac{1}{2} (J^2 - L^2 - S^2). \quad (11.8.14)$$

Recall, from Section [s11.2], that while  $J^2$  commutes with both  $L^2$  and  $S^2$ , it does not commute with either  $L_z$  or  $S_z$ . It follows that the perturbing Hamiltonian ([e12.127]) also commutes with both  $L^2$  and  $S^2$ , but does not commute with either  $L_z$  or  $S_z$ . Hence, the simultaneous eigenstates of the unperturbed Hamiltonian ([e12.58]) and the perturbing Hamiltonian ([e12.127]) are the same as the simultaneous eigenstates of  $L^2$ ,  $S^2$ , and  $J^2$  discussed in Section [s11.3]. It is important to know this because, according to Section 1.6, we can only safely apply perturbation theory to the simultaneous eigenstates of the unperturbed and perturbing Hamiltonians.

Adopting the notation introduced in Section [s11.3], let  $\psi_{l,s;j,m_j}^{(2)}$  be a simultaneous eigenstate of  $L^2$ ,  $S^2$ ,  $J^2$ , and  $J_z$  corresponding to the eigenvalues

$$\begin{aligned}
 L^2 \psi_{l,s;j,m_j}^{(2)} &= l(l+1) \hbar^2 \psi_{l,s;j,m_j}^{(2)}, \\
 S^2 \psi_{l,s;j,m_j}^{(2)} &= s(s+1) \hbar^2 \psi_{l,s;j,m_j}^{(2)}, \\
 J^2 \psi_{l,s;j,m_j}^{(2)} &= j(j+1) \hbar^2 \psi_{l,s;j,m_j}^{(2)}, \\
 J_z \psi_{l,s;j,m_j}^{(2)} &= m_j \hbar \psi_{l,s;j,m_j}^{(2)}.
 \end{aligned}$$

According to standard first-order perturbation theory, the energy-shift induced in such a state by spin-orbit coupling is given by

$$\begin{aligned}
 \Delta E_{l,1/2;j,m_j} &= \langle l, 1/2; j, m_j | H_1 | l, 1/2; j, m_j \rangle \\
 &= \frac{e^2}{16\pi \epsilon_0 m_e^2 c^2} \left\langle 1, 1/2; j, m_j \left| \frac{J^2 - L^2 - S^2}{r^3} \right| l, 1/2; j, m_j \right\rangle \\
 &= \frac{e^2 \hbar^2}{16\pi \epsilon_0 m_e^2 c^2} [j(j+1) - l(l+1) - 3/4] \left\langle \frac{1}{r^3} \right\rangle.
 \end{aligned}$$

Here, we have made use of the fact that  $s = 1/2$  for an electron. It follows from Equation ([e9.75a](#)) that

$$\Delta E_{l,1/2;j,m_j} = \frac{e^2 \hbar^2}{16\pi \epsilon_0 m_e^2 c^2 a_0^3} \left[ \frac{j(j+1) - l(l+1) - 3/4}{l(l+1/2)(l+1)n^3} \right], \quad (11.8.15)$$

where  $n$  is the radial quantum number. Finally, making use of Equations ([e9.55](#)), ([e9.56](#)), and ([e9.57](#)), the previous expression reduces to

$$\Delta E_{l,1/2;j,m_j} = E_n \frac{\alpha^2}{n^2} \left[ \frac{n \{3/4 + l(l+1) - j(j+1)\}}{2l(l+1/2)(l+1)} \right], \quad (11.8.16)$$

where  $\alpha$  is the fine structure constant. A comparison of this expression with Equation ([e12.121](#)) reveals that the energy-shift due to spin-orbit coupling is of the same order of magnitude as that due to the lowest-order relativistic correction to the Hamiltonian. We can add these two corrections together (making use of the fact that  $j = l \pm 1/2$  for a hydrogen atom—see Section [s11.3](#)) to obtain a net energy-shift of

$$\Delta E_{l,1/2;j,m_j} = E_n \frac{\alpha^2}{n^2} \left( \frac{n}{j+1/2} - \frac{3}{4} \right). \quad (11.8.17)$$

This modification of the energy levels of a hydrogen atom due to a combination of relativity and spin-orbit coupling is known as *fine structure*.


Now, it is conventional to refer to the energy eigenstates of a hydrogen atom that are also simultaneous eigenstates of  $J^2$  as  $nL_j$  states, where  $n$  is the radial quantum number,  $L = (S, P, D, F, \dots)$  as  $l = (0, 1, 2, 3, \dots)$ , and  $j$  is the total angular momentum quantum number. Let us examine the effect of the fine structure energy-shift ([e12.138](#)) on these eigenstates for  $n = 1, 2$  and 3.

For  $n = 1$ , in the absence of fine structure, there are two degenerate  $1S_{1/2}$  states. According to Equation ([e12.138](#)), the fine structure induced energy-shifts of these two states are the same. Hence, fine structure does not break the degeneracy of the two  $1S_{1/2}$  states of hydrogen.

For  $n = 2$ , in the absence of fine structure, there are two  $2S_{1/2}$  states, two  $2P_{1/2}$  states, and four  $2P_{3/2}$  states, all of which are degenerate. According to Equation ([e12.138](#)), the fine structure induced energy-shifts of the  $2S_{1/2}$  and  $2P_{1/2}$  states are the same as one another, but are different from the induced energy-shift of the  $2P_{3/2}$  states. Hence, fine structure does not break the degeneracy of the  $2S_{1/2}$  and  $2P_{1/2}$  states of hydrogen, but does break the degeneracy of these states relative to the  $2P_{3/2}$  states.

For  $n = 3$ , in the absence of fine structure, there are two  $3S_{1/2}$  states, two  $3P_{1/2}$  states, four  $3P_{3/2}$  states, four  $3D_{3/2}$  states, and six  $3D_{5/2}$  states, all of which are degenerate. According to Equation ([e12.138](#)), fine structure breaks these states into three groups: the  $3S_{1/2}$  and  $3P_{1/2}$  states, the  $3P_{3/2}$  and  $3D_{3/2}$  states, and the  $3D_{5/2}$  states.

The effect of the fine structure energy-shift on the  $n = 1, 2$ , and 3 energy states of a hydrogen atom is illustrated in Figure below:



unperturbed + fine structure

Figure 23: *Effect of the fine structure energy-shift on the  $n = 1, 2$  and  $3$  states of a hydrogen atom. Not to scale.*

Note, finally, that although expression ([e12.137]) does not have a well defined value for  $l = 0$ , when added to expression ([e12.121]) it, somewhat fortuitously, gives rise to an expression ([e12.138]) that is both well-defined and correct when  $l = 0$ .

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [11.8: Fine Structure of Hydrogen](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 11.9: Zeeman Effect

Consider a hydrogen atom placed in a uniform  $z$ -directed external magnetic field of magnitude  $|\mathbf{B}|$ . The modification to the Hamiltonian of the system is

$$H_1 = -\boldsymbol{\mu} \cdot \mathbf{B}, \quad (11.9.1)$$

where

$$\boldsymbol{\mu} = -\frac{e}{2m_e} (\mathbf{L} + 2\mathbf{S}) \quad (11.9.2)$$

is the total electron magnetic moment, including both orbital and spin contributions. [See Equations ([\[e10.57\]](#))–([\[e10.59\]](#)).] Thus,

$$H_1 = \frac{eB}{2m_e} (L_z + 2S_z). \quad (11.9.3)$$

Suppose that the applied magnetic field is much weaker than the atom's internal magnetic field, ([\[e12.124\]](#)). Because the magnitude of the internal field is about 25 tesla, this is a fairly reasonable assumption. In this situation, we can treat  $H_1$  as a small perturbation acting on the simultaneous eigenstates of the unperturbed Hamiltonian and the fine structure Hamiltonian. Of course, these states are the simultaneous eigenstates of  $L^2$ ,  $S^2$ ,  $J^2$ , and  $J_z$ . (See the previous section.) Hence, from standard perturbation theory, the first-order energy-shift induced by a weak external magnetic field is

$$\begin{aligned} \Delta E_{l,1/2;j,m_j} &= \langle l, 1/2; j, m_j | H_1 | l, 1/2; j, m_j \rangle \\ &= \frac{eB}{2m_e} (m_j \hbar + \langle l, 1/2; j, m_j | S_z | l, 1/2; j, m_j \rangle), \end{aligned}$$

because  $J_z = L_z + S_z$ . Now, according to Equations ([\[e11.47\]](#)) and ([\[e11.48\]](#)),

$$\psi_{j,m_j}^{(2)} = \left( \frac{j+m_j}{2l+1} \right)^{1/2} \psi_{m_j-1/2,1/2}^{(1)} + \left( \frac{j-m_j}{2l+1} \right)^{1/2} \psi_{m_j+1/2,-1/2}^{(1)} \quad (11.9.4)$$

when  $j = l + 1/2$ , and

$$\psi_{j,m_j}^{(2)} = \left( \frac{j+1-m_j}{2l+1} \right)^{1/2} \psi_{m_j-1/2,1/2}^{(1)} - \left( \frac{j+1+m_j}{2l+1} \right)^{1/2} \psi_{m_j+1/2,-1/2}^{(1)} \quad (11.9.5)$$

when  $j = l - 1/2$ . Here, the  $\psi_{m,m_s}^{(1)}$  are the simultaneous eigenstates of  $L^2$ ,  $S^2$ ,  $L_z$ , and  $S_z$ , whereas the  $\psi_{j,m_j}^{(2)}$  are the simultaneous eigenstates of  $L^2$ ,  $S^2$ ,  $J^2$ , and  $J_z$ . In particular,

$$S_z \psi_{m,\pm 1/2}^{(1)} = \pm \frac{\hbar}{2} \psi_{m,\pm 1/2}^{(1)}. \quad (11.9.6)$$

It follows from Equations ([\[e12.143\]](#))–([\[e12.145\]](#)), and the orthormality of the  $\psi^{(1)}$ , that

$$\langle l, 1/2; j, m_j | S_z | l, 1/2; j, m_j \rangle = \pm \frac{m_j \hbar}{2l+1} \quad (11.9.7)$$

when  $j = l \pm 1/2$ . Thus, the induced energy-shift when a hydrogen atom is placed in an external magnetic field—which is known as the *Zeeman effect*—becomes

$$\Delta E_{l,1/2;j,m_j} = \mu_B B m_j \left( 1 \pm \frac{1}{2l+1} \right) \quad (11.9.8)$$

where the  $\pm$  signs correspond to  $j = l \pm 1/2$ . Here,

$$\mu_B = \frac{e\hbar}{2m_e} = 5.788 \times 10^{-5} \text{ eV/T} \quad (11.9.9)$$

is known as the *Bohr magnetron*. Of course, the quantum number  $m_j$  takes values differing by unity in the range  $-j$  to  $j$ . It, thus, follows from Equation ([\[e12.147\]](#)) that the Zeeman effect splits degenerate states characterized by  $j = l + 1/2$  into  $2j + 1$  equally spaced states of interstate spacing

$$\Delta E_{j=l+1/2} = \mu_B B \left( \frac{2l+2}{2l+1} \right). \quad (11.9.10)$$

Likewise, the Zeeman effect splits degenerate states characterized by  $j = l - 1/2$  into  $2j + 1$  equally spaced states of interstate spacing

$$\Delta E_{j=l-1/2} = \mu_B B \left( \frac{2l}{2l+1} \right). \quad (11.9.11)$$

In conclusion, in the presence of a weak external magnetic field, the two degenerate  $1S_{1/2}$  states of the hydrogen atom are split by  $2\mu_B B$ . Likewise, the four degenerate  $2S_{1/2}$  and  $2P_{1/2}$  states are split by  $(2/3)\mu_B B$ , whereas the four degenerate  $2P_{3/2}$  states are split by  $(4/3)\mu_B B$ . This is illustrated in Figure [fzee]. Note, finally, that because the  $\psi_{l,m_j}^{(2)}$  are not simultaneous eigenstates of the unperturbed and perturbing Hamiltonians, Equations ([e12.149]) and ([e12.150]) can only be regarded as the expectation values of the magnetic-field induced energy-shifts. However, as long as the external magnetic field is much weaker than the internal magnetic field, these expectation values are almost identical to the actual measured values of the energy-shifts.




Figure 24: The Zeeman effect for the  $n = 1$  and  $2$  states of a hydrogen atom. Here,  $\epsilon = \mu_B B$ . Not to scale.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled 11.9: Zeeman Effect is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 11.10: Hyperfine Structure

The proton in a hydrogen atom is a spin one-half charged particle, and therefore possesses a magnetic moment. By analogy with Equation ([11.10.58](#)), we can write

$$\mu_p = \frac{g_p e}{2 m_p} \mathbf{S}_p, \quad (11.10.1)$$

where  $\mu_p$  is the proton magnetic moment,  $\mathbf{S}_p$  is the proton spin, and the proton gyromagnetic ratio  $g_p$  is found experimentally to take the value 5.59. Note that the magnetic moment of a proton is much smaller (by a factor of order  $m_e/m_p$ ) than that of an electron. According to classical electromagnetism, the proton's magnetic moment generates a magnetic field of the form

$$\mathbf{B} = \frac{\mu_0}{4\pi r^3} [3 (\mu_p \cdot \mathbf{e}_r) \mathbf{e}_r - \mu_p] + \frac{2\mu_0}{3} \mu_p \delta^3(\mathbf{r}), \quad (11.10.2)$$

where  $\mathbf{e}_r = \mathbf{r}/r$ . We can understand the origin of the delta-function term in the previous expression by thinking of the proton as a tiny current loop centred on the origin. All magnetic field-lines generated by the loop must pass through the loop. Hence, if the size of the loop goes to zero then the field will be infinite at the origin, and this contribution is what is reflected by the delta-function term. Now, the Hamiltonian of the electron in the magnetic field generated by the proton is simply

$$H_1 = -\mu_e \cdot \mathbf{B}, \quad (11.10.3)$$

where

$$\mu_e = -\frac{e}{m_e} \mathbf{S}_e. \quad (11.10.4)$$

Here,  $\mu_e$  is the electron magnetic moment [see Equations ([11.10.58](#)) and ([11.10.59](#))], and  $\mathbf{S}_e$  the electron spin. Thus, the perturbing Hamiltonian is written

$$H_1 = \frac{\mu_0 g_p e^2}{8\pi m_p m_e} \frac{3 (\mathbf{S}_p \cdot \mathbf{e}_r) (\mathbf{S}_e \cdot \mathbf{e}_r) - \mathbf{S}_p \cdot \mathbf{S}_e}{r^3} + \frac{\mu_0 g_p e^2}{3 m_p m_e} \mathbf{S}_p \cdot \mathbf{S}_e \delta^3(\mathbf{r}). \quad (11.10.5)$$

Note that, because we have neglected coupling between the proton spin and the magnetic field generated by the electron's orbital motion, the previous expression is only valid for  $l = 0$  states.

According to standard first-order perturbation theory, the energy-shift induced by spin-spin coupling between the proton and the electron is the expectation value of the perturbing Hamiltonian. Hence,

$$\Delta E = \frac{\mu_0 g_p e^2}{8\pi m_p m_e} \left\langle \frac{3 (\mathbf{S}_p \cdot \mathbf{e}_r) (\mathbf{S}_e \cdot \mathbf{e}_r) - \mathbf{S}_p \cdot \mathbf{S}_e}{r^3} \right\rangle + \frac{\mu_0 g_p e^2}{3 m_p m_e} \langle \mathbf{S}_p \cdot \mathbf{S}_e \rangle |\psi(0)|^2. \quad (11.10.6)$$

For the ground-state of hydrogen, which is spherically symmetric, the first term in the previous expression vanishes by symmetry. Moreover, it is easily demonstrated that  $|\psi_{000}(0)|^2 = 1/(\pi a_0^3)$ . Thus, we obtain

$$\Delta E = \frac{\mu_0 g_p e^2}{3\pi m_p m_e a_0^3} \langle \mathbf{S}_p \cdot \mathbf{S}_e \rangle. \quad (11.10.7)$$

Let

$$\mathbf{S} = \mathbf{S}_e + \mathbf{S}_p \quad (11.10.8)$$

be the total spin. We can show that

$$\mathbf{S}_p \cdot \mathbf{S}_e = \frac{1}{2} (S^2 - S_e^2 - S_p^2). \quad (11.10.9)$$

Thus, the simultaneous eigenstates of the perturbing Hamiltonian and the main Hamiltonian are the simultaneous eigenstates of  $S_e^2$ ,  $S_p^2$ , and  $S^2$ . However, both the proton and the electron are spin one-half particles. According to Section [\[shalf\]](#), when two spin one-half particles are combined (in the absence of orbital angular momentum) the net state has either spin 1 or spin 0. In fact, there are three spin 1 states, known as triplet states, and a single spin 0 state, known as the singlet state. For all states, the eigenvalues of  $S_e^2$  and  $S_p^2$  are  $(3/4) \hbar^2$ . The eigenvalue of  $S^2$  is 0 for the singlet state, and  $2 \hbar^2$  for the triplet states. Hence,



$$\langle \mathbf{S}_p \cdot \mathbf{S}_e \rangle = -\frac{3}{4} \hbar^2 \quad (11.10.10)$$

for the singlet state, and

$$\langle \mathbf{S}_p \cdot \mathbf{S}_e \rangle = \frac{1}{4} \hbar^2 \quad (11.10.11)$$

for the triplet states.

It follows, from the previous analysis, that spin-spin coupling breaks the degeneracy of the two  $1S_{1/2}$  states in hydrogen, lifting the energy of the triplet configuration, and lowering that of the singlet. This splitting is known as *hyperfine structure*. The net energy difference between the singlet and the triplet states is

$$\Delta E = \frac{8}{3} g_p \frac{m_e}{m_p} \alpha^2 E_0 = 5.88 \times 10^{-6} \text{ eV}, \quad (11.10.12)$$

where  $E_0 = 13.6 \text{ eV}$  is the (magnitude of the) ground-state energy. Note that the hyperfine energy-shift is much smaller, by a factor  $m_e/m_p$ , than a typical fine structure energy-shift. If we convert the previous energy into a wavelength then we obtain

$$\lambda = 21.1 \text{ cm}. \quad (11.10.13)$$

This is the wavelength of the radiation emitted by a hydrogen atom which is collisionally excited from the singlet to the triplet state, and then decays back to the lower energy singlet state. The 21 cm line is famous in radio astronomy because it was used to map out the spiral structure of our galaxy in the 1950's .

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [11.10: Hyperfine Structure](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## CHAPTER OVERVIEW

### 12: Time-Dependent Perturbation Theory

Consider a system whose Hamiltonian can be written

$$H(t) = H_0 + H_1(t). \quad (12.1)$$

Here,  $H_0$  is again a simple time-independent Hamiltonian whose eigenvalues and eigenstates are known exactly. However,  $H_1$  now represents a small time-dependent external perturbation. Let the eigenstates of  $H_0$  take the form

$$H_0 \psi_m = E_m \psi_m. \quad (12.2)$$

We know (see Section [\[sstat\]](#)) that if the system is in one of these eigenstates then, in the absence of an external perturbation, it remains in this state for ever. However, the presence of a small time-dependent perturbation can, in principle, give rise to a finite probability that if the system is initially in some eigenstate  $\psi_n$  of the unperturbed Hamiltonian then it is found in some other eigenstate at a subsequent time (because  $\psi_n$  is no longer an exact eigenstate of the total Hamiltonian). In other words, a time-dependent perturbation allows the system to make transitions between its unperturbed energy eigenstates. Let us investigate such transitions.

- [12.1: Preliminary Analysis](#)
- [12.2: Two-State System](#)
- [12.3: Spin Magnetic Resonance](#)
- [12.4: Perturbation Expansion](#)
- [12.5: Harmonic Perturbation](#)
- [12.6: Electromagnetic Radiation](#)
- [12.7: Electric Dipole Approximation](#)
- [12.8: Spontaneous Emission](#)
- [12.9: Radiation from Harmonic Oscillator](#)
- [12.10: Selection Rules \(Hydrogen Atoms\)](#)
- [12.11: 2P-1S Transitions in Hydrogen](#)
- [12.12: Intensity Rules](#)
- [12.13: Forbidden Transitions](#)
- [12.E: Time-Dependent Perturbation Theory \(Exercises\)](#)

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [12: Time-Dependent Perturbation Theory](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 12.1: Preliminary Analysis

Suppose that at  $t = 0$  the state of the system is represented by

$$\psi(0) = \sum_m c_m \psi_m, \quad (12.1.1)$$

where the  $c_m$  are complex numbers. Thus, the initial state is some linear superposition of the unperturbed energy eigenstates. In the absence of the time-dependent perturbation, the time evolution of the system is simply (see Section [\[sstat\]](#))

$$\psi(t) = \sum_m c_m \exp(-i E_m t / \hbar) \psi_m. \quad (12.1.2)$$

Now, the probability of finding the system in state  $n$  at time  $t$  is

$$P_n(t) = |\langle \psi_n | \psi \rangle|^2 = |c_n \exp(-i E_n t / \hbar)|^2 = |c_n|^2 = P_n(0), \quad (12.1.3)$$

because the unperturbed eigenstates are assumed to be orthonormal: that is,

$$\langle n | m \rangle = \delta_{nm}. \quad (12.1.4)$$

Clearly, with  $H_1 = 0$ , the probability of finding the system in state  $\psi_n$  at time  $t$  is exactly the same as the probability of finding the system in this state at the initial time,  $t = 0$ . However, with  $H_1 \neq 0$ , we expect the  $P_n$ —and, hence, the  $c_n$ —to vary with time. Thus, we can write

$$\psi(t) = \sum_m c_m(t) \exp(-i E_m t / \hbar) \psi_m, \quad (12.1.5)$$

where  $P_n(t) = |c_n(t)|^2$ . Here, we have carefully separated the fast phase oscillation of the eigenstates, which depends on the unperturbed Hamiltonian, from the slow variation of the amplitudes  $c_n(t)$ , which depends entirely on the perturbation (i.e.,  $c_n$  is constant in time if  $H_1 = 0$ ). Note that in Equation ([\[e13.7\]](#)) the eigenstates  $\psi_m$  are time-independent (they are actually the eigenstates of  $H_0$  evaluated at the initial time,  $t = 0$ ).

The time-dependent Schrödinger equation [see Equation ([\[etimed\]](#))] yields

$$i \hbar \frac{\partial \psi(t)}{\partial t} = H(t) \psi(t) = [H_0 + H_1(t)] \psi(t). \quad (12.1.6)$$

Now, it follows from Equation ([\[e13.7\]](#)) that

$$(H_0 + H_1) \psi = \sum_m c_m \exp(-i E_m t / \hbar) (E_m + H_1) \psi_m. \quad (12.1.7)$$

We also have

$$i \hbar \frac{\partial \psi}{\partial t} = \sum_m \left( i \hbar \frac{dc_m}{dt} + c_m E_m \right) \exp(-i E_m t / \hbar) \psi_m, \quad (12.1.8)$$

because the  $\psi_m$  are time-independent. According to Equation ([\[e13.8\]](#)), we can equate the right-hand sides of the previous two equations to obtain

$$\sum_m i \hbar \frac{dc_m}{dt} \exp(-i E_m t / \hbar) \psi_m = \sum_m c_m \exp(-i E_m t / \hbar) H_1 \psi_m. \quad (12.1.9)$$

Projecting out the component of the previous equation which is proportional to  $\psi_n$ , using Equation ([\[e13.6\]](#)), we obtain

$$i \hbar \frac{dc_n(t)}{dt} = \sum_m H_{nm}(t) \exp(i \omega_{nm} t) c_m(t), \quad (12.1.10)$$

where

$$H_{nm}(t) = \langle n | H_1(t) | m \rangle, \quad (12.1.11)$$

and

$$\omega_{nm} = \frac{E_n - E_m}{\hbar}. \quad (12.1.12)$$

Suppose that there are  $N$  linearly independent eigenstates of the unperturbed Hamiltonian. According to Equations ([e13.12]), the time dependence of the set of  $N$  coefficients  $c_n$ , which specify the probabilities of finding the system in these eigenstates at time  $t$ , is determined by  $N$  coupled first-order differential equations. Note that Equations ([e13.12]) are exact—we have made no approximations at this stage. Unfortunately, we cannot generally find exact solutions to these equations. Instead, we have to obtain approximate solutions via suitable expansions in small quantities. However, for the particularly simple case of a two-state system (i.e.,  $N = 2$ ), it is actually possible to solve Equations ([e13.12]) without approximation. This solution is of great practical importance.

---

This page titled [12.1: Preliminary Analysis](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 12.2: Two-State System

Consider a system in which the time-independent Hamiltonian possesses two eigenstates, denoted

$$\begin{aligned} H_0 \psi_1 &= E_1 \psi_1, \\ H_0 \psi_2 &= E_2 \psi_2. \end{aligned}$$

Suppose, for the sake of simplicity, that the diagonal elements of the interaction Hamiltonian,  $H_1$ , are zero: that is,

$$\langle 1|H_1|1\rangle = \langle 2|H_1|2\rangle = 0. \quad (12.2.1)$$

The off-diagonal elements are assumed to oscillate sinusoidally at some frequency  $\omega$ : that is,

$$\langle 1|H_1|2\rangle = \langle 2|H_1|1\rangle^* = \gamma \hbar \exp(i\omega t), \quad (12.2.2)$$

where  $\gamma$  and  $\omega$  are real. Note that it is only the off-diagonal matrix elements which give rise to the effect which we are interested in: namely, transitions between states 1 and 2.

For a two-state system, Equation ([e13.12]) reduces to

$$\begin{aligned} i \frac{dc_1}{dt} &= \gamma \exp[i(\omega - \omega_{21})t] c_2, \\ i \frac{dc_2}{dt} &= \gamma \exp[-i(\omega - \omega_{21})t] c_1, \end{aligned}$$

where  $\omega_{21} = (E_2 - E_1)/\hbar$ . The previous two equations can be combined to give a second-order differential equation for the time-variation of the amplitude  $c_2$ : that is,

$$\frac{d^2 c_2}{dt^2} + i(\omega - \omega_{21}) \frac{dc_2}{dt} + \gamma^2 c_2 = 0. \quad (12.2.3)$$

Once we have solved for  $c_2$ , we can use Equation ([e13.20]) to obtain the amplitude  $c_1$ . Let us search for a solution in which the system is certain to be in state 1 (and, thus, has no chance of being in state 2) at time  $t = 0$ . Thus, our initial conditions are  $c_1(0) = 1$  and  $c_2(0) = 0$ . It is easily demonstrated that the appropriate solutions to ([e13.21]) and ([e13.20]) are

$$c_2(t) = \left( \frac{-i\gamma}{\Omega} \right) \exp\left[ \frac{-i(\omega - \omega_{21})t}{2} \right] \sin(\Omega t) \quad (12.2.4)$$

$$\begin{aligned} c_1(t) = & \exp\left[ \frac{i(\omega - \omega_{21})t}{2} \right] \cos(\Omega t) \\ & - \left[ \frac{i(\omega - \omega_{21})}{2\Omega} \right] \exp\left[ \frac{i(\omega - \omega_{21})t}{2} \right] \sin(\Omega t) \end{aligned} \quad (12.2.5)$$

where

$$\Omega = \sqrt{\gamma^2 + (\omega - \omega_{21})^2/4} \quad (12.2.6)$$

Now, the probability of finding the system in state 1 at time  $t$  is simply  $P_1(t) = |c_1(t)|^2$ . Likewise, the probability of finding the system in state 2 at time  $t$  is  $P_2(t) = |c_2(t)|^2$ . It follows that

$$\begin{aligned} P_1(t) &= 1 - P_2(t), \\ P_2(t) &= \left[ \frac{\gamma^2}{\gamma^2 + (\omega - \omega_{21})^2/4} \right] \sin^2(\Omega t). \end{aligned}$$

This result is known as *Rabi's formula*.

Equation ([e13.25]) exhibits all the features of a classic resonance. At resonance, when the oscillation frequency of the perturbation,  $\omega$ , matches the frequency  $\omega_{21}$ , we find that

$$\begin{aligned} P_1(t) &= \cos^2(\gamma t), \\ P_2(t) &= \sin^2(\gamma t). \end{aligned}$$

According to the previous result, the system starts off in state 1 at  $t = 0$ . After a time interval  $\pi/(2\gamma)$  it is certain to be in state 2. After a further time interval  $\pi/(2\gamma)$  it is certain to be in state 1 again, and so on. Thus, the system periodically flip-flops between states 1 and 2 under the influence of the time-dependent perturbation. This implies that the system alternatively absorbs and emits energy from the source of the perturbation.

The absorption-emission cycle also takes place away from the resonance, when  $\omega \neq \omega_{21}$ . However, the amplitude of the oscillation in the coefficient  $c_2$  is reduced. This means that the maximum value of  $P_2(t)$  is no longer unity, nor is the minimum of  $P_1(t)$  zero. In fact, if we plot the maximum value of  $P_2(t)$  as a function of the applied frequency,  $\omega$ , then we obtain a resonance curve whose maximum (unity) lies at the resonance, and whose full-width half-maximum (in frequency) is  $4\gamma$ . Thus, if the applied frequency differs from the resonant frequency by substantially more than  $2\gamma$  then the probability of the system jumping from state 1 to state 2 is always very small. In other words, the time-dependent perturbation is only effective at causing transitions between states 1 and 2 if its frequency of oscillation lies in the approximate range  $\omega_{21} \pm 2\gamma$ . Clearly, the weaker the perturbation (i.e., the smaller  $\gamma$  becomes), the narrower the resonance.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [12.2: Two-State System](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 12.3: Spin Magnetic Resonance

Consider a system consisting of a spin one-half particle with no orbital angular momentum (e.g., a bound electron) placed in a uniform  $z$ -directed magnetic field, and then subject to a small time-dependent magnetic field rotating in the  $x$ - $y$  plane at the angular frequency  $\omega$ . Thus,

$$\mathbf{B} = B_0 \mathbf{e}_z + B_1 [\cos(\omega t) \mathbf{e}_x + \sin(\omega t) \mathbf{e}_y], \quad (12.3.1)$$

where  $B_0$  and  $B_1$  are constants, with  $B_1 \ll B_0$ . The rotating magnetic field usually represents the magnetic component of an electromagnetic wave propagating along the  $z$ -axis. In this system, the electric component of the wave has no effect. The Hamiltonian is written

$$H = -\boldsymbol{\mu} \cdot \mathbf{B} = H_0 + H_1, \quad (12.3.2)$$

where

$$H_0 = -\frac{geB_0}{2m} S_z, \quad (12.3.3)$$

and

$$H_1 = -\frac{geB_1}{2m} [\cos(\omega t) S_x + \sin(\omega t) S_y]. \quad (12.3.4)$$

Here,  $g$  and  $m$  are the gyromagnetic ratio [see Equation (12.151)] and mass of the particle in question, respectively.

The eigenstates of the unperturbed Hamiltonian are the “spin up” and “spin down” states, denoted  $\chi_+$  and  $\chi_-$ , respectively. Of course, these states are the eigenstates of  $S_z$  corresponding to the eigenvalues  $+\hbar/2$  and  $-\hbar/2$  respectively. (See Section [sspin].) Thus, we have

$$H_0 \chi_{\pm} = \mp \frac{ge\hbar B_0}{4m} \chi_{\pm}. \quad (12.3.5)$$

The time-dependent Hamiltonian can be written

$$H_1 = -\frac{geB_1}{4m} [\exp(i\omega t) S_- + \exp(-i\omega t) S_+], \quad (12.3.6)$$

where  $S_+$  and  $S_-$  are the conventional raising and lowering operators for spin angular momentum. (See Section [sspin].) It follows that

$$\langle + | H_1 | + \rangle = \langle - | H_1 | - \rangle = 0, \quad (12.3.7)$$

and

$$\langle - | H_1 | + \rangle = \langle + | H_1 | - \rangle^* = -\frac{geB_1}{4m} \exp(i\omega t). \quad (12.3.8)$$

It can be seen that this system is exactly the same as the two-state system discussed in the previous section, provided that we make the following identifications:

$$\begin{aligned} \psi_1 &\rightarrow \chi_+, \\ \psi_2 &\rightarrow \chi_-, \\ \omega_{21} &\rightarrow \frac{geB_0}{2m}, \\ \gamma &\rightarrow -\frac{geB_1}{4m}. \end{aligned}$$

The resonant frequency,  $\omega_{21}$ , is simply the spin precession frequency in a uniform magnetic field of strength  $B_0$ . (See Section [sspin].) In the absence of the perturbation, the expectation values of  $S_x$  and  $S_y$  oscillate because of the spin precession, but the expectation value of  $S_z$  remains invariant. If we now apply a magnetic perturbation rotating at the resonant frequency then, according to the analysis of the previous section, the system undergoes a succession of spin flips,  $\chi_+ \leftrightarrow \chi_-$ , in addition to the spin precession. We also know that if the oscillation frequency of the applied field is very different from the resonant frequency then

there is virtually zero probability of the field triggering a spin flip. The width of the resonance (in frequency) is determined by the strength of the oscillating magnetic perturbation. Experimentalists are able to measure the gyromagnetic ratios of spin one-half particles to a high degree of accuracy by placing the particles in a uniform magnetic field of known strength, and then subjecting them to an oscillating magnetic field whose frequency is gradually scanned. By determining the resonant frequency (i.e., the frequency at which the particles absorb energy from the oscillating field), it is possible to determine the gyromagnetic ratio (assuming that the mass is known) .

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [12.3: Spin Magnetic Resonance](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).



## 12.4: Perturbation Expansion

Let us recall the analysis of Section 1.2. The  $\psi_n$  are the stationary orthonormal eigenstates of the time-independent unperturbed Hamiltonian,  $H_0$ . Thus,  $H_0 \psi_n = E_n \psi_n$ , where the  $E_n$  are the unperturbed energy levels, and  $\langle n|m \rangle = \delta_{nm}$ . Now, in the presence of a small time-dependent perturbation to the Hamiltonian,  $H_1(t)$ , the wavefunction of the system takes the form

$$\psi(t) = \sum_n c_n(t) \exp(-i \omega_n t) \psi_n, \quad (12.4.1)$$

where  $\omega_n = E_n/\hbar$ . The amplitudes  $c_n(t)$  satisfy

$$i \hbar \frac{dc_n}{dt} = \sum_m H_{nm} \exp(i \omega_{nm} t) c_m, \quad (12.4.2)$$

where  $H_{nm}(t) = \langle n|H_1(t)|m \rangle$  and  $\omega_{nm} = (E_n - E_m)/\hbar$ . Finally, the probability of finding the system in the  $n$ th eigenstate at time  $t$  is simply

$$P_n(t) = |c_n(t)|^2 \quad (12.4.3)$$

(assuming that, initially,  $\sum_n |c_n|^2 = 1$ ).

Suppose that at  $t = 0$  the system is in some initial energy eigenstate labeled  $i$ . Equation (12.4.2) is, thus, subject to the initial condition

$$c_n(0) = \delta_{ni}. \quad (12.4.4)$$

Let us attempt a perturbative solution of Equation (12.4.2) using the ratio of  $H_1$  to  $H_0$  (or  $H_{nm}$  to  $\hbar \omega_{nm}$ , to be more exact) as our expansion parameter. Now, according to Equation (12.4.2), the  $c_n$  are constant in time in the absence of the perturbation. Hence, the zeroth-order solution is simply

$$c_n^{(0)}(t) = \delta_{ni}. \quad (12.4.5)$$

The first-order solution is obtained, via iteration, by substituting the zeroth-order solution into the right-hand side of Equation (12.4.2). Thus, we obtain

$$i \hbar \frac{dc_n^{(1)}}{dt} = \sum_m H_{nm} \exp(i \omega_{nm} t) c_m^{(0)} = H_{ni} \exp(i \omega_{ni} t), \quad (12.4.6)$$

subject to the boundary condition  $c_n^{(1)}(0) = 0$ . The solution to the previous equation is

$$c_n^{(1)} = -\frac{i}{\hbar} \int_0^t H_{ni}(t') \exp(i \omega_{ni} t') dt'. \quad (12.4.7)$$

It follows that, up to first-order in our perturbation expansion,

$$c_n(t) = \delta_{ni} - \frac{i}{\hbar} \int_0^t H_{ni}(t') \exp(i \omega_{ni} t') dt'. \quad (12.4.8)$$

Hence, the probability of finding the system in some final energy eigenstate labeled  $f$  at time  $t$ , given that it is definitely in a different initial energy eigenstate labeled  $i$  at time  $t = 0$ , is

$$P_{i \rightarrow f}(t) = |c_f(t)|^2 = \left| -\frac{i}{\hbar} \int_0^t H_{fi}(t') \exp(i \omega_{fi} t') dt' \right|^2. \quad (12.4.9)$$

Note, finally, that our perturbative solution is clearly only valid provided

$$P_{i \rightarrow f}(t) \ll 1. \quad (12.4.10)$$

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [12.4: Perturbation Expansion](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 12.5: Harmonic Perturbation

Consider a (Hermitian) perturbation that oscillates sinusoidally in time. This is usually termed a *harmonic perturbation*. Such a perturbation takes the form

$$H_1(t) = V \exp(i \omega t) + V^\dagger \exp(-i \omega t), \quad (12.5.1)$$

where  $V$  is, in general, a function of position, momentum, and spin operators.

It follows from Equations ([e13.48]) and ([e13.51]) that, to first-order,

$$c_f(t) = -\frac{i}{\hbar} \int_0^t \left[ V_{fi} \exp(i \omega t') + V_{fi}^\dagger \exp(-i \omega t') \right] \exp(i \omega_{fi} t') dt', \quad (12.5.2)$$

where

$$V_{fi} = \langle f | V | i \rangle, \\ V_{fi}^\dagger = \langle f | V^\dagger | i \rangle = \langle i | V | f \rangle^*.$$

Integration with respect to  $t'$  yields

$$c_f(t) = -\frac{i t}{\hbar} \left( V_{fi} \exp[i(\omega + \omega_{fi}) t/2] \text{sinc}[(\omega + \omega_{fi}) t/2] \right. \\ \left. + V_{fi}^\dagger \exp[-i(\omega - \omega_{fi}) t/2] \text{sinc}[(\omega - \omega_{fi}) t/2] \right),$$

where

$$\text{sinc } x \equiv \frac{\sin x}{x}. \quad (12.5.3)$$




Figure 25: The functions  $\text{sinc}(x)$  (dashed curve) and  $\text{sinc}^2(x)$  (solid curve). The vertical dotted lines denote the region  $|x| \leq \pi$

Now, the function  $\text{sinc}(x)$  takes its largest values when  $|x| \lesssim \pi$ , and is fairly negligible when  $|x| \gg \pi$ . (See Figure [fsinc].) Thus, the first and second terms on the right-hand side of Equation ([e13.55]) are only non-negligible when

$$|\omega + \omega_{fi}| \lesssim \frac{2\pi}{t} \quad (12.5.4)$$

and

$$|\omega - \omega_{fi}| \lesssim \frac{2\pi}{t} \quad (12.5.5)$$

respectively.

Clearly, as  $t$  increases, the ranges in  $\omega$  over which these two terms are non-negligible gradually shrink in size. Eventually, when  $t \gg 2\pi/|\omega_{fi}|$ , these two ranges become strongly non-overlapping. Hence, in this limit,  $P_{i \rightarrow f} = |c_f|^2$  yields

$$P_{i \rightarrow f}(t) = \frac{t^2}{\hbar^2} \left\{ |V_{fi}|^2 \text{sinc}^2[(\omega + \omega_{fi})t/2] + |V_{fi}^\dagger|^2 \text{sinc}^2[(\omega - \omega_{fi})t/2] \right\}. \quad (12.5.6)$$

Now, the function  $\text{sinc}^2(x)$  is very strongly peaked at  $x = 0$ , and is completely negligible for  $|x| \gg \pi$ . (See Figure [\[fsinc\]](#).) It follows that the previous expression exhibits a resonant response to the applied perturbation at the frequencies  $\omega = \pm\omega_{fi}$ . Moreover, the widths of these resonances decrease linearly as time increases. At each of the resonances (i.e., at  $\omega = \pm\omega_{fi}$ ), the transition probability  $P_{i \rightarrow f}(t)$  varies as  $t^2$  [because  $\text{sinc}(0) = 1$ ]. This behavior is entirely consistent with our earlier result ([\[e13.28\]](#)), for the two-state system, in the limit  $\gamma t \ll 1$  (recall that our perturbative solution is only valid as long as  $P_{i \rightarrow f} \ll 1$ ).

The resonance at  $\omega = -\omega_{fi}$  corresponds to

$$E_f - E_i = -\hbar\omega. \quad (12.5.7)$$

This implies that the system loses energy  $\hbar\omega$  to the perturbing field, while making a transition to a final state whose energy is less than the initial state by  $\hbar\omega$ . This process is known as *stimulated emission*. The resonance at  $\omega = \omega_{fi}$  corresponds to

$$E_f - E_i = \hbar\omega. \quad (12.5.8)$$

This implies that the system gains energy  $\hbar\omega$  from the perturbing field, while making a transition to a final state whose energy is greater than that of the initial state by  $\hbar\omega$ . This process is known as *absorption*.

Stimulated emission and absorption are mutually exclusive processes, because the first requires  $\omega_{fi} < 0$ , whereas the second requires  $\omega_{fi} > 0$ . Hence, we can write the transition probabilities for both processes separately. Thus, from Equation ([\[e13.49\]](#)), the transition probability for stimulated emission is

$$P_{i \rightarrow f}^{stm}(t) = \frac{t^2}{\hbar^2} |V_{if}^\dagger|^2 \text{sinc}^2[(\omega - \omega_{if})t/2], \quad (12.5.9)$$

where we have made use of the facts that  $\omega_{if} = -\omega_{fi} > 0$ , and  $|V_{fi}|^2 = |V_{if}^\dagger|^2$ . Likewise, the transition probability for absorption is

$$P_{i \rightarrow f}^{abs}(t) = \frac{t^2}{\hbar^2} |V_{fi}^\dagger|^2 \text{sinc}^2[(\omega - \omega_{fi})t/2]. \quad (12.5.10)$$

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [12.5: Harmonic Perturbation](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 12.6: Electromagnetic Radiation

Let us use the previous results to investigate the interaction of an atomic electron with classical (i.e., non-quantized) electromagnetic radiation.

The unperturbed Hamiltonian of the system is

$$H_0 = \frac{p^2}{2m_e} + V_0(r). \quad (12.6.1)$$

Now, the standard classical prescription for obtaining the Hamiltonian of a particle of charge  $q$  in the presence of an electromagnetic field is

$$\begin{aligned} \mathbf{p} &\rightarrow \mathbf{p} + q \mathbf{A}, \\ H &\rightarrow H - q \phi, \end{aligned}$$

where  $\mathbf{A}(\mathbf{r})$  is the vector potential, and  $\phi(\mathbf{r})$  the scalar potential. Note that

$$\begin{aligned} \mathbf{E} &= -\nabla\phi - \frac{\partial \mathbf{A}}{\partial t}, \\ \mathbf{B} &= \nabla \times \mathbf{A}. \end{aligned}$$

This prescription also works in quantum mechanics. Thus, the Hamiltonian of an atomic electron placed in an electromagnetic field is

$$H = \frac{(\mathbf{p} - e \mathbf{A})^2}{2m_e} + e\phi + V_0(r), \quad (12.6.2)$$

where  $\mathbf{A}$  and  $\phi$  are functions of the position operators. The previous equation can be written

$$H = \frac{(p^2 - e \mathbf{A} \cdot \mathbf{p} - e \mathbf{p} \cdot \mathbf{A} + e^2 A^2)}{2m_e} + e\phi + V_0(r). \quad (12.6.3)$$

Now,

$$\mathbf{p} \cdot \mathbf{A} = \mathbf{A} \cdot \mathbf{p}, \quad (12.6.4)$$

provided that we adopt the Coulomb gauge  $\nabla \cdot \mathbf{A} = 0$ . Hence,

$$H = \frac{p^2}{2m_e} - \frac{e \mathbf{A} \cdot \mathbf{p}}{m_e} + \frac{e^2 A^2}{2m_e} + e\phi + V_0(r). \quad (12.6.5)$$

Suppose that the perturbation corresponds to a linearly polarized, monochromatic, plane-wave. In this case,

$$\begin{aligned} \phi &= 0, \\ \mathbf{A} &= A_0 \epsilon \cos(\mathbf{k} \cdot \mathbf{r} - \omega t), \end{aligned}$$

where  $\mathbf{k}$  is the wavevector (note that  $\omega = kc$ ), and  $\epsilon$  a unit vector that specifies the direction of polarization (i.e., the direction of  $\mathbf{E}$ ). Note that  $\epsilon \cdot \mathbf{k} = 0$ . The Hamiltonian becomes

$$H = H_0 + H_1(t), \quad (12.6.6)$$

with

$$H_0 = \frac{p^2}{2m_e} + V_0(r), \quad (12.6.7)$$

and

$$H_1 \simeq -\frac{e \mathbf{A} \cdot \mathbf{p}}{m_e}, \quad (12.6.8)$$

where the  $A^2$  term, which is second order in  $A_0$ , has been neglected.

The perturbing Hamiltonian can be written

$$H_1 = -\frac{e A_0 \epsilon \cdot \mathbf{p}}{2 m_e} [\exp(i \mathbf{k} \cdot \mathbf{r} - i \omega t) + \exp(-i \mathbf{k} \cdot \mathbf{r} + i \omega t)]. \quad (12.6.9)$$

This has the same form as Equation (12.5.51), provided that

$$V^\dagger = -\frac{e A_0 \epsilon \cdot \mathbf{p}}{2 m_e} \exp(i \mathbf{k} \cdot \mathbf{r}). \quad (12.6.10)$$

It follows from Equations (12.5.53), (12.5.63), and (12.5.79) that the transition probability for radiation induced absorption is

$$P_{i \rightarrow f}^{abs}(t) = \frac{t^2}{\hbar^2} \frac{e^2 |A_0|^2}{4 m_e^2} |\langle f | \epsilon \cdot \mathbf{p} \exp(i \mathbf{k} \cdot \mathbf{r}) | i \rangle|^2 \text{sinc}^2[(\omega - \omega_{fi}) t / 2]. \quad (12.6.11)$$

Now, the mean energy density of an electromagnetic wave is

$$u = \frac{1}{2} \left( \frac{\epsilon_0 |E_0|^2}{2} + \frac{|B_0|^2}{2 \mu_0} \right) = \frac{1}{2} \epsilon_0 |E_0|^2, \quad (12.6.12)$$

where  $E_0 = A_0 \omega$  and  $B_0 = E_0 / c$  are the peak electric and magnetic field-strengths, respectively. It thus follows that

$$P_{i \rightarrow f}^{abs}(t) = \frac{t^2 e^2}{2 \epsilon_0 \hbar^2 m_e^2 \omega^2} |\langle f | \epsilon \cdot \mathbf{p} \exp(i \mathbf{k} \cdot \mathbf{r}) | i \rangle|^2 u \text{sinc}^2[(\omega - \omega_{fi}) t / 2]. \quad (12.6.13)$$

Thus, not surprisingly, the transition probability for radiation induced absorption (or stimulated emission) is directly proportional to the energy density of the incident radiation.

Suppose that the incident radiation is not monochromatic, but instead extends over a range of frequencies. We can write

$$u = \int_{-\infty}^{\infty} \rho(\omega) d\omega, \quad (12.6.14)$$

where  $\rho(\omega) d\omega$  is the energy density of radiation whose frequencies lie between  $\omega$  and  $\omega + d\omega$ . Equation (12.5.80) generalizes to

$$P_{i \rightarrow f}^{abs}(t) = \int_{-\infty}^{\infty} \frac{t^2 e^2}{2 \epsilon_0 \hbar^2 m_e^2 \omega^2} |\langle f | \epsilon \cdot \mathbf{p} \exp(i \mathbf{k} \cdot \mathbf{r}) | i \rangle|^2 \rho(\omega) \text{sinc}^2[(\omega - \omega_{fi}) t / 2] d\omega. \quad (12.6.15)$$

Note, however, that the previous expression is only valid provided the radiation in question is *incoherent*: that is, provided there are no phase correlations between waves of different frequencies. This follows because it is permissible to add the intensities of incoherent radiation, whereas we must always add the amplitudes of coherent radiation. Given that the function  $\text{sinc}^2[(\omega - \omega_{fi}) t / 2]$  is very strongly peaked (see Figure 12.5.1) about  $\omega = \omega_{fi}$  (assuming that  $t \gg 2\pi / \omega_{fi}$ ), and

$$\int_{-\infty}^{\infty} \text{sinc}^2(x) dx = \pi, \quad (12.6.16)$$

the previous equation reduces to

$$P_{i \rightarrow f}^{abs}(t) = \frac{\pi e^2 \rho(\omega_{fi})}{\epsilon_0 \hbar^2 m_e^2 \omega_{fi}^2} |\langle f | \epsilon \cdot \mathbf{p} \exp(i \mathbf{k} \cdot \mathbf{r}) | i \rangle|^2 t. \quad (12.6.17)$$

Note that in integrating over the frequencies of the incoherent radiation we have transformed a transition probability that is basically proportional to  $t^2$  [see Equation (12.5.80)] to one that is proportional to  $t$ . As has already been explained, the previous expression is only valid when  $P_{i \rightarrow f}^{abs} \ll 1$ . However, the result that

$$w_{i \rightarrow f}^{abs} \equiv \frac{dP_{i \rightarrow f}^{abs}}{dt} = \frac{\pi e^2 \rho(\omega_{fi})}{\epsilon_0 \hbar^2 m_e^2 \omega_{fi}^2} |\langle f | \epsilon \cdot \mathbf{p} \exp(i \mathbf{k} \cdot \mathbf{r}) | i \rangle|^2 \quad (12.6.18)$$

is constant in time is universally valid. Here,  $w_{i \rightarrow f}^{abs}$  is the transition probability per unit time interval, otherwise known as the *transition rate*. Given that the transition rate is constant, we can write (see Chapter 12.2)

$$P_{i \rightarrow f}^{abs}(t + dt) - P_{i \rightarrow f}^{abs}(t) = [1 - P_{i \rightarrow f}^{abs}(t)] w_{i \rightarrow f}^{abs} dt : \quad (12.6.19)$$

that is, the probability that the system makes a transition from state  $i$  to state  $f$  between times  $t$  and  $t + dt$  is equivalent to the probability that the system does not make a transition between times 0 and  $t$  and then makes a transition in a time interval  $dt$ —the probabilities of these two events are  $1 - P_{i \rightarrow f}^{abs}(t)$  and  $w_{i \rightarrow f}^{abs} dt$ , respectively. It follows that

$$\frac{dP_{i \rightarrow f}^{abs}}{dt} + w_{i \rightarrow f}^{abs} P_{i \rightarrow f}^{abs} = w_{i \rightarrow f}^{abs}, \quad (12.6.20)$$

with the initial condition  $P_{i \rightarrow f}^{abs}(0) = 0$ . The previous equation can be solved to give

$$P_{i \rightarrow f}^{abs}(t) = 1 - \exp(-w_{i \rightarrow f}^{abs} t). \quad (12.6.21)$$

This result is consistent with Equation ([e13.86]) provided  $w_{i \rightarrow f}^{abs} t \ll 1$ : that is, provided that  $P_{i \rightarrow f}^{abs} \ll 1$ .

Using similar arguments to those given previously, the transition probability for stimulated emission can be shown to take the form

$$P_{i \rightarrow f}^{stm}(t) = 1 - \exp(-w_{i \rightarrow f}^{stm} t), \quad (12.6.22)$$

where the corresponding transition rate is written

$$w_{i \rightarrow f}^{stm} = \frac{\pi e^2 \rho(\omega_{if})}{\epsilon_0 \hbar^2 m_e^2 \omega_{if}^2} |\langle i | \epsilon \cdot \mathbf{p} \exp(i \mathbf{k} \cdot \mathbf{r}) | f \rangle|^2. \quad (12.6.23)$$

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [12.6: Electromagnetic Radiation](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 12.7: Electric Dipole Approximation

In general, the wavelength of the type of electromagnetic radiation that induces, or is emitted during, transitions between different atomic energy levels is much larger than the typical size of an atom. Thus,

$$\exp(i \mathbf{k} \cdot \mathbf{r}) = 1 + i \mathbf{k} \cdot \mathbf{r} + \dots, \quad (12.7.1)$$

can be approximated by its first term, unity. This approach is known as the *electric dipole approximation*. It follows that

$$\langle f | \epsilon \cdot \mathbf{p} \exp(i \mathbf{k} \cdot \mathbf{r}) | i \rangle \simeq \epsilon \cdot \langle f | \mathbf{p} | i \rangle. \quad (12.7.2)$$

Now, it is readily demonstrated that

$$[\mathbf{r}, H_0] = \frac{i \hbar \mathbf{p}}{m_e} \quad (12.7.3)$$

so

$$\langle f | \mathbf{p} | i \rangle = -i \frac{m_e}{\hbar} \langle f | [\mathbf{r}, H_0] | i \rangle = i m_e \omega_{fi} \langle f | \mathbf{r} | i \rangle. \quad (12.7.4)$$

Thus, our previous expressions for the transition rates for radiation induced absorption and stimulated emission reduce to

$$w_{i \rightarrow f}^{abs} = \frac{\pi}{\epsilon_0 \hbar^2} |\epsilon \cdot \mathbf{d}_{if}|^2 \rho(\omega_{fi}),$$

$$w_{i \rightarrow f}^{stm} = \frac{\pi}{\epsilon_0 \hbar^2} |\epsilon \cdot \mathbf{d}_{if}|^2 \rho(\omega_{if}),$$

respectively. Here,

$$\mathbf{d}_{if} = \langle f | e \mathbf{r} | i \rangle \quad (12.7.5)$$

is the effective electric dipole moment of the atom when making a transition from state  $i$  to state  $f$ .

Equations ([e13.97]) and ([e13.98]) give the transition rates for absorption and stimulated emission, respectively, induced by a linearly polarized plane-wave. Actually, we are more interested in the transition rates induced by unpolarized isotropic radiation. To obtain these we must average Equations ([e13.97]) and ([e13.98]) over all possible polarizations and propagation directions of the wave. To facilitate this process, we can define a set of Cartesian coordinates such that the wavevector  $\mathbf{k}$ , which specifies the direction of wave propagation, points along the  $z$ -axis, and the vector  $\mathbf{d}_{if}$ , which specifies the direction of the atomic dipole moment, lies in the  $x$ - $z$  plane. It follows that the vector  $\epsilon$ , which specifies the direction of wave polarization, must lie in the  $x$ - $y$  plane, because it has to be orthogonal to  $\mathbf{k}$ . Thus, we can write

$$\begin{aligned} \mathbf{k} &= (0, 0, k), \\ \mathbf{d}_{if} &= (d_{if} \sin \theta, 0, d_{if} \cos \theta), \\ \epsilon &= (\cos \phi, \sin \phi, 0), \end{aligned}$$

which implies that

$$|\epsilon \cdot \mathbf{d}_{if}|^2 = d_{if}^2 \sin^2 \theta \cos^2 \phi. \quad (12.7.6)$$

We must now average the previous quantity over all possible values of  $\theta$  and  $\phi$ . Thus,

$$\langle |\epsilon \cdot \mathbf{d}_{if}|^2 \rangle_{av} = d_{if}^2 \frac{\int \int \sin^2 \theta \cos^2 \phi d\Omega}{4\pi}, \quad (12.7.7)$$

where  $d\Omega = \sin \theta d\theta d\phi$ , and the integral is taken over all solid angle. It is easily demonstrated that

$$\langle |\epsilon \cdot \mathbf{d}_{if}|^2 \rangle_{av} = \frac{d_{if}^2}{3}. \quad (12.7.8)$$

Here,  $d_{if}^2$  stands for

$$d_{if}^2 = |\langle f | e x | i \rangle|^2 + |\langle f | e y | i \rangle|^2 + |\langle f | e z | i \rangle|^2. \quad (12.7.9)$$



Hence, the transition rates for absorption and stimulated emission induced by unpolarized isotropic radiation are

$$w_{i \rightarrow f}^{abs} = \frac{\pi}{3 \epsilon_0 \hbar^2} d_{if}^2 \rho(\omega_{fi}),$$
$$w_{i \rightarrow f}^{stm} = \frac{\pi}{3 \epsilon_0 \hbar^2} d_{if}^2 \rho(\omega_{if}),$$

respectively.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [12.7: Electric Dipole Approximation](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 12.8: Spontaneous Emission

So far, we have calculated the rates of radiation induced transitions between two atomic states. This process is known as absorption when the energy of the final state exceeds that of the initial state, and stimulated emission when the energy of the final state is less than that of the initial state. Now, in the absence of any external radiation, we would not expect an atom in a given state to spontaneously jump into an state with a higher energy. On the other hand, it should be possible for such an atom to spontaneously jump into an state with a lower energy via the emission of a photon whose energy is equal to the difference between the energies of the initial and final states. This process is known as *spontaneous emission*.

It is possible to derive the rate of spontaneous emission between two atomic states from a knowledge of the corresponding absorption and stimulated emission rates using a famous thermodynamic argument due to Einstein . Consider a very large ensemble of similar atoms placed inside a closed cavity whose walls (which are assumed to be perfect emitters and absorbers of radiation) are held at the constant temperature  $T$ . Let the system have attained thermal equilibrium. According to statistical thermodynamics, the cavity is filled with so-called “black-body” electromagnetic radiation whose energy spectrum is

$$\rho(\omega) = \frac{\hbar}{\pi^2 c^3} \frac{\omega^3}{\exp(\hbar\omega/k_B T) - 1}, \quad (12.8.1)$$

where  $k_B$  is the Boltzmann constant. This well-known result was first obtained by Max Planck in 1900 .

Consider two atomic states, labeled  $i$  and  $f$ , with  $E_i > E_f$ . One of the tenants of statistical thermodynamics is that in thermal equilibrium we have so-called *detailed balance* . This means that, irrespective of any other atomic states, the rate at which atoms in the ensemble leave state  $i$  due to transitions to state  $f$  is exactly balanced by the rate at which atoms enter state  $i$  due to transitions from state  $f$ . The former rate (i.e., number of transitions per unit time in the ensemble) is written

$$W_{i \rightarrow f} = N_i (w_{i \rightarrow f}^{spn} + w_{i \rightarrow f}^{stm}), \quad (12.8.2)$$

where  $w_{i \rightarrow f}^{spn}$  is the rate of spontaneous emission (for a single atom) between states  $i$  and  $f$ , and  $N_i$  is the number of atoms in the ensemble in state  $i$ . Likewise, the latter rate takes the form

$$W_{f \rightarrow i} = N_f w_{f \rightarrow i}^{abs}, \quad (12.8.3)$$

where  $N_f$  is the number of atoms in the ensemble in state  $f$ . The previous expressions describe how atoms in the ensemble make transitions from state  $i$  to state  $f$  due to a combination of spontaneous and stimulated emission, and make the opposite transition as a consequence of absorption. In thermal equilibrium, we have  $W_{i \rightarrow f} = W_{f \rightarrow i}$ , which gives

$$w_{i \rightarrow f}^{spn} = \frac{N_f}{N_i} w_{f \rightarrow i}^{abs} - w_{i \rightarrow f}^{stm}. \quad (12.8.4)$$

According to Equations ([e13.107]) and ([e13.108]), we can also write

$$w_{i \rightarrow f}^{spn} = \left( \frac{N_f}{N_i} - 1 \right) \frac{\pi}{3 \epsilon_0 \hbar^2} d_{if}^2 \rho(\omega_{if}). \quad (12.8.5)$$

Now, another famous result in statistical thermodynamics is that in thermal equilibrium the number of atoms in an ensemble occupying a state of energy  $E$  is proportional to  $\exp(-E/k_B T)$ . This implies that

$$\frac{N_f}{N_i} = \frac{\exp(-E_f/k_B T)}{\exp(-E_i/k_B T)} = \exp(\hbar\omega_{if}/k_B T). \quad (12.8.6)$$

Thus, it follows from Equation ([e13.109]), ([e13.113]), and ([e13.114]) that the rate of spontaneous emission between states  $i$  and  $f$  takes the form

$$w_{i \rightarrow f}^{spn} = \frac{\omega_{if}^3 d_{if}^2}{3\pi \epsilon_0 \hbar c^3}. \quad (12.8.7)$$

Note, that, although the previous result has been derived for an atom in a radiation-filled cavity, it remains correct even in the absence of radiation. Finally, the corresponding absorption and stimulated emission rates for an atom in a radiation-filled cavity are

$$w_{i \rightarrow f}^{abs} = \frac{\omega_{fi}^3 d_{if}^2}{3\pi \epsilon_0 \hbar c^3} \frac{1}{\exp(\hbar \omega_{fi}/k_B T) - 1},$$

$$w_{i \rightarrow f}^{stm} = \frac{\omega_{if}^3 d_{if}^2}{3\pi \epsilon_0 \hbar c^3} \frac{1}{\exp(\hbar \omega_{if}/k_B T) - 1},$$

respectively.

Let us estimate the typical value of the spontaneous emission rate for a hydrogen atom. We expect the dipole moment  $d_{if}$  to be of order  $e a_0$ , where  $a_0$  is the Bohr radius. [See Equation ([\[e9.57\]](#)).] We also expect  $\omega_{if}$  to be of order  $|E_0|/\hbar$ , where  $E_0$  is the energy of the ground-state [see Equation ([\[e9.56\]](#))]. It thus follows from Equation ([\[e3.115\]](#)) that

$$w_{i \rightarrow f}^{spn} \sim \alpha^3 \omega_{if}, \quad (12.8.8)$$

where  $\alpha = e^2/(4\pi \epsilon_0 \hbar c) \simeq 1/137$  is the fine-structure constant. This is an important result, because our perturbation expansion is based on the assumption that the transition rate between different energy eigenstates is much slower than the frequency of phase oscillation of these states: that is, that  $w_{i \rightarrow f}^{spn} \ll \omega_{if}$ . (See Section [1.2.](#)) This is indeed the case.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [12.8: Spontaneous Emission](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 12.9: Radiation from Harmonic Oscillator

Consider an electron in a one-dimensional harmonic oscillator potential aligned along the  $x$ -axis. According to Section [sosc], the unperturbed energy eigenvalues of the system are

$$E_n = (n + 1/2) \hbar \omega_0, \quad (12.9.1)$$

where  $\omega_0$  is the frequency of the corresponding classical oscillator. Here, the quantum number  $n$  takes the values  $0, 1, 2, \dots$ . Let the  $\psi_n(x)$  be the (real) properly normalized unperturbed eigenstates of the system.

Suppose that the electron is initially in an excited state: that is,  $n > 0$ . In principle, the electron can decay to a lower energy state via the spontaneous emission of a photon of the appropriate frequency. Let us investigate this effect. Now, according to Equation ([e3.115]), the system can only make a spontaneous transition from an energy state corresponding to the quantum number  $n$  to one corresponding to the quantum number  $n'$  if the associated electric dipole moment

$$(d_x)_{n,n'} = \langle n | e x | n' \rangle = e \int_{-\infty}^{\infty} \psi_n(x) x \psi_{n'}(x) dx \quad (12.9.2)$$

is non-zero [because  $d_{if} \equiv (d_x)_{n,n'}$  for the case in hand]. However, according to Equation ([e5.xxx]),

$$\int_{-\infty}^{\infty} \psi_n x \psi_{n'} dx = \sqrt{\frac{\hbar}{2 m_e \omega_0}} \left( \sqrt{n} \delta_{n,n'+1} + \sqrt{n'} \delta_{n,n'-1} \right). \quad (12.9.3)$$

Because we are dealing with emission, we must have  $n > n'$ . Hence, we obtain

$$(d_x)_{n,n'} = e \sqrt{\frac{\hbar n}{2 m_e \omega_0}} \delta_{n,n'+1}. \quad (12.9.4)$$

It is clear that (in the electric dipole approximation) we can only have spontaneous emission between states whose quantum numbers differ by unity. Thus, the frequency of the photon emitted when the  $n$ th excited state decays is

$$\omega_{n,n-1} = \frac{E_n - E_{n-1}}{\hbar} = \omega_0. \quad (12.9.5)$$

Hence, we conclude that, no matter which state decays, the emitted photon always has the same frequency as the classical oscillator.

According to Equation ([e3.115]), the decay rate of the  $n$ th excited state is given by

$$w_n = \frac{\omega_{n,n-1}^3 (d_x)_{n,n-1}^2}{3\pi \epsilon_0 \hbar c^3}. \quad (12.9.6)$$

It follows that

$$w_n = \frac{n e^2 \omega_0^2}{6\pi \epsilon_0 m_e c^3}. \quad (12.9.7)$$

The mean radiated power is simply

$$P_n = \hbar \omega_0 w_n = \frac{e^2 \omega_0^2}{6\pi \epsilon_0 m_e c^3} [E_n - (1/2) \hbar \omega_0]. \quad (12.9.8)$$

Classically, an electron in a one-dimensional oscillator potential radiates at the oscillation frequency  $\omega_0$  with the mean power

$$P = \frac{e^2 \omega_0^2}{6\pi \epsilon_0 m_e c^3} E, \quad (12.9.9)$$

where  $E$  is the oscillator energy. It can be seen that a quantum oscillator radiates in an almost exactly analogous manner to the equivalent classical oscillator. The only difference is the factor  $(1/2) \hbar \omega_0$  in Equation ([e13.126])—this is needed to ensure that the ground-state of the quantum oscillator does not radiate.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [12.9: Radiation from Harmonic Oscillator](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 12.10: Selection Rules (Hydrogen Atoms)

Let us now consider spontaneous transitions between the different energy levels of a hydrogen atom. Because the perturbing Hamiltonian ([e13.77]) does not contain any spin operators, we can neglect electron spin in our analysis. Thus, according to Section [s10.4], the various energy eigenstates of the hydrogen atom are labeled by the familiar quantum numbers  $n$ ,  $l$ , and  $m$ .

According to Equations ([e3.106]) and ([e3.115]), a hydrogen atom can only make a spontaneous transition from an energy state corresponding to the quantum numbers  $n, l, m$  to one corresponding to the quantum numbers  $n', l', m'$  if the modulus squared of the associated electric dipole moment

$$d^2 = |\langle n, l, m | e x | n', l', m' \rangle|^2 + |\langle n, l, m | e y | n', l', m' \rangle|^2 + |\langle n, l, m | e z | n', l', m' \rangle|^2 \quad (12.10.1)$$

is non-zero. Now, we have already seen, in Section [s12.5], that the matrix element  $\langle n, l, m | z | n', l', m' \rangle$  is only non-zero provided that  $m' = m$  and  $l' = l \pm 1$ . It turns out that the proof that this matrix element is zero unless  $l' = l \pm 1$  can, via a trivial modification, also be used to demonstrate that  $\langle n, l, m | x | n', l', m' \rangle$  and  $\langle n, l, m | y | n', l', m' \rangle$  are also zero unless  $l' = l \pm 1$ . Consider

$$x_{\pm} = x \pm i y. \quad (12.10.2)$$

It is easily demonstrated that

$$[L_z, x_{\pm}] = \pm \hbar x_{\pm}. \quad (12.10.3)$$

Hence,

$$\langle n, l, m | [L_z, x_+] - \hbar x_+ | n', l', m' \rangle = \hbar (m - m' - 1) \langle n, l, m | x_+ | n', l', m' \rangle = 0, \quad (12.10.4)$$

and

$$\langle n, l, m | [L_z, x_-] + \hbar x_- | n', l', m' \rangle = \hbar (m - m' + 1) \langle n, l, m | x_- | n', l', m' \rangle = 0. \quad (12.10.5)$$

Clearly,  $\langle n, l, m | x_+ | n', l', m' \rangle$  is zero unless  $m' = m - 1$ , and  $\langle n, l, m | x_- | n', l', m' \rangle$  is zero unless  $m' = m + 1$ . Now,  $\langle n, l, m | x | n', l', m' \rangle$  and  $\langle n, l, m | y | n', l', m' \rangle$  are obviously both zero if  $\langle n, l, m | x_+ | n', l', m' \rangle$  and  $\langle n, l, m | x_- | n', l', m' \rangle$  are both zero. Hence, we conclude that  $\langle n, l, m | x | n', l', m' \rangle$  and  $\langle n, l, m | y | n', l', m' \rangle$  are only non-zero if  $m' = m \pm 1$ .

The previous arguments demonstrate that spontaneous transitions between different energy levels of a hydrogen atom are only possible provided

$$l' = l \pm 1, \\ m' = m, m \pm 1.$$

These are termed the *selection rules* for electric dipole transitions (i.e., transitions calculated using the electric dipole approximation). Note, finally, that because the perturbing Hamiltonian does not contain any spin operators, the spin quantum number  $m_s$  cannot change during a transition. Hence, we have the additional selection rule that  $m'_s = m_s$ .

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [12.10: Selection Rules \(Hydrogen Atoms\)](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 12.11: 2P-1S Transitions in Hydrogen

Let us calculate the rate of spontaneous emission between the first excited state (i.e.,  $n = 2$ ) and the ground-state (i.e.,  $n' = 1$ ) of a hydrogen atom. Now, the ground-state is characterized by  $l' = m' = 0$ . Hence, in order to satisfy the selection rules ([e13.133]) and ([e13.134]), the excited state must have the quantum numbers  $l = 1$  and  $m = 0, \pm 1$ . Thus, we are dealing with a spontaneous transition from a  $2P$  to a  $1S$  state. Note, incidentally, that a spontaneous transition from a  $2S$  to a  $1S$  state is forbidden by our selection rules.

According to Section [s10.4], the wavefunction of a hydrogen atom takes the form

$$\psi_{n,l,m}(r, \theta, \phi) = R_{n,l}(r) Y_{l,m}(\theta, \phi), \quad (12.11.1)$$

where the radial functions  $R_{n,l}$  are given in Section [s10.4], and the spherical harmonics  $Y_{l,m}$  are given in Section [sharm]. Some straightforward, but tedious, integration reveals that

$$\begin{aligned} \langle 1, 0, 0 | x | 2, 1, \pm 1 \rangle &= \pm \frac{2^7}{3^5} a_0, \\ \langle 1, 0, 0 | y | 2, 1, \pm 1 \rangle &= i \frac{2^7}{3^5} a_0, \\ \langle 1, 0, 0 | z | 2, 1, 0 \rangle &= \sqrt{2} \frac{2^7}{3^5} a_0, \end{aligned}$$

where  $a_0$  is the Bohr radius specified in Equation ([e9.57]). All of the other possible  $2P \rightarrow 1S$  matrix elements are zero because of the selection rules. It follows from Equation ([e13.128]) that the modulus squared of the dipole moment for the  $2P \rightarrow 1S$  transition takes the same value

$$d^2 = \frac{2^{15}}{3^{10}} (e a_0)^2 \quad (12.11.2)$$

for  $m = 0, 1$ , or  $-1$ . Clearly, the transition rate is independent of the quantum number  $m$ . It turns out that this is a general result.

Now, the energy of the eigenstate of the hydrogen atom characterized by the quantum numbers  $n, l, m$  is  $E = E_0/n^2$ , where the ground-state energy  $E_0$  is specified in Equation ([e9.56]). Hence, the energy of the photon emitted during a  $2P \rightarrow 1S$  transition is

$$\hbar \omega = E_0/4 - E_0 = -\frac{3}{4} E_0 = 10.2 \text{ eV}. \quad (12.11.3)$$

This corresponds to a wavelength of  $1.215 \times 10^{-7} \text{ m}$ .

Finally, according to Equation ([e3.115]), the  $2P \rightarrow 1S$  transition rate is written

$$w_{2P \rightarrow 1S} = \frac{\omega^3 d^2}{3\pi \epsilon_0 \hbar c^3}, \quad (12.11.4)$$

which reduces to

$$w_{2P \rightarrow 1S} = \left(\frac{2}{3}\right)^8 \alpha^5 \frac{m_e c^2}{\hbar} = 6.27 \times 10^8 \text{ s}^{-1} \quad (12.11.5)$$

with the aid of Equations ([e13.139]) and ([e13.140]). Here,  $\alpha = 1/137$  is the fine-structure constant. Hence, the mean life-time of a hydrogen  $2P$  state is

$$\tau_{2P} = (w_{2P \rightarrow 1S})^{-1} = 1.6 \text{ ns}. \quad (12.11.6)$$

Incidentally, because the  $2P$  state only has a finite life-time, it follows from the energy-time uncertainty relation that the energy of this state is uncertain by an amount

$$\Delta E_{2P} \sim \frac{\hbar}{\tau_{2P}} \sim 4 \times 10^{-7} \text{ eV}. \quad (12.11.7)$$

This uncertainty gives rise to a finite width of the spectral line associated with the  $2P \rightarrow 1S$  transition. This natural line-width is of order

$$\frac{\Delta\lambda}{\lambda} \sim \frac{\Delta E_{2P}}{\hbar\omega} \sim 4 \times 10^{-8} \quad (12.11.8)$$

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [12.11: 2P-1S Transitions in Hydrogen](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).



## 12.12: Intensity Rules

Now, we know, from Section [\[s12.8\]](#), that when we take electron spin and spin-orbit coupling into account the degeneracy of the six  $2P$  states of the hydrogen atom is broken. In fact, these states are divided into two groups with slightly different energies. There are four states characterized by the overall angular momentum quantum number  $j = 3/2$ —these are called the  $2P_{3/2}$  states. The remaining two states are characterized by  $j = 1/2$ , and are thus called the  $2P_{1/2}$  states. The energy of the  $2P_{3/2}$  states is slightly higher than that of the  $2P_{1/2}$  states. In fact, the energy difference is

$$\Delta E = -\frac{\alpha^2}{16} E_0 = 4.53 \times 10^{-5} \text{ eV}. \quad (12.12.1)$$

Thus, the wavelength of the spectral line associated with the  $2P \rightarrow 1S$  transition in hydrogen is split by a relative amount

$$\frac{\Delta\lambda}{\lambda} = \frac{\Delta E}{\hbar\omega} = 4.4 \times 10^{-6}. \quad (12.12.2)$$

Note that this splitting is much greater than the natural line-width estimated in Equation ([\[e13.144a\]](#)), so there really are two spectral lines. How does all of this affect the rate of the  $2P \rightarrow 1S$  transition?

Well, we have seen that the transition rate is independent of spin, and hence of the spin quantum number  $m_s$ , and is also independent of the quantum number  $m$ . It follows that the transition rate is independent of the  $z$ -component of total angular momentum quantum number  $m_j = m + m_s$ . However, if this is the case then the transition rate is plainly also independent of the total angular momentum quantum number  $j$ . Hence, we expect the  $2P_{3/2} \rightarrow 1S$  and  $2P_{1/2} \rightarrow 1S$  transition rates to be the same. However, there are four  $2P_{3/2}$  states and only two  $2P_{1/2}$  states. If these states are equally populated—which we would certainly expect to be the case in thermal equilibrium, because they have almost the same energies—and given that they decay to the  $1S$  state at the same rate, it stands to reason that the spectral line associated with the  $2P_{3/2} \rightarrow 1S$  transition is twice as bright as that associated with the  $2P_{1/2} \rightarrow 1S$  transition.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [12.12: Intensity Rules](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 12.13: Forbidden Transitions

Atomic transitions which are forbidden by the electric dipole selection rules ([\[e13.133\]](#)) and ([\[e13.134\]](#)) are unsurprisingly known as *forbidden transitions*. It is clear from the analysis in Section [1.8](#) that a forbidden transition is one for which the matrix element  $\langle f | \epsilon \cdot \mathbf{p} | i \rangle$  is zero. However, this matrix element is only an approximation to the true matrix element for radiative transitions, which takes the form  $\langle f | \epsilon \cdot \mathbf{p} \exp(i \mathbf{k} \cdot \mathbf{r}) | i \rangle$ . Expanding  $\exp(i \mathbf{k} \cdot \mathbf{r})$ , and keeping the first two terms, the matrix element for a forbidden transition becomes

$$\langle f | \epsilon \cdot \mathbf{p} \exp(i \mathbf{k} \cdot \mathbf{r}) | i \rangle \simeq i \langle f | (\epsilon \cdot \mathbf{p}) (\mathbf{k} \cdot \mathbf{r}) | i \rangle. \quad (12.13.1)$$

Hence, if the residual matrix element on the right-hand side of the previous expression is non-zero then a “forbidden” transition can take place, albeit at a much reduced rate. In fact, in Section [1.9](#), we calculated that the typical rate of an electric dipole transition is

$$w_{i \rightarrow f} \sim \alpha^3 \omega_{if}. \quad (12.13.2)$$

Because the transition rate is proportional to the square of the radiative matrix element, it is clear that the transition rate for a forbidden transition enabled by the residual matrix element ([\[e13.146\]](#)) is smaller than that of an electric dipole transition by a factor  $(kr)^2$ . Estimating  $r$  as the Bohr radius, and  $k$  as the wavenumber of a typical spectral line of hydrogen, it is easily demonstrated that

$$w_{i \rightarrow f} \sim \alpha^5 \omega_{if} \quad (12.13.3)$$

for such a transition. Of course, there are some transitions (in particular, the  $2S \rightarrow 1S$  transition) for which the true radiative matrix element  $\langle f | \epsilon \cdot \mathbf{p} \exp(i \mathbf{k} \cdot \mathbf{r}) | i \rangle$  is zero. Such transitions are absolutely forbidden.

Finally, it is fairly obvious that excited states which decay via forbidden transitions have much longer life-times than those which decay via electric dipole transitions. Because the natural width of a spectral line is inversely proportional to the life-time of the associated decaying state, it follows that spectral lines associated with forbidden transitions are generally much sharper than those associated with electric dipole transitions.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [12.13: Forbidden Transitions](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 12.E: Time-Dependent Perturbation Theory (Exercises)

1. Consider the two-state system examined in Section 1.3.[ex8.1] Suppose that

$$\begin{aligned}\langle 1 | H_1 | 1 \rangle &= e_{11}, \\ \langle 2 | H_1 | 2 \rangle &= e_{22}, \\ \langle 1 | H_1 | 2 \rangle &= \langle 2 | H_1 | 1 \rangle^* = \frac{1}{2} \gamma \hbar \exp(i \omega t),\end{aligned}$$

where  $e_{11}$ ,  $e_{22}$ ,  $\gamma$ , and  $\omega$  are real. Show that

$$\begin{aligned}i \frac{d\hat{c}_1}{dt} &= \frac{\gamma}{2} \exp[+i(\omega - \hat{\omega}_{21})t] \hat{c}_2, \\ i \frac{d\hat{c}_2}{dt} &= \frac{\gamma}{2} \exp[-i(\omega - \hat{\omega}_{21})t] \hat{c}_1,\end{aligned}$$

where  $\hat{c}_1 = c_1 \exp(i e_{11} t / \hbar)$ ,  $\hat{c}_2 = c_2 \exp(i e_{22} t / \hbar)$ , and

$$\hat{\omega}_{21} = \frac{E_2 + e_{22} - E_1 - e_{11}}{\hbar}. \quad (12.E.1)$$

Hence, deduce that if the system is definitely in state 1 at time  $t = 0$  then the probability of finding it in state 2 at some subsequent time,  $t$ , is

$$P_2(t) = \frac{\gamma^2}{\gamma^2 + (\omega - \hat{\omega}_{21})^2} \sin^2 \left( \left[ \gamma^2 + (\omega - \hat{\omega}_{21})^2 \right]^{1/2} \frac{t}{2} \right). \quad (12.E.2)$$

2. Consider an atomic nucleus of spin- $s$  and gyromagnetic ratio  $g$  placed in the magnetic field

$$\mathbf{B} = B_0 \mathbf{e}_z + B_1 [\cos(\omega t) \mathbf{e}_x - \sin(\omega t) \mathbf{e}_y], \quad (12.E.3)$$

where  $B_1 \ll B_0$ . Let  $\chi_{s,m}$  be a properly normalized simultaneous eigenstate of  $S^2$  and  $S_z$ , where  $\mathbf{S}$  is the nuclear spin. Thus,  $S^2 \chi_{s,m} = s(s+1) \hbar^2 \chi_{s,m}$  and  $S_z \chi_{s,m} = m \hbar \chi_{s,m}$ , where  $-s \leq m \leq s$ . Furthermore, the instantaneous nuclear spin state is written

$$\chi = \sum_{m=-s,s} c_m(r) \chi_{s,m}, \quad (12.E.4)$$

where  $\sum_{m=-s,s} |c_m|^2 = 1$ .

1. Demonstrate that

$$\begin{aligned}\frac{dc_m}{dt} &= \frac{i\gamma}{2} \left( [s(s+1) - m(m-1)]^{1/2} e^{i(\omega - \omega_0)t} c_{m-1} \right. \\ &\quad \left. + [s(s+1) - m(m+1)]^{1/2} e^{-i(\omega - \omega_0)t} c_{m+1} \right)\end{aligned}$$

for  $-s \leq m \leq s$ , where  $\omega_0 = g \mu_N B_0 / \hbar$ ,  $\gamma = g \mu_N B_1 / \hbar$ , and  $\mu_N = e \hbar / (2 m_p)$ .

2. Consider the case  $s = 1/2$ . Demonstrate that if  $\omega = \omega_0$  and  $c_{1/2}(0) = 1$  then

$$c_{1/2}(t) = \cos(\gamma t / 2), \quad c_{-1/2}(t) = i \sin(\gamma t / 2).$$

3. Consider the case  $s = 1$ . Demonstrate that if  $\omega = \omega_0$  and  $c_1(0) = 1$  then

$$\begin{aligned}c_1(t) &= \cos^2(\gamma t / 2), \\ c_0(t) &= i\sqrt{2} \cos(\gamma t / 2) \sin(\gamma t / 2), \\ c_{-1}(t) &= -\sin^2(\gamma t / 2).\end{aligned}$$

4. Consider the case  $s = 3/2$ . Demonstrate that if  $\omega = \omega_0$  and  $c_{3/2}(0) = 1$  then

$$\begin{aligned}c_{3/2}(t) &= \cos^3(\gamma t/2), \\c_{1/2}(t) &= i\sqrt{3} \cos(\gamma t/2) \sin^2(\gamma t/2), \\c_{-1/2}(t) &= -\sqrt{3} \cos^2(\gamma t/2) \sin(\gamma t/2), \\c_{-3/2}(t) &= -i \sin^3(\gamma t/2).\end{aligned}$$

3. Demonstrate that a spontaneous transition between two atomic states of zero orbital angular momentum is absolutely forbidden. (Actually, a spontaneous transition between two zero orbital angular momentum states is possible via the simultaneous emission of two photons, but takes place at a very slow rate .)

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [12.E: Time-Dependent Perturbation Theory \(Exercises\)](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## CHAPTER OVERVIEW

### 13: Variational Methods

We have seen, in Sect. 8.3, that we can solve Schrödinger's equation *exactly* to find the stationary eigenstates of a hydrogen atom. Unfortunately, it is not possible to find exact solutions of Schrödinger's equation for atoms more complicated than hydrogen, or for molecules. In such systems, the best that we can do is to find *approximate* solutions. Most of the methods which have been developed for finding such solutions employ the so-called *variational principle* discussed below.

[13.1: Variational Principle](#)

[13.2: Helium Atom](#)

[13.3: Hydrogen Molecule Ion](#)

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [13: Variational Methods](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 13.1: Variational Principle

Suppose that we wish to solve the time-independent Schrödinger equation

$$H \psi = E \psi, \quad (13.1.1)$$

where  $H$  is a known (presumably complicated) time-independent Hamiltonian. Let  $\psi$  be a properly normalized trial solution to the previous equation. The variational principle states, quite simply, that the ground-state energy,  $E_0$ , is always less than or equal to the expectation value of  $H$  calculated with the trial wavefunction: that is,

$$E_0 \leq \langle \psi | H | \psi \rangle. \quad (13.1.2)$$

Thus, by varying  $\psi$  until the expectation value of  $H$  is minimized, we can obtain approximations to the wavefunction and the energy of the ground-state.

Let us prove the variational principle. Suppose that the  $\psi_n$  and the  $E_n$  are the true eigenstates and eigenvalues of  $H$ : that is,

$$H \psi_n = E_n \psi_n. \quad (13.1.3)$$

Furthermore, let

$$E_0 < E_1 < E_2 < \dots, \quad (13.1.4)$$

so that  $\psi_0$  is the ground-state,  $\psi_1$  the first excited state, et cetera. The  $\psi_n$  are assumed to be orthonormal: that is,

$$\langle \psi_n | \psi_m \rangle = \delta_{nm}. \quad (13.1.5)$$

If our trial wavefunction  $\psi$  is properly normalized then we can write

$$\psi = \sum_n c_n \psi_n, \quad (13.1.6)$$

where

$$\sum_n |c_n|^2 = 1. \quad (13.1.7)$$

Now, the expectation value of  $H$ , calculated with  $\psi$ , takes the form

$$\begin{aligned} \langle \psi | H | \psi \rangle &= \left\langle \sum_n c_n \psi_n \left| H \right| \sum_m c_m \psi_m \right\rangle = \sum_{n,m} c_n^* c_m \langle \psi_n | H | \psi_m \rangle \\ &= \sum_n c_n^* c_m E_m \langle \psi_n | \psi_m \rangle = \sum_n E_n |c_n|^2, \end{aligned}$$

where use has been made of Equations 13.1.3 and 13.1.5. So, we can write

$$\langle \psi | H | \psi \rangle = |c_0|^2 E_0 + \sum_{n>0} |c_n|^2 E_n. \quad (13.1.8)$$

However, Equation 13.1.7 can be rearranged to give

$$|c_0|^2 = 1 - \sum_{n>0} |c_n|^2. \quad (13.1.9)$$

Combining the previous two equations, we obtain

$$\langle \psi | H | \psi \rangle = E_0 + \sum_{n>0} |c_n|^2 (E_n - E_0). \quad (13.1.10)$$

The second term on the right-hand side of the previous expression is positive definite, because  $E_n - E_0 > 0$  for all  $n > 0$  (Equation 13.1.4). Hence, we obtain the desired result

$$\langle \psi | H | \psi \rangle \geq E_0. \quad (13.1.11)$$

## Excited States

Suppose that we have found a good approximation,  $\tilde{\psi}_0$ , to the ground-state wavefunction. If  $\psi$  is a normalized trial wavefunction that is orthogonal to  $\tilde{\psi}_0$  (i.e.,  $\langle \psi | \tilde{\psi}_0 \rangle = 0$ ) then, by repeating the previous analysis, we can easily demonstrate that

$$\langle \psi | H | \psi \rangle \geq E_1. \quad (13.1.12)$$

Thus, by varying  $\psi$  until the expectation value of  $H$  is minimized, we can obtain approximations to the wavefunction and the energy of the first excited state. Obviously, we can continue this process until we have approximations to all of the stationary eigenstates. Note, however, that the errors are clearly cumulative in this method, so that any approximations to highly excited states are unlikely to be very accurate. For this reason, the variational method is generally only used to calculate the ground-state and first few excited states of complicated quantum systems.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [13.1: Variational Principle](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 13.2: Helium Atom

A helium atom consists of a nucleus of charge  $+2e$  surrounded by two electrons. Let us attempt to calculate its ground-state energy.

Let the nucleus lie at the origin of our coordinate system, and let the position vectors of the two electrons be  $\mathbf{r}_1$  and  $\mathbf{r}_2$ , respectively. The Hamiltonian of the system thus takes the form

$$H = -\frac{\hbar^2}{2m_e}(\nabla_1^2 + \nabla_2^2) - \frac{e^2}{4\pi\epsilon_0}\left(\frac{2}{r_1} + \frac{2}{r_2} - \frac{1}{|\mathbf{r}_2 - \mathbf{r}_1|}\right), \quad (13.2.1)$$

where we have neglected any reduced mass effects. The terms in the previous expression represent the kinetic energy of the first electron, the kinetic energy of the second electron, the electrostatic attraction between the nucleus and the first electron, the electrostatic attraction between the nucleus and the second electron, and the electrostatic repulsion between the two electrons, respectively. It is the final term that causes all of the difficulties. Indeed, if this term is neglected then we can write

$$H = H_1 + H_2, \quad (13.2.2)$$

where

$$H_{1,2} = -\frac{\hbar^2}{2m_e}\nabla_{1,2}^2 - \frac{2e^2}{4\pi\epsilon_0 r_{1,2}}. \quad (13.2.3)$$

In other words, the Hamiltonian just becomes the sum of separate Hamiltonians for each electron. In this case, we would expect the wavefunction to be separable: that is,

$$\psi(\mathbf{r}_1, \mathbf{r}_2) = \psi_1(\mathbf{r}_1)\psi_2(\mathbf{r}_2). \quad (13.2.4)$$

Hence, Schrödinger's equation,

$$H\psi = E\psi, \quad (13.2.5)$$

reduces to

$$H_{1,2}\psi_{1,2} = E_{1,2}\psi_{1,2}, \quad (13.2.6)$$

where

$$E = E_1 + E_2. \quad (13.2.7)$$

Of course, Equation ??? is the Schrödinger equation of a hydrogen atom whose nuclear charge is  $+2e$ , instead of  $+e$ . It follows, from Section [s10.4] (making the substitution  $e^2 \rightarrow 2e^2$ ), that if both electrons are in their lowest energy states then

$$\begin{aligned} \psi_1(\mathbf{r}_1) &= \psi_0(\mathbf{r}_1), \\ \psi_2(\mathbf{r}_2) &= \psi_0(\mathbf{r}_2), \end{aligned}$$

where

$$\psi_0(\mathbf{r}) = \frac{4}{\sqrt{2\pi}a_0^{3/2}}\exp\left(-\frac{2r}{a_0}\right). \quad (13.2.8)$$

Here,  $a_0$  is the Bohr radius. [See Equation ([e9.57]).] Note that  $\psi_0$  is properly normalized. Furthermore,

$$E_1 = E_2 = 4E_0, \quad (13.2.9)$$

where  $E_0 = -13.6$  eV is the hydrogen ground-state energy. [See Equation ([e9.56]).] Thus, our crude estimate for the ground-state energy of helium becomes

$$E = 4E_0 + 4E_0 = 8E_0 = -108.8 \text{ eV}. \quad (13.2.10)$$

Unfortunately, this estimate is significantly different from the experimentally determined value, which is  $-78.98$  eV. This fact demonstrates that the neglected electron-electron repulsion term makes a large contribution to the helium ground-state energy. Fortunately, however, we can use the variational principle to estimate this contribution.



Let us employ the separable wavefunction discussed previously as our trial solution. Thus,

$$\psi(\mathbf{r}_1, \mathbf{r}_2) = \psi_0(\mathbf{r}_1) \psi_0(\mathbf{r}_2) = \frac{8}{\pi a_0^3} \exp\left(-\frac{2[r_1 + r_2]}{a_0}\right). \quad (13.2.11)$$

The expectation value of the Hamiltonian 13.2.1 thus becomes

$$\langle H \rangle = 8 E_0 + \langle V_{ee} \rangle, \quad (13.2.12)$$

where

$$\langle V_{ee} \rangle = \left\langle \psi \left| \frac{e^2}{4\pi \epsilon_0 |\mathbf{r}_2 - \mathbf{r}_1|} \right| \psi \right\rangle = \frac{e^2}{4\pi \epsilon_0} \int \frac{|\psi(\mathbf{r}_1, \mathbf{r}_2)|^2}{|\mathbf{r}_2 - \mathbf{r}_1|} d^3 \mathbf{r}_1 d^3 \mathbf{r}_2. \quad (13.2.13)$$

The variation principle only guarantees that Equation 13.2.12 yields an upper bound on the ground-state energy. In reality, we hope that it will give a reasonably accurate estimate of this energy.

It follows from Equations ([e9.56]), 13.2.11, and 13.2.13 that

$$\langle V_{ee} \rangle = -\frac{4 E_0}{\pi^2} \int \frac{e^{-2(\hat{r}_1 + \hat{r}_2)}}{|\hat{\mathbf{r}}_1 - \hat{\mathbf{r}}_2|} d^3 \hat{\mathbf{r}}_1 d^3 \hat{\mathbf{r}}_2, \quad (13.2.14)$$

where  $\hat{\mathbf{r}}_{1,2} = 2 \mathbf{r}_{1,2}/a_0$ . Neglecting the hats, for the sake of clarity, the previous expression can also be written

$$\langle V_{ee} \rangle = -\frac{4 E_0}{\pi^2} \int \frac{e^{-2(r_1 + r_2)}}{\sqrt{r_1^2 + r_2^2 - 2 r_1 r_2 \cos \theta}} d^3 \mathbf{r}_1 d^3 \mathbf{r}_2, \quad (13.2.15)$$

where  $\theta$  is the angle subtended between vectors  $\mathbf{r}_1$  and  $\mathbf{r}_2$ . If we perform the integral in  $\mathbf{r}_1$  space before that in  $\mathbf{r}_2$  space then

$$\langle V_{ee} \rangle = -\frac{4 E_0}{\pi^2} \int e^{-2 r_2} I(\mathbf{r}_2) d^3 \mathbf{r}_2, \quad (13.2.16)$$

where

$$I(\mathbf{r}_2) = \int \frac{e^{-2 r_1}}{\sqrt{r_1^2 + r_2^2 - 2 r_1 r_2 \cos \theta}} d^3 \mathbf{r}_1. \quad (13.2.17)$$

Our first task is to evaluate the function  $I(\mathbf{r}_2)$ . Let  $(r_1, \theta_1, \phi_1)$  be a set of spherical coordinates in  $\mathbf{r}_1$  space whose axis of symmetry runs in the direction of  $\mathbf{r}_2$ . It follows that  $\theta = \theta_1$ . Hence,

$$I(\mathbf{r}_2) = \int_0^\infty \int_0^\pi \int_0^{2\pi} \frac{e^{-2 r_1}}{\sqrt{r_1^2 + r_2^2 - 2 r_1 r_2 \cos \theta_1}} r_1^2 dr_1 \sin \theta_1 d\theta_1 d\phi_1, \quad (13.2.18)$$

which trivially reduces to

$$I(\mathbf{r}_2) = 2\pi \int_0^\infty \int_0^\pi \frac{e^{-2 r_1}}{\sqrt{r_1^2 + r_2^2 - 2 r_1 r_2 \cos \theta_1}} r_1^2 dr_1 \sin \theta_1 d\theta_1. \quad (13.2.19)$$

Making the substitution  $\mu = \cos \theta_1$ , we can see that

$$\int_0^\pi \frac{1}{\sqrt{r_1^2 + r_2^2 - 2 r_1 r_2 \cos \theta_1}} \sin \theta_1 d\theta_1 = \int_{-1}^1 \frac{d\mu}{\sqrt{r_1^2 + r_2^2 - 2 r_1 r_2 \mu}}. \quad (13.2.20)$$

Now,

$$\begin{aligned}\int_{-1}^1 \frac{d\mu}{\sqrt{r_1^2 + r_2^2 - 2 r_1 r_2 \mu}} &= \left[ \frac{\sqrt{r_1^2 + r_2^2 - 2 r_1 r_2 \mu}}{r_1 r_2} \right]_{-1}^{-1} \\ &= \frac{(r_1 + r_2) - |r_1 - r_2|}{r_1 r_2} \\ &= \begin{cases} 2/r_1 & \text{for } r_1 > r_2 \\ 2/r_2 & \text{for } r_1 < r_2 \end{cases},\end{aligned}$$

giving

$$I(\mathbf{r}_2) = 4\pi \left( \frac{1}{r_2} \int_0^{r_2} e^{-2r_1} r_1^2 dr_1 + \int_{r_2}^{\infty} e^{-2r_1} r_1 dr_1 \right). \quad (13.2.21)$$

But ,

$$\begin{aligned}\int e^{-\beta x} x dx &= -\frac{e^{-\beta x}}{\beta^2} (1 + \beta x), \\ \int e^{-\beta x} x^2 dx &= -\frac{e^{-\beta x}}{\beta^3} (2 + 2\beta x + \beta^2 x^2),\end{aligned}$$

yielding

$$I(\mathbf{r}_2) = \frac{\pi}{r_2} [1 - e^{-2r_2} (1 + r_2)]. \quad (13.2.22)$$

Because the function  $I(\mathbf{r}_2)$  only depends on the magnitude of  $\mathbf{r}_2$ , the integral in Equation 13.2.16 reduces to

$$\langle V_{ee} \rangle = -\frac{16 E_0}{\pi} \int_0^{\infty} e^{-2r_2} I(r_2) r_2^2 dr_2, \quad (13.2.23)$$

which yields

$$\langle V_{ee} \rangle = -16 E_0 \int_0^{\infty} e^{-2r_2} [1 - e^{-2r_2} (1 + r_2)] r_2 dr_2 = -\frac{5}{2} E_0. \quad (13.2.24)$$

Hence, from Equation 13.2.12 our estimate for the ground-state energy of helium is

$$\langle H \rangle = 8 E_0 - \frac{5}{2} E_0 = \frac{11}{2} E_0 = -74.8 \text{ eV}. \quad (13.2.25)$$

This is remarkably close to the correct result.

## Shielding and Effective Nuclear Charge

We can actually refine our estimate further. The trial wavefunction 13.2.11 essentially treats the two electrons as non-interacting particles. In reality, we would expect one electron to partially shield the nuclear charge from the other, and vice versa. Hence, a better trial wavefunction might be

$$\psi(\mathbf{r}_1, \mathbf{r}_2) = \frac{Z^3}{\pi a_0^3} \exp\left(-\frac{Z[r_1 + r_2]}{a_0}\right), \quad (13.2.26)$$

where  $Z < 2$  is effective nuclear charge number seen by each electron. Let us recalculate the ground-state energy of helium as a function of  $Z$ , using the previous trial wavefunction, and then minimize the result with respect to  $Z$ . According to the variational principle, this should give us an even better estimate for the ground-state energy.

We can rewrite the expression 13.2.1 for the Hamiltonian of the helium atom in the form

$$H = H_1(Z) + H_2(Z) + V_{ee} + U(Z), \quad (13.2.27)$$

where

$$H_{1,2}(Z) = -\frac{\hbar^2}{2m_e} \nabla_{1,2}^2 - \frac{Ze^2}{4\pi\epsilon_0 r_{1,2}} \quad (13.2.28)$$

is the Hamiltonian of a hydrogen atom with nuclear charge  $+Ze$ ,

$$V_{ee} = \frac{e^2}{4\pi\epsilon_0} \frac{1}{|\mathbf{r}_2 - \mathbf{r}_1|} \quad (13.2.29)$$

is the electron-electron repulsion term, and

$$U(Z) = \frac{e^2}{4\pi\epsilon_0} \left( \frac{[Z-2]}{r_1} + \frac{[Z-2]}{r_2} \right). \quad (13.2.30)$$

It follows that

$$\langle H \rangle(Z) = 2E_0(Z) + \langle V_{ee} \rangle(Z) + \langle U \rangle(Z), \quad (13.2.31)$$

where  $E_0(Z) = Z^2 E_0$  is the ground-state energy of a hydrogen atom with nuclear charge  $+Ze$ ,  $\langle V_{ee} \rangle(Z) = -(5Z/4)E_0$  is the value of the electron-electron repulsion term when recalculated with the wavefunction in Equation 13.2.26 [actually, all we need to do is to make the substitution  $a_0 \rightarrow (2/Z)a_0$ ], and

$$\langle U \rangle(Z) = 2(Z-2) \left( \frac{e^2}{4\pi\epsilon_0} \right) \left\langle \frac{1}{r} \right\rangle. \quad (13.2.32)$$

Here,  $\langle 1/r \rangle$  is the expectation value of  $1/r$  calculated for a hydrogen atom with nuclear charge  $+Ze$ . It follows from Equation ([e9.74]) [with  $n=1$ , and making the substitution  $a_0 \rightarrow a_0/Z$ ] that

$$\left\langle \frac{1}{r} \right\rangle = \frac{Z}{a_0}. \quad (13.2.33)$$

Hence,

$$\langle U \rangle(Z) = -4Z(Z-2)E_0, \quad (13.2.34)$$

because  $E_0 = -e^2/(8\pi\epsilon_0 a_0)$ . Collecting the various terms, our new expression for the expectation value of the Hamiltonian becomes

$$\langle H \rangle(Z) = \left[ 2Z^2 - \frac{5}{4}Z - 4Z(Z-2) \right] E_0 = \left( -2Z^2 + \frac{27}{4}Z \right) E_0. \quad (13.2.35)$$

The value of  $Z$  that minimizes this expression is the root of

$$\frac{d\langle H \rangle}{dZ} = \left( -4Z + \frac{27}{4} \right) E_0 = 0. \quad (13.2.36)$$

It follows that

$$Z = \frac{27}{16} = 1.69. \quad (13.2.37)$$

The fact that  $Z < 2$  confirms our earlier conjecture that the electrons partially shield the nuclear charge from one another. Our new estimate for the ground-state energy of helium is

$$\langle H \rangle(1.69) = \frac{1}{2} \left( \frac{3}{2} \right)^6 E_0 = -77.5 \text{ eV}. \quad (13.2.38)$$

This is clearly an improvement on our previous estimate in Equation 13.2.25 (Recall that the correct result is  $-78.98 \text{ eV}$ .)

Obviously, we could get even closer to the correct value of the helium ground-state energy by using a more complicated trial wavefunction with more adjustable parameters.

Note, finally, that because the two electrons in a helium atom are indistinguishable fermions, the overall wavefunction must be *anti-symmetric* with respect to exchange of particles. (See Chapter [smany].) Now, the overall wavefunction is the product of the spatial wavefunction and the spinor representing the spin-state. Our spatial wavefunction ([e14.44]) is obviously symmetric with

respect to exchange of particles. This means that the spinor must be anti-symmetric. It is clear, from Section [\[shalf\]](#), that if the spin-state of an  $l = 0$  system consisting of two spin one-half particles (i.e., two electrons) is anti-symmetric with respect to interchange of particles then the system is in the so-called singlet state with overall spin zero. Hence, the ground-state of helium has overall electron spin zero.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [13.2: Helium Atom](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 13.3: Hydrogen Molecule Ion

The hydrogen molecule ion consists of an electron orbiting about two protons, and is the simplest imaginable molecule. Let us investigate whether or not this molecule possesses a bound state: that is, whether or not it possesses a ground-state whose energy is less than that of a hydrogen atom and a free proton. According to the variation principle, we can deduce that the  $H_2^+$  ion has a bound state if we can find any trial wavefunction for which the total Hamiltonian of the system has an expectation value less than that of a hydrogen atom and a free proton.




Figure 26: The hydrogen molecule ion.

Suppose that the two protons are separated by a distance  $R$ . In fact, let them lie on the  $z$ -axis, with the first at the origin, and the second at  $z = R$ . See Figure [fh2p]. In the following, we shall treat the protons as essentially stationary. This is reasonable because the electron moves far more rapidly than the protons. Incidentally, the neglect of nuclear motion when calculating the electronic energy of the molecule is known as the *Born-Oppenheimer approximation*.

Let us try

$$\psi(\mathbf{r})_{\pm} = A [\psi_0(\mathbf{r}_1) \pm \psi_0(\mathbf{r}_2)] \quad (13.3.1)$$

as our trial wavefunction, where

$$\psi_0(\mathbf{r}) = \frac{1}{\sqrt{\pi} a_0^{3/2}} e^{-r/a_0} \quad (13.3.2)$$

is a normalized hydrogen ground-state wavefunction centered on the origin, and  $\mathbf{r}_{1,2}$  are the position vectors of the electron with respect to each of the protons. See Figure [fh2p]. Obviously, this is a very simplistic wavefunction, because it is just a linear combination of hydrogen ground-state wavefunctions centered on each proton. Note, however, that the wavefunction respects the obvious symmetries in the problem.

Our first task is to normalize our trial wavefunction. We require that

$$\int |\psi_{\pm}|^2 d^3\mathbf{r} = 1. \quad (13.3.3)$$

Hence, from Equation ([e14.57]),  $A = I^{-1/2}$ , where

$$I = \int \left[ |\psi_0(\mathbf{r}_1)|^2 + |\psi_0(\mathbf{r}_2)|^2 \pm 2 \psi_0(\mathbf{r}_1) \psi_0(\mathbf{r}_2) \right] d^3\mathbf{r}. \quad (13.3.4)$$

It follows that

$$I = 2 (1 \pm J), \quad (13.3.5)$$

with

$$J = \int \psi_0(\mathbf{r}_1) \psi_0(\mathbf{r}_2) d^3\mathbf{r}. \quad (13.3.6)$$

Let us employ the standard spherical coordinates  $(r, \theta, \phi)$ . Now, it is easily seen that  $r_1 = r$  and  $r_2 = (r^2 + R^2 - 2rR \cos \theta)^{1/2}$ . Hence,

$$J = 2 \int_0^\infty \int_0^\pi \exp[-x - (x^2 + X^2 - 2xX \cos \theta)^{1/2}] x^2 dx \sin \theta d\theta, \quad (13.3.7)$$

where  $X = R/a_0$ . Here, we have already performed the trivial  $\phi$  integral. Let  $y = (x^2 + X^2 - 2xX \cos \theta)^{1/2}$ . It follows that  $d(y^2) = 2y dy = 2xX \sin \theta d\theta$ , giving

$$\begin{aligned} \int_0^\pi e^{(x^2 + X^2 - 2xX \cos \theta)^{1/2}} \sin \theta d\theta &= \frac{1}{xX} \int_{|x-X|}^{x+X} e^{-y} y dy \\ &= -\frac{1}{xX} \left[ e^{-(x+X)} (1+x+X) - e^{-|x-X|} (1+|x-X|) \right]. \end{aligned}$$

Thus,

$$\begin{aligned} J &= -\frac{2}{X} e^{-X} \int_0^X [e^{-2x} (1+X+x) - (1+X-x)] x dx \\ &\quad - \frac{2}{X} \int_X^\infty e^{-2x} [e^{-X} (1+X+x) - e^X (1-X+x)] x dx, \end{aligned}$$

which evaluates to

$$J = e^{-X} \left( 1 + X + \frac{X^3}{3} \right). \quad (13.3.8)$$

Now, the Hamiltonian of the electron is written

$$H = -\frac{\hbar^2}{2m_e} \nabla^2 - \frac{e^2}{4\pi\epsilon_0} \left( \frac{1}{r_1} + \frac{1}{r_2} \right). \quad (13.3.9)$$

Note, however, that

$$\left( -\frac{\hbar^2}{2m_e} \nabla^2 - \frac{e^2}{4\pi\epsilon_0 r_{1,2}} \right) \psi_0(\mathbf{r}_{1,2}) = E_0 \psi_0(\mathbf{r}_{1,2}), \quad (13.3.10)$$

because  $\psi_0(\mathbf{r}_{1,2})$  are hydrogen ground-state wavefunctions. It follows that

$$\begin{aligned} H \psi_\pm &= A \left[ -\frac{\hbar^2}{2m_e} \nabla^2 - \frac{e^2}{4\pi\epsilon_0} \left( \frac{1}{r_1} + \frac{1}{r_2} \right) \right] [\psi_0(\mathbf{r}_1) \pm \psi_0(\mathbf{r}_2)] \\ &= E_0 \psi - A \left( \frac{e^2}{4\pi\epsilon_0} \right) \left[ \frac{\psi_0(\mathbf{r}_1)}{r_2} \pm \frac{\psi_0(\mathbf{r}_2)}{r_1} \right]. \end{aligned}$$

Hence,

$$\langle H \rangle = E_0 + 4A^2 (D \pm E) E_0, \quad (13.3.11)$$

where

$$\begin{aligned} D &= \left\langle \psi_0(\mathbf{r}_1) \left| \frac{a_0}{r_2} \right| \psi_0(\mathbf{r}_1) \right\rangle, \\ E &= \left\langle \psi_0(\mathbf{r}_1) \left| \frac{a_0}{r_1} \right| \psi_0(\mathbf{r}_2) \right\rangle. \end{aligned}$$

Now,

$$D = 2 \int_0^\infty \int_0^\pi \frac{e^{-2x}}{(x^2 + X^2 - 2xX \cos \theta)^{1/2}} x^2 dx \sin \theta d\theta, \quad (13.3.12)$$

which reduces to

$$D = \frac{4}{X} \int_0^X e^{-2x} x^2 dx + 4 \int_X^\infty e^{-2x} x dx, \quad (13.3.13)$$

giving

$$D = \frac{1}{X} (1 - [1 + X] e^{-2X}). \quad (13.3.14)$$

Furthermore,

$$E = 2 \int_0^\infty \int_0^\pi \exp[-x - (x^2 + X^2 - 2xX \cos \theta)^{1/2}] x dx \sin \theta d\theta, \quad (13.3.15)$$

which reduces to

$$\begin{aligned} E &= -\frac{2}{X} e^{-X} \int_0^X [e^{-2x} (1 + X + x) - (1 + X - x)] dx \\ &\quad - \frac{2}{X} \int_X^\infty e^{-2x} [e^{-X} (1 + X + x) - e^X (1 - X + x)] dx, \end{aligned}$$

yielding

$$E = (1 + X) e^{-X}. \quad (13.3.16)$$

Our expression for the expectation value of the electron Hamiltonian is

$$\langle H \rangle = \left[ 1 + 2 \frac{(D \pm E)}{(1 \pm J)} \right] E_0, \quad (13.3.17)$$

where  $J$ ,  $D$ , and  $E$  are specified as functions of  $X = R/a_0$  in Equations ([e14.66]), ([e14.75]), and ([e14.78]), respectively. In order to obtain the total energy of the molecule, we must add to this the potential energy of the two protons. Thus,

$$E_{\text{total}} = \langle H \rangle + \frac{e^2}{4\pi \epsilon_0 R} = \langle H \rangle - \frac{2}{X} E_0, \quad (13.3.18)$$

because  $E_0 = -e^2/(8\pi \epsilon_0 a_0)$ . Hence, we can write

$$E_{\text{total}} = -F_\pm(R/a_0) E_0, \quad (13.3.19)$$

where  $E_0$  is the hydrogen ground-state energy, and

$$F_\pm(X) = -1 + \frac{2}{X} \left[ \frac{(1 + X) e^{-2X} \pm (1 - 2X^2/3) e^{-X}}{1 \pm (1 + X + X^2/3) e^{-X}} \right]. \quad (13.3.20)$$

The functions  $F_+(X)$  and  $F_-(X)$  are both plotted in Figure [fh2pa]. Recall that in order for the  $H_2^+$  ion to be in a bound state it must have a lower energy than a hydrogen atom and a free proton: that is,  $E_{\text{total}} < E_0$ . It follows from Equation ([e14.81]) that a bound state corresponds to  $F_\pm < -1$ . Clearly, the even trial wavefunction  $\psi_+$  possesses a bound state, whereas the odd trial wavefunction  $\psi_-$  does not. [See Equation ([e14.57]).] This is hardly surprising, because the even wavefunction maximizes the electron probability density between the two protons, thereby reducing their mutual electrostatic repulsion. On the other hand, the odd wavefunction does exactly the opposite. The *binding energy* of the  $H_2^+$  ion is defined as the difference between its energy and that of a hydrogen atom and a free proton: that is,

$$E_{\text{bind}} = E_{\text{total}} - E_0 = -(F_+ + 1) E_0. \quad (13.3.21)$$

According to the variational principle, the binding energy is less than or equal to the minimum binding energy that can be inferred from Figure [fh2pa]. This minimum occurs when  $X \simeq 2.5$  and  $F_+ \simeq -1.13$ . Thus, our estimates for the separation between the two protons, and the binding energy, for the  $H_2^+$  ion are  $R = 2.5 a_0 = 1.33 \times 10^{-10}$  m and  $E_{\text{bind}} = 0.13 E_0 = -1.77$  eV,

respectively. The experimentally determined values are  $R = 1.06 \times 10^{-10}$  m, and  $E_{\text{bind}} = -2.8$  eV, respectively. Clearly, our estimates are not particularly accurate. However, our calculation does establish, beyond any doubt, the existence of a bound state of the  $H_2^+$  ion, which is all that we set out to achieve.




Figure 27: The functions  $F_+(X)$  (solid curve) and  $F_-(X)$  (dashed curve).

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

This page titled [13.3: Hydrogen Molecule Ion](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).



## CHAPTER OVERVIEW

### 14: Scattering Theory

Historically, data regarding quantum phenomena has been obtained from two main sources. Firstly, from the study of spectroscopic lines, and, secondly, from scattering experiments. We have already developed theories that account for some aspects of the spectra of hydrogen, and hydrogen-like, atoms. Let us now examine the quantum theory of scattering.

[14.1: Fundamentals of Scattering Theory](#)

[14.2: Born Approximation](#)

[14.3: Partial Waves](#)

[14.4: Optical Theorem](#)

[14.5: Determination of Phase-Shifts](#)

[14.6: Hard-Sphere Scattering](#)

[14.7: Low-Energy Scattering](#)

[14.8: Resonances](#)

---

This page titled [14: Scattering Theory](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 14.1: Fundamentals of Scattering Theory

Consider time-independent, energy conserving scattering in which the Hamiltonian of the system is written

$$H = H_0 + V(\mathbf{r}), \quad (14.1.1)$$

where

$$H_0 = \frac{p^2}{2m} \equiv -\frac{\hbar^2}{2m} \nabla^2 \quad (14.1.2)$$

is the Hamiltonian of a free particle of mass  $m$ , and  $V(\mathbf{r})$  the scattering potential. This potential is assumed to only be non-zero in a fairly localized region close to the origin. Let

$$\psi_0(\mathbf{r}) = \sqrt{n} e^{i\mathbf{k}\cdot\mathbf{r}} \quad (14.1.3)$$

represent an incident beam of particles, of number density  $n$ , and velocity  $\mathbf{v} = \hbar\mathbf{k}/m$ . [See Equation ([e14.14gg](#)).] Of course,

$$H_0 \psi_0 = E \psi_0, \quad (14.1.4)$$

where  $E = \hbar^2 k^2 / (2m)$  is the particle energy. Schrödinger's equation for the scattering problem is

$$(H_0 + V) \psi = E \psi, \quad (14.1.5)$$

subject to the boundary condition  $\psi \rightarrow \psi_0$  as  $V \rightarrow 0$ .

The previous equation can be rearranged to give

$$(\nabla^2 + k^2) \psi = \frac{2m}{\hbar^2} V \psi. \quad (14.1.6)$$

Now,

$$(\nabla^2 + k^2) u(\mathbf{r}) = \rho(\mathbf{r}) \quad (14.1.7)$$

is known as the *Helmholtz equation*. The solution to this equation is well known

$$u(\mathbf{r}) = u_0(\mathbf{r}) - \int \frac{e^{ik|\mathbf{r}-\mathbf{r}'|}}{4\pi|\mathbf{r}-\mathbf{r}'|} \rho(\mathbf{r}') d^3\mathbf{r}'. \quad (14.1.8)$$

Here,  $u_0(\mathbf{r})$  is any solution of  $(\nabla^2 + k^2) u_0 = 0$ . Hence, Equation ([e15.6](#)) can be inverted, subject to the boundary condition  $\psi \rightarrow \psi_0$  as  $V \rightarrow 0$ , to give

$$\psi(\mathbf{r}) = \psi_0(\mathbf{r}) - \frac{2m}{\hbar^2} \int \frac{e^{ik|\mathbf{r}-\mathbf{r}'|}}{4\pi|\mathbf{r}-\mathbf{r}'|} V(\mathbf{r}') \psi(\mathbf{r}') d^3\mathbf{r}' \quad (14.1.9)$$

Let us calculate the value of the wavefunction  $\psi(\mathbf{r})$  well outside the scattering region. Now, if  $r \gg r'$  then

$$|\mathbf{r} - \mathbf{r}'| \simeq r - \hat{\mathbf{r}} \cdot \mathbf{r}' \quad (14.1.10)$$

to first-order in  $r'/r$ , where  $\hat{\mathbf{r}}/r$  is a unit vector that points from the scattering region to the observation point. It is helpful to define  $\mathbf{k}' = k\hat{\mathbf{r}}$ . This is the wavevector for particles with the same energy as the incoming particles (i.e.,  $k' = k$ ) that propagate from the scattering region to the observation point. Equation ([e15.9](#)) reduces to

$$\psi(\mathbf{r}) \simeq \sqrt{n} \left[ e^{i\mathbf{k}\cdot\mathbf{r}} + \frac{e^{ikr}}{r} f(\mathbf{k}, \mathbf{k}') \right], \quad (14.1.11)$$

where

$$f(\mathbf{k}, \mathbf{k}') = -\frac{m}{2\pi\sqrt{n}\hbar^2} \int e^{-i\mathbf{k}'\cdot\mathbf{r}'} V(\mathbf{r}') \psi(\mathbf{r}') d^3\mathbf{r}'. \quad (14.1.12)$$

The first term on the right-hand side of Equation ([e15.11](#)) represents the incident particle beam, whereas the second term represents an outgoing spherical wave of scattered particles.

The *differential scattering cross-section*,  $d\sigma/d\Omega$ , is defined as the number of particles per unit time scattered into an element of solid angle  $d\Omega$ , divided by the incident particle flux. From Section [s7.2], the probability flux (i.e., the particle flux) associated with a wavefunction  $\psi$  is

$$\mathbf{j} = \frac{\hbar}{m} \text{Im}(\psi^* \nabla \psi). \quad (14.1.13)$$

Thus, the particle flux associated with the incident wavefunction  $\psi_0$  is

$$\mathbf{j} = n \mathbf{v}, \quad (14.1.14)$$

where  $\mathbf{v} = \hbar \mathbf{k}/m$  is the velocity of the incident particles. Likewise, the particle flux associated with the scattered wavefunction  $\psi - \psi_0$  is

$$\mathbf{j}' = n \frac{|f(\mathbf{k}, \mathbf{k}')|^2}{r^2} \mathbf{v}', \quad (14.1.15)$$

where  $\mathbf{v}' = \hbar \mathbf{k}'/m$  is the velocity of the scattered particles. Now,

$$\frac{d\sigma}{d\Omega} d\Omega = \frac{r^2 d\Omega |\mathbf{j}'|}{|\mathbf{j}|}, \quad (14.1.16)$$

which yields

$$\frac{d\sigma}{d\Omega} = |f(\mathbf{k}, \mathbf{k}')|^2. \quad (14.1.17)$$

Thus,  $|f(\mathbf{k}, \mathbf{k}')|^2$  gives the differential cross-section for particles with incident velocity  $\mathbf{v} = \hbar \mathbf{k}/m$  to be scattered such that their final velocities are directed into a range of solid angles  $d\Omega$  about  $\mathbf{v}' = \hbar \mathbf{k}'/m$ . Note that the scattering conserves energy, so that  $|\mathbf{v}'| = |\mathbf{v}|$  and  $|\mathbf{k}'| = |\mathbf{k}|$ .

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [14.1: Fundamentals of Scattering Theory](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 14.2: Born Approximation

Equation ([e15.17]) is not particularly useful, as it stands, because the quantity  $f(\mathbf{k}, \mathbf{k}')$  depends on the, as yet, unknown wavefunction  $\psi(\mathbf{r})$ . [See Equation ([e5.12]).] Suppose, however, that the scattering is not particularly strong. In this case, it is reasonable to suppose that the total wavefunction,  $\psi(\mathbf{r})$ , does not differ substantially from the incident wavefunction,  $\psi_0(\mathbf{r})$ . Thus, we can obtain an expression for  $f(\mathbf{k}, \mathbf{k}')$  by making the substitution  $\psi(\mathbf{r}) \rightarrow \psi_0(\mathbf{r}) = \sqrt{n} \exp(i \mathbf{k} \cdot \mathbf{r})$  in Equation ([e5.12]). This procedure is called the *Born approximation*.

The Born approximation yields

$$f(\mathbf{k}, \mathbf{k}') \simeq \frac{m}{2\pi \hbar^2} \int e^{i(\mathbf{k}-\mathbf{k}') \cdot \mathbf{r}'} V(\mathbf{r}') d^3 \mathbf{r}'. \quad (14.2.1)$$

Thus,  $f(\mathbf{k}, \mathbf{k}')$  becomes proportional to the Fourier transform of the scattering potential  $V(\mathbf{r})$  with respect to the wavevector  $\mathbf{q} = \mathbf{k} - \mathbf{k}'$ .

For a spherically symmetric potential,

$$f(\mathbf{k}', \mathbf{k}) \simeq -\frac{m}{2\pi \hbar^2} \int \int \int \exp(i q r' \cos \theta') V(r') r'^2 dr' \sin \theta' d\theta' d\phi', \quad (14.2.2)$$

giving

$$f(\mathbf{k}', \mathbf{k}) \simeq -\frac{2m}{\hbar^2 q} \int_0^\infty r' V(r') \sin(q r') dr'. \quad (14.2.3)$$

Note that  $f(\mathbf{k}', \mathbf{k})$  is just a function of  $q$  for a spherically symmetric potential. It is easily demonstrated that

$$q \equiv |\mathbf{k} - \mathbf{k}'| = 2k \sin(\theta/2), \quad (14.2.4)$$

where  $\theta$  is the angle subtended between the vectors  $\mathbf{k}$  and  $\mathbf{k}'$ . In other words,  $\theta$  is the scattering angle. Recall that the vectors  $\mathbf{k}$  and  $\mathbf{k}'$  have the same length, via energy conservation.

Consider scattering by a *Yukawa potential*,

$$V(r) = \frac{V_0 \exp(-\mu r)}{\mu r}, \quad (14.2.5)$$

where  $V_0$  is a constant, and  $1/\mu$  measures the “range” of the potential. It follows from Equation ([e17.38]) that

$$f(\theta) = -\frac{2m V_0}{\hbar^2 \mu} \frac{1}{q^2 + \mu^2}, \quad (14.2.6)$$

because

$$\int_0^\infty \exp(-\mu r') \sin(q r') dr' = \frac{q}{q^2 + \mu^2}. \quad (14.2.7)$$

Thus, in the Born approximation, the differential cross-section for scattering by a Yukawa potential is

$$\frac{d\sigma}{d\Omega} \simeq \left( \frac{2m V_0}{\hbar^2 \mu} \right)^2 \frac{1}{[2k^2 (1 - \cos \theta) + \mu^2]^2}, \quad (14.2.8)$$

given that

$$q^2 = 4k^2 \sin^2(\theta/2) = 2k^2 (1 - \cos \theta). \quad (14.2.9)$$

The Yukawa potential reduces to the familiar Coulomb potential as  $\mu \rightarrow 0$ , provided that  $V_0/\mu \rightarrow Z Z' e^2/(4\pi \epsilon_0)$ . In this limit, the Born differential cross-section becomes

$$\frac{d\sigma}{d\Omega} \simeq \left( \frac{2m Z Z' e^2}{4\pi \epsilon_0 \hbar^2} \right)^2 \frac{1}{16k^4 \sin^4(\theta/2)}. \quad (14.2.10)$$

Recall that  $\hbar k$  is equivalent to  $|\mathbf{p}|$ , so the previous equation can be rewritten

$$\frac{d\sigma}{d\Omega} \simeq \left( \frac{Z Z' e^2}{16\pi \epsilon_0 E} \right)^2 \frac{1}{\sin^4(\theta/2)}, \quad (14.2.11)$$

where  $E = p^2/2m$  is the kinetic energy of the incident particles. Of course, Equation ([e17.46]) is identical to the famous *Rutherford scattering cross-section* formula of classical physics.

The Born approximation is valid provided that  $\psi(\mathbf{r})$  is not too different from  $\psi_0(\mathbf{r})$  in the scattering region. It follows, from Equation ([e15.9]), that the condition for  $\psi(\mathbf{r}) \simeq \psi_0(\mathbf{r})$  in the vicinity of  $\mathbf{r} = \mathbf{0}$  is

$$\left| \frac{m}{2\pi \hbar^2} \int \frac{\exp(i k r')}{r'} V(\mathbf{r}') d^3 \mathbf{r}' \right| \ll 1. \quad (14.2.12)$$

Consider the special case of the Yukawa potential. At low energies, (i.e.,  $k \ll \mu$ ) we can replace  $\exp(i k r')$  by unity, giving

$$\frac{2m}{\hbar^2} \frac{|V_0|}{\mu^2} \ll 1 \quad (14.2.13)$$

as the condition for the validity of the Born approximation. The condition for the Yukawa potential to develop a bound state is

$$\frac{2m}{\hbar^2} \frac{|V_0|}{\mu^2} \geq 2.7, \quad (14.2.14)$$

where  $V_0$  is negative. Thus, if the potential is strong enough to form a bound state then the Born approximation is likely to break down. In the high- $k$  limit, Equation ([e17.47]) yields

$$\frac{2m}{\hbar^2} \frac{|V_0|}{\mu k} \ll 1. \quad (14.2.15)$$

This inequality becomes progressively easier to satisfy as  $k$  increases, implying that the Born approximation is more accurate at high incident particle energies.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [14.2: Born Approximation](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 14.3: Partial Waves

We can assume, without loss of generality, that the incident wavefunction is characterized by a wavevector  $\mathbf{k}$  that is aligned parallel to the  $z$ -axis. The scattered wavefunction is characterized by a wavevector  $\mathbf{k}'$  that has the same magnitude as  $\mathbf{k}$ , but, in general, points in a different direction. The direction of  $\mathbf{k}'$  is specified by the polar angle  $\theta$  (i.e., the angle subtended between the two wavevectors), and an azimuthal angle  $\phi$  about the  $z$ -axis. Equations ([e17.38]) and ([e17.39]) strongly suggest that for a spherically symmetric scattering potential [i.e.,  $V(\mathbf{r}) = V(r)$ ] the scattering amplitude is a function of  $\theta$  only: that is,

$$f(\theta, \phi) = f(\theta). \quad (14.3.1)$$

It follows that neither the incident wavefunction,

$$\psi_0(\mathbf{r}) = \sqrt{n} \exp(i k z) = \sqrt{n} \exp(i k r \cos \theta), \quad (14.3.2)$$

nor the large- $r$  form of the total wavefunction,

$$\psi(\mathbf{r}) = \sqrt{n} \left[ \exp(i k r \cos \theta) + \frac{\exp(i k r) f(\theta)}{r} \right], \quad (14.3.3)$$

depend on the azimuthal angle  $\phi$ .

Outside the range of the scattering potential, both  $\psi_0(\mathbf{r})$  and  $\psi(\mathbf{r})$  satisfy the free-space Schrödinger equation,

$$(\nabla^2 + k^2) \psi = 0. \quad (14.3.4)$$

What is the most general solution to this equation in spherical polar coordinates that does not depend on the azimuthal angle  $\phi$ ? Separation of variables yields

$$\psi(r, \theta) = \sum_l R_l(r) P_l(\cos \theta), \quad (14.3.5)$$

because the Legendre functions,  $P_l(\cos \theta)$ , form a complete set in  $\theta$ -space. The Legendre functions are related to the *spherical harmonics*, introduced in Chapter [sorb], via

$$P_l(\cos \theta) = \sqrt{\frac{4\pi}{2l+1}} Y_{l,0}(\theta, \varphi). \quad (14.3.6)$$

Equations ([e17.54]) and ([e17.55]) can be combined to give

$$r^2 \frac{d^2 R_l}{dr^2} + 2r \frac{dR_l}{dr} + [k^2 r^2 - l(l+1)] R_l = 0. \quad (14.3.7)$$

The two independent solutions to this equation are the spherical Bessel functions,  $j_l(kr)$  and  $y_l(kr)$ , introduced in Section [rwell]. Recall that

$$j_l(z) = z^l \left( -\frac{1}{z} \frac{d}{dz} \right)^l \left( \frac{\sin z}{z} \right),$$

$$y_l(z) = -z^l \left( -\frac{1}{z} \frac{d}{dz} \right)^l \left( \frac{\cos z}{z} \right).$$

Note that the  $j_l(z)$  are well behaved in the limit  $z \rightarrow 0$ , whereas the  $y_l(z)$  become singular. The asymptotic behavior of these functions in the limit  $z \rightarrow \infty$  is

$$j_l(z) \rightarrow \frac{\sin(z - l\pi/2)}{z},$$

$$y_l(z) \rightarrow -\frac{\cos(z - l\pi/2)}{z}.$$

We can write

$$\exp(i k r \cos \theta) = \sum_l a_l j_l(kr) P_l(\cos \theta), \quad (14.3.8)$$

where the  $a_l$  are constants. Note there are no  $y_l(kr)$  functions in this expression because they are not well-behaved as  $r \rightarrow 0$ . The Legendre functions are orthonormal ,

$$\int_{-1}^1 P_n(\mu) P_m(\mu) d\mu = \frac{\delta_{nm}}{n+1/2}, \quad (14.3.9)$$

so we can invert the previous expansion to give

$$a_l j_l(kr) = (l+1/2) \int_{-1}^1 \exp(i k r \mu) P_l(\mu) d\mu. \quad (14.3.10)$$

It is well known that

$$j_l(y) = \frac{(-i)^l}{2} \int_{-1}^1 \exp(i y \mu) P_l(\mu) d\mu, \quad (14.3.11)$$

where  $l = 0, 1, 2, \dots$ . Thus,

$$a_l = i^l (2l+1), \quad (14.3.12)$$

giving

$$\psi_0(\mathbf{r}) = \sqrt{n} \exp(i k r \cos \theta) = \sqrt{n} \sum_l i^l (2l+1) j_l(kr) P_l(\cos \theta). \quad (14.3.13)$$

The previous expression tells us how to decompose the incident plane-wave into a series of spherical waves. These waves are usually termed “partial waves”.

The most general expression for the total wavefunction outside the scattering region is

$$\psi(\mathbf{r}) = \sqrt{n} \sum_l [A_l j_l(kr) + B_l y_l(kr)] P_l(\cos \theta), \quad (14.3.14)$$

where the  $A_l$  and  $B_l$  are constants. Note that the  $y_l(kr)$  functions are allowed to appear in this expansion because its region of validity does not include the origin. In the large- $r$  limit, the total wavefunction reduces to

$$\psi(\mathbf{r}) \simeq \sqrt{n} \sum_l \left[ A_l \frac{\sin(kr - l\pi/2)}{kr} - B_l \frac{\cos(kr - l\pi/2)}{kr} \right] P_l(\cos \theta), \quad (14.3.15)$$

where use has been made of Equations ([e17.59a]) and ([e17.59b]). The previous expression can also be written

$$\psi(\mathbf{r}) \simeq \sqrt{n} \sum_l C_l \frac{\sin(kr - l\pi/2 + \delta_l)}{kr} P_l(\cos \theta), \quad (14.3.16)$$

where the sine and cosine functions have been combined to give a sine function which is phase-shifted by  $\delta_l$ . Note that  $A_l = C_l \cos \delta_l$  and  $B_l = -C_l \sin \delta_l$ .

Equation ([e17.68]) yields

$$\psi(\mathbf{r}) \simeq \sqrt{n} \sum_l C_l \left[ \frac{e^{i(kr - l\pi/2 + \delta_l)} - e^{-i(kr - l\pi/2 + \delta_l)}}{2i kr} \right] P_l(\cos \theta), \quad (14.3.17)$$

which contains both incoming and outgoing spherical waves. What is the source of the incoming waves? Obviously, they must be part of the large- $r$  asymptotic expansion of the incident wavefunction. In fact, it is easily seen from Equations ([e17.59a]) and ([e15.49]) that

$$\psi_0(\mathbf{r}) \simeq \sqrt{n} \sum_l i^l (2l+1) \left[ \frac{e^{i(kr - l\pi/2)} - e^{-i(kr - l\pi/2)}}{2i kr} \right] P_l(\cos \theta) \quad (14.3.18)$$

in the large- $r$  limit. Now, Equations ([e17.52]) and ([e17.53]) give

$$\frac{\psi(\mathbf{r}) - \psi_0(\mathbf{r})}{\sqrt{n}} = \frac{\exp(i kr)}{r} f(\theta). \quad (14.3.19)$$

Note that the right-hand side consists of an outgoing spherical wave only. This implies that the coefficients of the incoming spherical waves in the large- $r$  expansions of  $\psi(\mathbf{r})$  and  $\psi_0(\mathbf{r})$  must be the same. It follows from Equations ([e17.69]) and ([e17.70]) that

$$C_l = (2l + 1) \exp[i(\delta_l + l\pi/2)]. \quad (14.3.20)$$

Thus, Equations ([e17.69])–([e17.71]) yield

$$f(\theta) = \sum_{l=0, \infty} (2l + 1) \frac{\exp(i\delta_l)}{k} \sin \delta_l P_l(\cos \theta). \quad (14.3.21)$$

Clearly, determining the scattering amplitude,  $f(\theta)$ , via a decomposition into partial waves (i.e., spherical waves) is equivalent to determining the phase-shifts,  $\delta_l$ .

Now, the differential scattering cross-section,  $d\sigma/d\Omega$ , is simply the modulus squared of the scattering amplitude,  $f(\theta)$ . [See Equation ([e15.17]).] The total cross-section is thus given by

$$\begin{aligned} \sigma_{\text{total}} &= \int |f(\theta)|^2 d\Omega \\ &= \frac{1}{k^2} \oint d\phi \int_{-1}^1 d\mu \sum_l \sum_{l'} (2l + 1) (2l' + 1) \exp[i(\delta_l - \delta_{l'})] \sin \delta_l \sin \delta_{l'} P_l(\mu) P_{l'}(\mu), \end{aligned}$$

where  $\mu = \cos \theta$ . It follows that

$$\sigma_{\text{total}} = \frac{4\pi}{k^2} \sum_l (2l + 1) \sin^2 \delta_l, \quad (14.3.22)$$

where use has been made of Equation ([e17.61]).

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [14.3: Partial Waves](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).



## 14.4: Optical Theorem

---

A comparison between Equations ([\[e17.73\]](#)) and ([\[e17.75\]](#)) reveals that

$$\sigma_{\text{total}} = \frac{4\pi}{k} \text{Im}[f(0)] = \frac{4\pi}{k} \text{Im}(f(\mathbf{k}, \mathbf{k})), \quad (14.4.1)$$

because  $P_l(0) = 1$ . This result is known as the *optical theorem*, and is a consequence of the fact that the very existence of scattering requires scattering in the forward ( $\theta = 0$ ) direction, in order to interfere with the incident wave, and thereby reduce the probability current in that direction.

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [14.4: Optical Theorem](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 14.5: Determination of Phase-Shifts

Let us now consider how the phase-shifts,  $\delta_l$ , in Equation ([e17.73]) can be evaluated. Consider a spherically symmetric potential,  $V(r)$ , that vanishes for  $r > a$ , where  $a$  is termed the *range* of the potential. In the region  $r > a$ , the wavefunction  $\psi(\mathbf{r})$  satisfies the free-space Schrödinger equation ([e17.54]). The most general solution that is consistent with no incoming spherical-waves is

$$\psi(\mathbf{r}) = \sqrt{n} \sum_{l=0, \infty} i^l (2l+1) \mathcal{R}_l(r) P_l(\cos \theta), \quad (14.5.1)$$

where

$$\mathcal{R}_l(r) = \exp(i \delta_l) [\cos \delta_l j_l(kr) - \sin \delta_l y_l(kr)]. \quad (14.5.2)$$

Note that  $y_l(kr)$  functions are allowed to appear in the previous expression because its region of validity does not include the origin (where  $V \neq 0$ ). The logarithmic derivative of the  $l$ th radial wavefunction,  $\mathcal{R}_l(r)$ , just outside the range of the potential is given by

$$\beta_{l+} = k a \left[ \frac{\cos \delta_l j'_l(ka) - \sin \delta_l y'_l(ka)}{\cos \delta_l j_l(ka) - \sin \delta_l y_l(ka)} \right], \quad (14.5.3)$$

where  $j'_l(x)$  denotes  $dj_l(x)/dx$ , et cetera. The previous equation can be inverted to give

$$\tan \delta_l = \frac{k a j'_l(ka) - \beta_{l+} j_l(ka)}{k a y'_l(ka) - \beta_{l+} y_l(ka)}. \quad (14.5.4)$$

Thus, the problem of determining the phase-shift,  $\delta_l$ , is equivalent to that of obtaining  $\beta_{l+}$ .

The most general solution to Schrödinger's equation inside the range of the potential ( $r < a$ ) that does not depend on the azimuthal angle  $\phi$  is

$$\psi(\mathbf{r}) = \sqrt{n} \sum_{l=0, \infty} i^l (2l+1) \mathcal{R}_l(r) P_l(\cos \theta), \quad (14.5.5)$$

where

$$\mathcal{R}_l(r) = \frac{u_l(r)}{r}, \quad (14.5.6)$$

and

$$\frac{d^2 u_l}{dr^2} + \left[ k^2 - \frac{l(l+1)}{r^2} - \frac{2m}{\hbar^2} V \right] u_l = 0. \quad (14.5.7)$$

The boundary condition

$$u_l(0) = 0 \quad (14.5.8)$$

ensures that the radial wavefunction is well behaved at the origin. We can launch a well-behaved solution of the previous equation from  $r = 0$ , integrate out to  $r = a$ , and form the logarithmic derivative

$$\beta_{l-} = \frac{1}{(u_l/r)} \left. \frac{d(u_l/r)}{dr} \right|_{r=a}. \quad (14.5.9)$$

Because  $\psi(\mathbf{r})$  and its first derivatives are necessarily continuous for physically acceptable wavefunctions, it follows that

$$\beta_{l+} = \beta_{l-}. \quad (14.5.10)$$

The phase-shift,  $\delta_l$ , is then obtainable from Equation ([e17.82]).

### Contributors and Attributions

- Richard Fitzpatrick (Professor of Physics, The University of Texas at Austin)

This page titled [14.5: Determination of Phase-Shifts](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 14.6: Hard-Sphere Scattering

Let us test out this scheme using a particularly simple example. Consider scattering by a *hard sphere*, for which the potential is infinite for  $r < a$ , and zero for  $r > a$ . It follows that  $\psi(\mathbf{r})$  is zero in the region  $r < a$ , which implies that  $u_l = 0$  for all  $l$ . Thus,

$$\beta_{l-} = \beta_{l+} = \infty, \quad (14.6.1)$$

for all  $l$ . Equation ([e17.82]) thus gives

$$\tan \delta_l = \frac{j_l(ka)}{y_l(ka)}. \quad (14.6.2)$$

Consider the  $l = 0$  partial wave, which is usually referred to as the *S-wave*. Equation ([e17.90]) yields

$$\tan \delta_0 = \frac{\sin(ka)/ka}{-\cos(ka)/ka} = -\tan(ka), \quad (14.6.3)$$

where use has been made of Equations ([e17.58a]) and ([e17.58b]). It follows that

$$\delta_0 = -ka. \quad (14.6.4)$$

The *S-wave* radial wave function is [see Equation ([e17.80])]

$$\begin{aligned} \mathcal{R}_0(r) &= \exp(-ika) \frac{[\cos(ka) \sin(kr) - \sin(ka) \cos(kr)]}{kr} \\ &= \exp(-ika) \frac{\sin[k(r-a)]}{kr}. \end{aligned}$$

The corresponding radial wavefunction for the incident wave takes the form [see Equation ([e15.49])]

$$\tilde{\mathcal{R}}_0(r) = \frac{\sin(kr)}{kr}. \quad (14.6.5)$$

Thus, the actual  $l = 0$  radial wavefunction is similar to the incident  $l = 0$  wavefunction, except that it is phase-shifted by  $ka$ .

Let us examine the low- and high-energy asymptotic limits of  $\tan \delta_l$ . Low energy implies that  $ka \ll 1$ . In this regime, the spherical Bessel functions reduce to:

$$\begin{aligned} j_l(kr) &\simeq \frac{(kr)^l}{(2l+1)!!}, \\ y_l(kr) &\simeq -\frac{(2l-1)!!}{(kr)^{l+1}}, \end{aligned}$$

where  $n!! = n(n-2)(n-4) \cdots 1$ . It follows that

$$\tan \delta_l = \frac{-(ka)^{2l+1}}{(2l+1)[(2l-1)!!]^2}. \quad (14.6.6)$$

It is clear that we can neglect  $\delta_l$ , with  $l > 0$ , with respect to  $\delta_0$ . In other words, at low energy, only *S-wave* scattering (i.e., spherically symmetric scattering) is important. It follows from Equations ([e15.17]), ([e17.73]), and ([e17.92]) that

$$\frac{d\sigma}{d\Omega} = \frac{\sin^2 ka}{k^2} \simeq a^2 \quad (14.6.7)$$

for  $ka \ll 1$ . Note that the total cross-section

$$\sigma_{\text{total}} = \int \frac{d\sigma}{d\Omega} d\Omega = 4\pi a^2 \quad (14.6.8)$$

is four times the geometric cross-section  $\pi a^2$  (i.e., the cross-section for classical particles bouncing off a hard sphere of radius  $a$ ). However, low energy scattering implies relatively long wavelengths, so we would not expect to obtain the classical result in this limit.

Consider the high-energy limit  $ka \gg 1$ . At high energies, all partial waves up to  $l_{\max} = ka$  contribute significantly to the scattering cross-section. It follows from Equation (14.6.75) that

$$\sigma_{\text{total}} \simeq \frac{4\pi}{k^2} \sum_{l=0, l_{\max}} (2l+1) \sin^2 \delta_l. \quad (14.6.9)$$

With so many  $l$  values contributing, it is legitimate to replace  $\sin^2 \delta_l$  by its average value  $1/2$ . Thus,

$$\sigma_{\text{total}} \simeq \sum_{l=0, ka} \frac{2\pi}{k^2} (2l+1) \simeq 2\pi a^2. \quad (14.6.10)$$

This is twice the classical result, which is somewhat surprising, because we might expect to obtain the classical result in the short-wavelength limit. For hard-sphere scattering, incident waves with impact parameters less than  $a$  must be deflected. However, in order to produce a “shadow” behind the sphere, there must also be some scattering in the forward direction in order to produce destructive interference with the incident plane-wave. (Recall the optical theorem.) In fact, the interference is not completely destructive, and the shadow has a bright spot (the so-called “Poisson spot”) in the forward direction. The effective cross-section associated with this bright spot is  $\pi a^2$  which, when combined with the cross-section for classical reflection,  $\pi a^2$ , gives the actual cross-section of  $2\pi a^2$ .

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [14.6: Hard-Sphere Scattering](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 14.7: Low-Energy Scattering

In general, at low energies (i.e., when  $1/k$  is much larger than the range of the potential), partial waves with  $l > 0$  make a negligible contribution to the scattering cross-section. It follows that, at these energies, with a finite range potential, only  $S$ -wave scattering is important.

As a specific example, let us consider scattering by a finite potential well, characterized by  $V = V_0$  for  $r < a$ , and  $V = 0$  for  $r \geq a$ . Here,  $V_0$  is a constant. The potential is repulsive for  $V_0 > 0$ , and attractive for  $V_0 < 0$ . The outside wavefunction is given by [see Equation ([e17.80])]

$$\begin{aligned}\mathcal{R}_0(r) &= \exp(i\delta_0) [\cos\delta_0 j_0(kr) - \sin\delta_0 y_0(kr)] \\ &= \frac{\exp(i\delta_0) \sin(kr + \delta_0)}{kr},\end{aligned}$$

where use has been made of Equations ([e17.58a]) and ([e17.58b]). The inside wavefunction follows from Equation ([e17.85]). We obtain

$$\mathcal{R}_0(r) = B \frac{\sin(k'r)}{r}, \quad (14.7.1)$$

where use has been made of the boundary condition ([e17.86]). Here,  $B$  is a constant, and

$$E - V_0 = \frac{\hbar^2 k'^2}{2m}. \quad (14.7.2)$$

Note that Equation ([e17.103]) only applies when  $E > V_0$ . For  $E < V_0$ , we have

$$\mathcal{R}_0(r) = B \frac{\sinh(\kappa r)}{r}, \quad (14.7.3)$$

where

$$V_0 - E = \frac{\hbar^2 \kappa^2}{2m}. \quad (14.7.4)$$

Matching  $\mathcal{R}_0(r)$ , and its radial derivative, at  $r = a$  yields

$$\tan(ka + \delta_0) = \frac{k}{k'} \tan(k'a) \quad (14.7.5)$$

for  $E > V_0$ , and

$$\tan(ka + \delta_0) = \frac{k}{\kappa} \tanh(\kappa a) \quad (14.7.6)$$

for  $E < V_0$ .

Consider an attractive potential, for which  $E > V_0$ . Suppose that  $|V_0| \gg E$  (i.e., the depth of the potential well is much larger than the energy of the incident particles), so that  $k' \gg k$ . We can see from Equation ([e17.107]) that, unless  $\tan(k'a)$  becomes extremely large, the right-hand side is much less than unity, so replacing the tangent of a small quantity with the quantity itself, we obtain

$$ka + \delta_0 \simeq \frac{k}{k'} \tan(k'a). \quad (14.7.7)$$

This yields

$$\delta_0 \simeq ka \left[ \frac{\tan(k'a)}{k'a} - 1 \right]. \quad (14.7.8)$$

According to Equation ([e17.99]), the scattering cross-section is given by

$$\sigma_{\text{total}} \simeq \frac{4\pi}{k^2} \sin^2 \delta_0 = 4\pi a^2 \left[ \frac{\tan(k'a)}{k'a} - 1 \right]^2. \quad (14.7.9)$$

Now,

$$k' a = \sqrt{k^2 a^2 + \frac{2 m |V_0| a^2}{\hbar^2}}, \quad (14.7.10)$$

so for sufficiently small values of  $k a$ ,

$$k' a \simeq \sqrt{\frac{2 m |V_0| a^2}{\hbar^2}}. \quad (14.7.11)$$

It follows that the total ( $S$ -wave) scattering cross-section is independent of the energy of the incident particles (provided that this energy is sufficiently small).

Note that there are values of  $k' a$  (e.g.,  $k' a \simeq 4.49$ ) at which  $\delta_0 \rightarrow \pi$ , and the scattering cross-section ([\[e17.111\]](#)) vanishes, despite the very strong attraction of the potential. In reality, the cross-section is not exactly zero, because of contributions from  $l > 0$  partial waves. But, at low incident energies, these contributions are small. It follows that there are certain values of  $V_0$  and  $k$  that give rise to almost perfect transmission of the incident wave. This is called the *Ramsauer-Townsend effect*, and has been observed experimentally.

## Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [14.7: Low-Energy Scattering](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## 14.8: Resonances

There is a significant exception to the independence of the cross-section on energy mentioned previously. Suppose that the quantity  $(2m|V_0|a^2/\hbar^2)^{1/2}$  is slightly less than  $\pi/2$ . As the incident energy increases,  $k'a$ , which is given by Equation ([e17.112]), can reach the value  $\pi/2$ . In this case,  $\tan(k'a)$  becomes infinite, so we can no longer assume that the right-hand side of Equation ([e17.107]) is small. In fact, it follows from Equation ([e17.107]) that if the value of the incident energy is such that  $k'a = \pi/2$  then we also have  $ka + \delta_0 = \pi/2$ , or  $\delta_0 \simeq \pi/2$  (because we are assuming that  $ka \ll 1$ ). This implies that

$$\sigma_{\text{total}} = \frac{4\pi}{k^2} \sin^2 \delta_0 = 4\pi a^2 \left( \frac{1}{k^2 a^2} \right). \quad (14.8.1)$$

Note that the cross-section now depends on the energy. Furthermore, the magnitude of the cross-section is much larger than that given in Equation ([e17.111]) for  $k'a \neq \pi/2$  (because  $ka \ll 1$ ).

The origin of this rather strange behavior is quite simple. The condition

$$\sqrt{\frac{2m|V_0|a^2}{\hbar^2}} = \frac{\pi}{2} \quad (14.8.2)$$

is equivalent to the condition that a spherical well of depth  $V_0$  possesses a bound state at zero energy. Thus, for a potential well that satisfies the previous equation, the energy of the scattering system is essentially the same as the energy of the bound state. In this situation, an incident particle would like to form a bound state in the potential well. However, the bound state is not stable, because the system has a small positive energy. Nevertheless, this sort of *resonance scattering* is best understood as the capture of an incident particle to form a metastable bound state, and the subsequent decay of the bound state and release of the particle. The cross-section for resonance scattering is generally much larger than that for non-resonance scattering.

We have seen that there is a resonant effect when the phase-shift of the  $S$ -wave takes the value  $\pi/2$ . There is nothing special about the  $l = 0$  partial wave, so it is reasonable to assume that there is a similar resonance when the phase-shift of the  $l$ th partial wave is  $\pi/2$ . Suppose that  $\delta_l$  attains the value  $\pi/2$  at the incident energy  $E_0$ , so that

$$\delta_l(E_0) = \frac{\pi}{2}. \quad (14.8.3)$$

Let us expand  $\cot \delta_l$  in the vicinity of the resonant energy:

$$\begin{aligned} \cot \delta_l(E) &= \cot \delta_l(E_0) + \left( \frac{d \cot \delta_l}{dE} \right)_{E=E_0} (E - E_0) + \dots \\ &= - \left( \frac{1}{\sin^2 \delta_l} \frac{d\delta_l}{dE} \right)_{E=E_0} (E - E_0) + \dots \end{aligned}$$

Defining

$$\left( \frac{d\delta_l(E)}{dE} \right)_{E=E_0} = \frac{2}{\Gamma} \quad (14.8.4)$$

we obtain

$$\cot \delta_l(E) = -\frac{2}{\Gamma} (E - E_0) + \dots \quad (14.8.5)$$

Recall, from Equation ([e17.75]), that the contribution of the  $l$ th partial wave to the scattering cross-section is

$$\sigma_l = \frac{4\pi}{k^2} (2l+1) \sin^2 \delta_l = \frac{4\pi}{k^2} (2l+1) \frac{1}{1 + \cot^2 \delta_l}. \quad (14.8.6)$$

Thus,

$$\sigma_l \simeq \frac{4\pi}{k^2} (2l+1) \frac{\Gamma^2/4}{(E - E_0)^2 + \Gamma^2/4}. \quad (14.8.7)$$



This is the famous *Breit-Wigner formula* . The variation of the partial cross-section  $\sigma_l$  with the incident energy has the form of a classical resonance curve. The quantity  $\Gamma$  is the width of the resonance (in energy). We can interpret the Breit-Wigner formula as describing the absorption of an incident particle to form a metastable state, of energy  $E_0$ , and lifetime  $\tau = \hbar/\Gamma$ .

### Contributors and Attributions

- [Richard Fitzpatrick](#) (Professor of Physics, The University of Texas at Austin)

---

This page titled [14.8: Resonances](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

## CHAPTER OVERVIEW

### 15: Bibliography

#### Topic hierarchy

- [15.1: Section 1-](#)
- [15.2: Section 2-](#)
- [15.3: Section 3-](#)
- [15.4: Section 4-](#)
- [15.5: Section 5-](#)
- [15.6: Section 6-](#)

---

This page titled [15: Bibliography](#) is shared under a [not declared](#) license and was authored, remixed, and/or curated by [Richard Fitzpatrick](#).

Welcome to the Physics Library. This Living Library is a principal hub of the [LibreTexts project](#), which is a multi-institutional collaborative venture to develop the next generation of open-access texts to improve postsecondary education at all levels of higher learning. The LibreTexts approach is highly collaborative where an Open Access textbook environment is under constant revision by students, faculty, and outside experts to supplant conventional paper-based books.

Welcome to the Physics Library. This Living Library is a principal hub of the [LibreTexts project](#), which is a multi-institutional collaborative venture to develop the next generation of open-access texts to improve postsecondary education at all levels of higher learning. The LibreTexts approach is highly collaborative where an Open Access textbook environment is under constant revision by students, faculty, and outside experts to supplant conventional paper-based books.

Welcome to the Physics Library. This Living Library is a principal hub of the [LibreTexts project](#), which is a multi-institutional collaborative venture to develop the next generation of open-access texts to improve postsecondary education at all levels of higher learning. The LibreTexts approach is highly collaborative where an Open Access textbook environment is under constant revision by students, faculty, and outside experts to supplant conventional paper-based books.

Welcome to the Physics Library. This Living Library is a principal hub of the [LibreTexts project](#), which is a multi-institutional collaborative venture to develop the next generation of open-access texts to improve postsecondary education at all levels of higher learning. The LibreTexts approach is highly collaborative where an Open Access textbook environment is under constant revision by students, faculty, and outside experts to supplant conventional paper-based books.

Welcome to the Physics Library. This Living Library is a principal hub of the [LibreTexts project](#), which is a multi-institutional collaborative venture to develop the next generation of open-access texts to improve postsecondary education at all levels of higher learning. The LibreTexts approach is highly collaborative where an Open Access textbook environment is under constant revision by students, faculty, and outside experts to supplant conventional paper-based books.

Welcome to the Physics Library. This Living Library is a principal hub of the [LibreTexts project](#), which is a multi-institutional collaborative venture to develop the next generation of open-access texts to improve postsecondary education at all levels of higher learning. The LibreTexts approach is highly collaborative where an Open Access textbook environment is under constant revision by students, faculty, and outside experts to supplant conventional paper-based books.



## Index

### B

Born approximation  
14.2: Born Approximation  
bosons  
5.4: Identical Particles

### C

Central Potentials  
8: Central Potentials

### E

effective nuclear charge  
13.2: Helium Atom  
Ehrenfest's Theorem  
3.4: Ehrenfest's Theorem  
EPR  
12.3: Spin Magnetic Resonance  
expectation value  
3.3: Expectation Values (Averages) and Variances

### F

fermions  
5.4: Identical Particles  
Forbidden Transitions  
12.13: Forbidden Transitions

### G

gyromagnetic ratio  
12.3: Spin Magnetic Resonance

### H

Hydrogen atom  
8.3: Hydrogen Atom  
hyperfine structure  
11.10: Hyperfine Structure

### I

Infinite Spherical Potential Well  
8.2: Infinite Spherical Potential Well

### N

NMR  
12.3: Spin Magnetic Resonance  
normalization condition  
3.2: Normalization of the Wavefunction

### O

Operators  
3.5: Operators  
Optical Theorem  
14.4: Optical Theorem

### P

Partial Waves  
14.3: Partial Waves  
Pauli matrices  
9.4: Pauli Representation  
Pauli representation  
9.4: Pauli Representation  
Photoelectric effect  
2.5: Photoelectric Effect

### R

reduced mass  
5.3: Two-Particle Systems  
Rydberg formula  
8.4: Rydberg Formula

### S

Scattering  
14: Scattering Theory

### Selection rules

12.10: Selection Rules (Hydrogen Atoms)  
shielding

13.2: Helium Atom

Slater determinant

5.4: Identical Particles

Spherical Bessel Functions

8.2: Infinite Spherical Potential Well

Spherical harmonics

7.6: Spherical Harmonics

spin angular momentum

9: Spin Angular Momentum

spin operators

9.1: Spin Operators

Spin Precession

9.5: Spin Precession

Spin Space

9.2: Spin Space

spinor

9.4: Pauli Representation

Spontaneous Emission

12.8: Spontaneous Emission

### V

Variational Principle  
13.1: Variational Principle

### W

wavefunctions  
2.1: Wavefunctions

### Y

Yukawa potential  
14.2: Born Approximation

## Index

### B

Born approximation  
14.2: Born Approximation  
bosons  
5.4: Identical Particles

### C

Central Potentials  
8: Central Potentials

### E

effective nuclear charge  
13.2: Helium Atom  
Ehrenfest's Theorem  
3.4: Ehrenfest's Theorem  
EPR  
12.3: Spin Magnetic Resonance  
expectation value  
3.3: Expectation Values (Averages) and Variances

### F

fermions  
5.4: Identical Particles  
Forbidden Transitions  
12.13: Forbidden Transitions

### G

gyromagnetic ratio  
12.3: Spin Magnetic Resonance

### H

Hydrogen atom  
8.3: Hydrogen Atom  
hyperfine structure  
11.10: Hyperfine Structure

### I

Infinite Spherical Potential Well  
8.2: Infinite Spherical Potential Well

### N

NMR  
12.3: Spin Magnetic Resonance  
normalization condition  
3.2: Normalization of the Wavefunction

### O

Operators  
3.5: Operators  
Optical Theorem  
14.4: Optical Theorem

### P

Partial Waves  
14.3: Partial Waves  
Pauli matrices  
9.4: Pauli Representation  
Pauli representation  
9.4: Pauli Representation  
Photoelectric effect  
2.5: Photoelectric Effect

### R

reduced mass  
5.3: Two-Particle Systems  
Rydberg formula  
8.4: Rydberg Formula

### S

Scattering  
14: Scattering Theory

### Selection rules

12.10: Selection Rules (Hydrogen Atoms)  
shielding

13.2: Helium Atom  
Slater determinant

5.4: Identical Particles  
Spherical Bessel Functions  
8.2: Infinite Spherical Potential Well

Spherical harmonics  
7.6: Spherical Harmonics  
spin angular momentum  
9: Spin Angular Momentum

spin operators  
9.1: Spin Operators

Spin Precession  
9.5: Spin Precession

Spin Space  
9.2: Spin Space

spinor  
9.4: Pauli Representation  
Spontaneous Emission  
12.8: Spontaneous Emission

### V

Variational Principle  
13.1: Variational Principle

### W

wavefunctions  
2.1: Wavefunctions

### Y

Yukawa potential  
14.2: Born Approximation

## Glossary

---

**Sample Word 1** | Sample Definition 1

## Detailed Licensing

### Overview

**Title:** [Introductory Quantum Mechanics \(Fitzpatrick\)](#)

**Webpages:** 136

**All licenses found:**

- [Undeclared](#): 100% (136 pages)

### By Page

- [Introductory Quantum Mechanics \(Fitzpatrick\)](#) - *Undeclared*
  - [Front Matter](#) - *Undeclared*
    - [TitlePage](#) - *Undeclared*
    - [InfoPage](#) - *Undeclared*
    - [Table of Contents](#) - *Undeclared*
    - [Licensing](#) - *Undeclared*
  - [1: Probability Theory](#) - *Undeclared*
    - [1.1: What is Probability?](#) - *Undeclared*
    - [1.2: Combining Probabilities](#) - *Undeclared*
    - [1.3: Mean, Variance, and Standard Deviation](#) - *Undeclared*
    - [1.4: Continuous Probability Distributions](#) - *Undeclared*
    - [1.5: Exercises](#) - *Undeclared*
  - [2: Wave-Particle Duality](#) - *Undeclared*
    - [2.1: Wavefunctions](#) - *Undeclared*
    - [2.2: Plane-Waves](#) - *Undeclared*
    - [2.3: Representation of Waves via Complex Functions](#) - *Undeclared*
    - [2.4: Classical Light-Waves](#) - *Undeclared*
    - [2.5: Photoelectric Effect](#) - *Undeclared*
    - [2.6: Quantum Theory of Light](#) - *Undeclared*
    - [2.7: Classical Interferences of Light Waves](#) - *Undeclared*
    - [2.8: Quantum Interference of Light](#) - *Undeclared*
    - [2.9: Particles](#) - *Undeclared*
    - [2.10: Wave-Packets](#) - *Undeclared*
    - [2.11: Evolution of Wave-Packets](#) - *Undeclared*
    - [2.12: Schrodinger's Equation and Wavefunction Collapse](#) - *Undeclared*
    - [2.13: Exercises](#) - *Undeclared*
  - [3: Fundamentals of Quantum Mechanics](#) - *Undeclared*
    - [3.1: Schrodinger's Equation](#) - *Undeclared*
    - [3.2: Normalization of the Wavefunction](#) - *Undeclared*
    - [3.3: Expectation Values \(Averages\) and Variances](#) - *Undeclared*
    - [3.4: Ehrenfest's Theorem](#) - *Undeclared*
    - [3.5: Operators](#) - *Undeclared*
    - [3.6: Momentum Representation](#) - *Undeclared*
    - [3.7: Heisenberg's Uncertainty Principle](#) - *Undeclared*
    - [3.8: Eigenstates and Eigenvalues](#) - *Undeclared*
    - [3.9: Measurement](#) - *Undeclared*
    - [3.10: Stationary States](#) - *Undeclared*
    - [3.11: Exercises](#) - *Undeclared*
  - [4: One-Dimensional Potentials](#) - *Undeclared*
    - [4.1: Infinite Potential Well](#) - *Undeclared*
    - [4.2: Square Potential Barrier](#) - *Undeclared*
    - [4.3: WKB Approximation](#) - *Undeclared*
    - [4.4: Cold Emission](#) - *Undeclared*
    - [4.5: Alpha Decay](#) - *Undeclared*
    - [4.6: Square Potential Well](#) - *Undeclared*
    - [4.7: Simple Harmonic Oscillator](#) - *Undeclared*
    - [4.E: One-Dimensional Potentials \(Exercises\)](#) - *Undeclared*
  - [5: Multi-Particle Systems](#) - *Undeclared*
    - [5.1: Fundamental Concepts of Multi-Particle Systems](#) - *Undeclared*
    - [5.2: Non-interacting Particles](#) - *Undeclared*
    - [5.3: Two-Particle Systems](#) - *Undeclared*
    - [5.4: Identical Particles](#) - *Undeclared*
    - [5.E: Multi-Particle Systems \(Exercises\)](#) - *Undeclared*
  - [6: Three-Dimensional Quantum Mechanics](#) - *Undeclared*
    - [6.1: Fundamental Concepts](#) - *Undeclared*
    - [6.2: Particle in Box](#) - *Undeclared*
    - [6.3: Degenerate Electron Gases](#) - *Undeclared*
    - [6.4: White Dwarf Stars](#) - *Undeclared*
    - [6.5: Exercises](#) - *Undeclared*
  - [7: Orbital Angular Momentum](#) - *Undeclared*
    - [7.1: Angular Momentum Operators](#) - *Undeclared*
    - [7.2: Representation of Angular Momentum](#) - *Undeclared*
    - [7.3: Eigenstates of Angular Momentum](#) - *Undeclared*
    - [7.4: Eigenvalues of  \$L\_z\$](#)  - *Undeclared*
    - [7.5: Eigenvalues of  \$L^2\$](#)  - *Undeclared*
    - [7.6: Spherical Harmonics](#) - *Undeclared*
    - [7.E: Orbital Angular Momentum \(Exercises\)](#) - *Undeclared*
  - [8: Central Potentials](#) - *Undeclared*

- 8.1: Derivation of Radial Equation - *Undeclared*
- 8.2: Infinite Spherical Potential Well - *Undeclared*
- 8.3: Hydrogen Atom - *Undeclared*
- 8.4: Rydberg Formula - *Undeclared*
- 8.E: Central Potentials (Exercises) - *Undeclared*
- 9: Spin Angular Momentum - *Undeclared*
  - 9.1: Spin Operators - *Undeclared*
  - 9.2: Spin Space - *Undeclared*
  - 9.3: Eigenstates of  $S_z$  and  $S^2$  - *Undeclared*
  - 9.4: Pauli Representation - *Undeclared*
  - 9.5: Spin Precession - *Undeclared*
  - 9.E: Spin Angular Momentum (Exercises) - *Undeclared*
- 10: Addition of Angular Momentum - *Undeclared*
  - 10.1: General Principles of Angular Momentum - *Undeclared*
  - 10.2: Angular Momentum in Hydrogen Atom - *Undeclared*
  - 10.3: Two Spin One-Half Particles - *Undeclared*
  - 10.E: Addition of Angular Momentum (Exercises) - *Undeclared*
- 11: Time-Independent Perturbation Theory - *Undeclared*
  - 11.1: Exercises - *Undeclared*
  - 11.2: Improved Notation - *Undeclared*
  - 11.3: Two-State System - *Undeclared*
  - 11.4: Non-Degenerate Perturbation Theory - *Undeclared*
  - 11.5: Quadratic Stark Effect - *Undeclared*
  - 11.6: Degenerate Perturbation Theory - *Undeclared*
  - 11.7: Linear Stark Effect - *Undeclared*
  - 11.8: Fine Structure of Hydrogen - *Undeclared*
  - 11.9: Zeeman Effect - *Undeclared*
  - 11.10: Hyperfine Structure - *Undeclared*
- 12: Time-Dependent Perturbation Theory - *Undeclared*
  - 12.1: Preliminary Analysis - *Undeclared*
  - 12.2: Two-State System - *Undeclared*
  - 12.3: Spin Magnetic Resonance - *Undeclared*
  - 12.4: Perturbation Expansion - *Undeclared*
  - 12.5: Harmonic Perturbation - *Undeclared*
  - 12.6: Electromagnetic Radiation - *Undeclared*
  - 12.7: Electric Dipole Approximation - *Undeclared*
  - 12.8: Spontaneous Emission - *Undeclared*
  - 12.9: Radiation from Harmonic Oscillator - *Undeclared*
  - 12.10: Selection Rules (Hydrogen Atoms) - *Undeclared*
  - 12.11: 2P-1S Transitions in Hydrogen - *Undeclared*
  - 12.12: Intensity Rules - *Undeclared*
  - 12.13: Forbidden Transitions - *Undeclared*
  - 12.E: Time-Dependent Perturbation Theory (Exercises) - *Undeclared*
- 13: Variational Methods - *Undeclared*
  - 13.1: Variational Principle - *Undeclared*
  - 13.2: Helium Atom - *Undeclared*
  - 13.3: Hydrogen Molecule Ion - *Undeclared*
- 14: Scattering Theory - *Undeclared*
  - 14.1: Fundamentals of Scattering Theory - *Undeclared*
  - 14.2: Born Approximation - *Undeclared*
  - 14.3: Partial Waves - *Undeclared*
  - 14.4: Optical Theorem - *Undeclared*
  - 14.5: Determination of Phase-Shifts - *Undeclared*
  - 14.6: Hard-Sphere Scattering - *Undeclared*
  - 14.7: Low-Energy Scattering - *Undeclared*
  - 14.8: Resonances - *Undeclared*
- 15: Bibliography - *Undeclared*
  - 15.1: Section 1- - *Undeclared*
  - 15.2: Section 2- - *Undeclared*
  - 15.3: Section 3- - *Undeclared*
  - 15.4: Section 4- - *Undeclared*
  - 15.5: Section 5- - *Undeclared*
  - 15.6: Section 6- - *Undeclared*
- Back Matter - *Undeclared*
  - Index - *Undeclared*
  - Index - *Undeclared*
  - Glossary - *Undeclared*
  - Detailed Licensing - *Undeclared*