

6.1: Projectile Motion

The simplest case of two-dimensional motion occurs when a particle experiences a force only in one direction. The prime example of this case is the motion of a projectile in Earth's (or any other planet's) gravitational field as locally described by Galilean gravity (Equation 2.2.2): $\mathbf{F} = m\mathbf{g}$. Once a projectile has been fired with a certain initial velocity \mathbf{v}_0 , we can find its trajectory by solving the equation of motion that follows from Newton's second law: $m\mathbf{g} = m\ddot{\mathbf{r}}$. We can decompose \mathbf{r} and \mathbf{v}_0 in horizontal (x) and vertical (z) components; each of them has its own one-dimensional equation of motion, which we already solved in Section 2.3. The horizontal component experiences no force and thus executes a simple linear motion with uniform velocity $v_0 \cos \theta_0$, where $\theta_0 = \arccos(\mathbf{v}_0 \cdot \hat{\mathbf{x}})/v_0$ is the angle with the horizontal under which the projectile was fired and $v_0 = |\mathbf{v}_0|$ the initial speed. Likewise, because the acceleration due to gravitation is constant, our projectile will execute a uniformly accelerated motion in the vertical direction with initial velocity $v_0 \sin \theta_0$. If the projectile's initial position is (x_0, z_0) , its motion is thus described by:

$$\mathbf{r}(t) = \begin{pmatrix} x(t) \\ z(t) \end{pmatrix} = \begin{pmatrix} x_0 \\ z_0 \end{pmatrix} + v_0 \begin{pmatrix} \cos \theta_0 \\ \sin \theta_0 \end{pmatrix} t - \begin{pmatrix} 0 \\ g \end{pmatrix} \frac{1}{2} t^2 \quad (6.1.1)$$

This page titled [6.1: Projectile Motion](#) is shared under a [CC BY-NC-SA 4.0](#) license and was authored, remixed, and/or curated by [Timon Idema \(TU Delft Open\)](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.