

CHAPTER OVERVIEW

4: Motion in Two and Three Dimensions

To give a complete description of kinematics, we must explore motion in two and three dimensions. After all, most objects in our universe do not move in straight lines; rather, they follow curved paths. From kicked footballs to the flight paths of birds to the orbital motions of celestial bodies and down to the flow of blood plasma in your veins, most motion follows curved trajectories. In this chapter we also explore two special types of motion in two dimensions: projectile motion and circular motion. Last, we conclude with a discussion of relative motion. In the chapter-opening picture, each jet has a relative motion with respect to any other jet in the group or to the people observing the air show on the ground.

[4.1: Prelude to Motion in Two and Three Dimensions](#)

[4.2: Displacement and Velocity Vectors](#)

[4.3: Acceleration Vector](#)

[4.4: Projectile Motion](#)

[4.5: Uniform Circular Motion](#)

[4.6: Relative Motion in One and Two Dimensions](#)

[4.E: Motion in Two and Three Dimensions \(Exercises\)](#)

[4.S: Motion in Two and Three Dimensions \(Summary\)](#)

Thumbnail: The Red Arrows is the aerobatics display team of Britain's Royal Air Force. Based in Lincolnshire, England, they perform precision flying shows at high speeds, which requires accurate measurement of position, velocity, and acceleration in three dimensions. (credit: modification of work by Phil Long).

Contributors and Attributions

- Samuel J. Ling (Truman State University), Jeff Sanny (Loyola Marymount University), and Bill Moebs with many contributing authors. This work is licensed by OpenStax University Physics under a [Creative Commons Attribution License \(by 4.0\)](#).

This page titled [4: Motion in Two and Three Dimensions](#) is shared under a [CC BY](#) license and was authored, remixed, and/or curated by [OpenStax](#).