

CHAPTER OVERVIEW

3: Integrals

If we have a function $f(x)$ which is well-defined for some $a \leq x \leq b$, its integral over those two values is defined as

$$\int_a^b dx f(x) = \lim_{N \rightarrow \infty} \sum_{n=0}^N \Delta x f(x_n) \quad \text{where } x_n = a + n\Delta x, \quad \Delta x \equiv \left(\frac{b-a}{N} \right). \quad (3.1)$$

This is called a **definite integral**, and represents the area under the graph of $f(x)$ in the region between $x = a$ and $x = b$, as shown in the figure below. The function $f(x)$ is called the **integrand**, and the two points a and b are called the **bounds** of the integral. The interval between the two bounds is divided into N segments, of length $(b-a)/N$ each. Each term in the sum represents the area of a rectangle, and as $N \rightarrow \infty$, the sum converges to the area under the curve.




Figure 3.1

A **multiple integral** involves integration over more than one variable. For instance, when we have a function $f(x_1, x_2)$ that depends on two independent variables, x_1 and x_2 , we can perform a double integral by integrating over one variable first, then the other variable:

$$\int_{a_1}^{b_1} dx_1 \int_{a_2}^{b_2} dx_2 f(x_1, x_2) \equiv \int_{a_1}^{b_1} dx_1 F(x_1) \quad \text{where } F(x_1) \equiv \int_{a_2}^{b_2} dx_2 f(x_1, x_2). \quad (3.2)$$

[3.1: Basic Properties of Definite Integrals](#)

[3.2: Integrals as Antiderivatives](#)

[3.3: Integration by Parts](#)

[3.4: Change of Variables](#)

[3.5: The Gaussian Integral](#)

[3.6: Differentiating Under the Integral Sign](#)

[3.7: Exercises](#)

This page titled [3: Integrals](#) is shared under a [CC BY-SA 4.0](#) license and was authored, remixed, and/or curated by [Y. D. Chong](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.