

## CHAPTER OVERVIEW

### 4: Diffraction

In the preceding chapter, we implicitly regarded slits as objects with positions but no size. The widths of the slits were considered negligible. When the slits have finite widths, each point along the opening can be considered a point source of light—a foundation of Huygens's principle. Because real-world optical instruments must have finite apertures (otherwise, no light can enter), diffraction plays a major role in the way we interpret the output of these optical instruments. For example, diffraction places limits on our ability to resolve images or objects. This is a problem that we will study later in this chapter.

[4.1: Prelude to Diffraction](#)

[4.2: Single-Slit Diffraction](#)

[4.3: Intensity in Single-Slit Diffraction](#)

[4.4: Double-Slit Diffraction](#)

[4.5: Diffraction Gratings](#)

[4.6: Circular Apertures and Resolution](#)

[4.7: X-Ray Diffraction](#)

[4.8: Holography](#)

[4.A: Diffraction \(Answers\)](#)

[4.E: Diffraction \(Exercises\)](#)

[4.S: Diffraction \(Summary\)](#)

---

This page titled [4: Diffraction](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [OpenStax](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.