

4.2: Linear Momentum and Force

Learning Objectives

By the end of this section, you will be able to:

- Define linear momentum.
- Explain the relationship between momentum and force.
- State Newton's second law of motion in terms of momentum.
- Calculate momentum given mass and velocity.

The scientific definition of linear momentum is consistent with most people's intuitive understanding of momentum: a large, fast-moving object has greater momentum than a smaller, slower object. Linear momentum is defined as the product of a system's mass multiplied by its velocity.

Linear Momentum

Linear momentum is defined as the product of a system's mass multiplied by its velocity:

$$p = mv \quad (4.2.1)$$

Momentum is directly proportional to the object's mass and also its velocity. Thus the greater an object's mass or the greater its velocity, the greater its momentum. Momentum p is a vector having the same direction as the velocity v . The SI unit for momentum is $\text{kg} \cdot \text{m/s}$.

Example 4.2.1: Calculating Momentum: A Football Player and a Football

- Calculate the momentum of a 110-kg football player running at 8.00 m/s.
- Compare the player's momentum with the momentum of a hard-thrown 0.410-kg football that has a speed of 25.0 m/s.

Strategy

No information is given regarding direction, and so we can calculate only the magnitude of the momentum, p (As usual, a symbol that is in italics is a magnitude, whereas one that is italicized, boldfaced, and has an arrow is a vector.) In both parts of this example, the magnitude of momentum can be calculated directly from the definition of momentum given in Equation 4.2.1, which becomes

$$p = mv$$

when only magnitudes are considered.

Solution for (a)

To determine the momentum of the player, substitute the known values for the player's mass and speed into the equation.

$$\begin{aligned} p_{\text{player}} &= (110 \text{ kg})(8.00 \text{ m/s}) \\ &= 880 \text{ kg} \cdot \text{m/s} \end{aligned}$$

Solution for (b)

To determine the momentum of the ball, substitute the known values for the ball's mass and speed into the equation.

$$\begin{aligned} p_{\text{ball}} &= (0.410 \text{ kg})(25.0 \text{ m/s}) \\ &= 10.3 \text{ kg} \cdot \text{m/s} \end{aligned}$$

The ratio of the player's momentum to that of the ball is

$$\frac{p_{\text{player}}}{p_{\text{ball}}} = \frac{880}{10.3} = 85.0$$

Discussion

Although the ball has greater velocity, the player has a much greater mass. Thus the momentum of the player is much greater than the momentum of the football, as you might guess. As a result, the player's motion is only slightly affected if he catches the ball. We shall quantify what happens in such collisions in terms of momentum in later sections.

Momentum and Newton's Second Law

The importance of momentum, unlike the importance of energy, was recognized early in the development of classical physics. Momentum was deemed so important that it was called the "quantity of motion." Newton actually stated his second law of motion in terms of momentum: The net external force equals the change in momentum of a system divided by the time over which it changes.

Newton's Second Law of Motion in Terms of Momentum

The net external force equals the change in momentum of a system divided by the time over which it changes.

$$F_{net} = \frac{\Delta p}{\Delta t} \quad (4.2.2)$$

where F_{net} is the net external force, Δp is the change in momentum, and Δt is the change in time.

Making Connections: Force and Momentum

Force and momentum are intimately related. Force acting over time can change momentum, and Newton's second law of motion, can be stated in its most broadly applicable form in terms of momentum. Momentum continues to be a key concept in the study of atomic and subatomic particles in quantum mechanics.

This statement of Newton's second law of motion includes the more familiar $F_{net} = ma$ as a special case. We can derive this form as follows. First, note that the change in momentum Δp is given by

$$\Delta p = \Delta(mv) \quad (4.2.3)$$

If the mass of the system is constant, then

$$\Delta(mv) = m\Delta v. \quad (4.2.4)$$

So that for constant mass, Newton's second law of motion becomes

$$F_{net} = \frac{\Delta p}{\Delta t} = \frac{m\Delta v}{\Delta t}. \quad (4.2.5)$$

Because $\frac{\Delta v}{\Delta t} = a$, we get the familiar equation

$$F_{net} = ma \quad (4.2.6)$$

when the mass of the system is *constant*.

Newton's second law of motion stated in terms of momentum is more generally applicable because it can be applied to systems where the mass is changing, such as rockets, as well as to systems of constant mass. We will consider systems with varying mass in some detail; however, the relationship between momentum and force remains useful when mass is constant, such as in the following example.

Example 4.2.2: Calculating Force: Venus Williams' Racquet

During the 2007 French Open, Venus Williams hit the fastest recorded serve in a premier women's match, reaching a speed of 58 m/s (209 km/h). What is the average force exerted on the 0.057-kg tennis ball by Venus Williams' racquet, assuming that the ball's speed just after impact is 58 m/s, that the initial horizontal component of the velocity before impact is negligible, and that the ball remained in contact with the racquet for 5.0 ms (milliseconds)?

Strategy

This problem involves only one dimension because the ball starts from having no horizontal velocity component before impact. Newton's second law stated in terms of momentum is then written as

$$F_{net} = \frac{\Delta p}{\Delta t}$$

As noted above, when mass is constant, the change in momentum is given by

$$\Delta p = m\Delta v = m(v_f - v_i).$$

In this example, the velocity just after impact and the change in time are given; thus, once Δp is calculated, $F_{net} = \frac{\Delta p}{\Delta t}$ can be used to find the force.

Solution

To determine the change in momentum, substitute the values for the initial and final velocities into the equation above.

$$\begin{aligned}\Delta p &= m(v_f - v_i) \\ &= (0.057 \text{ kg})(58 \text{ m/s} - 0 \text{ m/s}) \\ &= 3.306 \text{ kg} \cdot \text{m/s} = 3.3 \text{ kg} \cdot \text{m/s}\end{aligned}$$

Now the magnitude of the net external force can be determined by using $F_{net} = \frac{\Delta p}{\Delta t}$

$$\begin{aligned}F_{net} &= \frac{\Delta p}{\Delta t} = \frac{3.306 \text{ kg}}{5.0 \times 10^{-3}} \\ &= 661 \text{ N},\end{aligned}$$

where we have retained only two significant figures in the final step.

Discussion

This quantity was the average force exerted by Venus Williams' racquet on the tennis ball during its brief impact (note that the ball also experienced the 0.56-N force of gravity, but that force was not due to the racquet). This problem could also be solved by first finding the acceleration and then using $F = ma$ but one additional step would be required compared with the strategy used in this example.

Summary

- Linear momentum (*momentum* for brevity) is defined as the product of a system's mass multiplied by its velocity.
- In symbols, linear momentum p is defined to be

$$p = mv$$

where m is the mass of the system and v is its velocity.

- The SI unit for momentum is $\text{kg} \cdot \text{m/s}$.
- Newton's second law of motion in terms of momentum states that the net external force equals the change in momentum of a system divided by the time over which it changes.
- In symbols, Newton's second law of motion is defined to be

$$F_{net} = \frac{\Delta p}{\Delta t}$$

where F_{net} is the net external force, Δp is the change in momentum, and Δt is change in time.

Glossary

linear momentum

the product of mass and velocity

second law of motion

physical law that states that the net external force equals the change in momentum of a system divided by the time over which it changes

This page titled [4.2: Linear Momentum and Force](#) is shared under a [CC BY](#) license and was authored, remixed, and/or curated by [OpenStax](#).

- **8.1: Linear Momentum and Force** by OpenStax is licensed [CC BY 4.0](#). Original source: <https://openstax.org/details/books/college-physics>.