

CHAPTER OVERVIEW

5: The Central Limit Theorem

In this chapter, you will study means and the **central limit theorem**, which is one of the most powerful and useful ideas in all of statistics. There are two alternative forms of the theorem, and both alternatives are concerned with drawing finite samples size n from a population with a known mean, μ , and a known standard deviation, σ . The first alternative says that if we collect samples of size n with a "large enough n ," calculate each sample's mean, and create a histogram of those means, then the resulting histogram will tend to have an approximate normal bell shape. The second alternative says that if we again collect samples of size n that are "large enough," calculate the sum of each sample and create a histogram, then the resulting histogram will again tend to have a normal bell-shape.

[5.1: Prelude to the Central Limit Theorem](#)

[5.2: The Central Limit Theorem for Sample Means \(Averages\)](#)

[5.2E: The Central Limit Theorem for Sample Means \(Exercises\)](#)

[5.3: The Central Limit Theorem for Sums](#)

[5.4: Using the Central Limit Theorem](#)

[5.4E: Using the Central Limit Theorem \(Exercises\)](#)

[5.5: Central Limit Theorem - Pocket Change \(Worksheet\)](#)

[5.6: Central Limit Theorem - Cookie Recipes \(Worksheet\)](#)

[5.E: The Central Limit Theorem \(Exercises\)](#)

Barbara Illowsky and Susan Dean (De Anza College) with many other contributing authors. Content produced by OpenStax College is licensed under a Creative Commons Attribution License 4.0 license. Download for free at <http://cnx.org/contents/30189442-699...b91b9de@18.114>.

This page titled [5: The Central Limit Theorem](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [OpenStax](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.