

CHAPTER OVERVIEW

6: Hypothesis Testing

One job of a statistician is to make statistical inferences about populations based on samples taken from the population. Confidence intervals are one way to estimate a population parameter. Another way to make a statistical inference is to make a decision about a parameter. For instance, a car dealer advertises that its new small truck gets 35 miles per gallon, on average. A tutoring service claims that its method of tutoring helps 90% of its students get an A or a B. A company says that women managers in their company earn an average of \$60,000 per year.

- [6.1: Prelude to Hypothesis Testing](#)
- [6.2: Null and Alternative Hypotheses](#)
- [6.3: Outcomes and the Type I and Type II Errors](#)
- [6.4: Distribution Needed for Hypothesis Testing](#)
- [6.5: Rare Events, the Sample, Decision and Conclusion](#)
- [6.6: Additional Information and Full Hypothesis Test Examples](#)
- [6.7: Prelude to Hypothesis Testing with Two Samples](#)
- [6.8: Two Population Means with Unknown Standard Deviations](#)
- [6.9: Two Population Means with Known Standard Deviations](#)
- [6.10: Comparing Two Independent Population Proportions](#)
- [6.11: Matched or Paired Samples](#)

This page titled [6: Hypothesis Testing](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [OpenStax](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.