

CHAPTER OVERVIEW

6: Hypothesis Testing with One Sample

One job of a statistician is to make statistical inferences about populations based on samples taken from the population. Confidence intervals are one way to estimate a population parameter. Another way to make a statistical inference is to make a decision about a parameter. For instance, a car dealer advertises that its new small truck gets 35 miles per gallon, on average. A tutoring service claims that its method of tutoring helps 90% of its students get an A or a B. A company says that women managers in their company earn an average of \$60,000 per year.

[6.1: Prelude to Hypothesis Testing](#)

[6.2: Null and Alternative Hypotheses](#)

[6.3: Outcomes and the Type I and Type II Errors](#)

[6.4: Distribution Needed for Hypothesis Testing](#)

[6.5: Rare Events, the Sample, Decision and Conclusion](#)

[6.6: Additional Information and Full Hypothesis Test Examples](#)

Contributors

- Barbara Illowsky and Susan Dean (De Anza College) with many other contributing authors. Content produced by OpenStax College is licensed under a Creative Commons Attribution License 4.0 license. Download for free at <http://cnx.org/contents/30189442-699...b91b9de@18.114>.

This page titled [6: Hypothesis Testing with One Sample](#) is shared under a [CC BY 4.0](#) license and was authored, remixed, and/or curated by [OpenStax](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.